- Anderson, S.C. & Leviton, A.E. Amphibians and reptiles collected by the Street expedition to Afghanistan, 1965. Proc. California Academy of Sciences. 4th series 37(2): 25-56.
- Annandale, N. & Carter, H.E. Notes on the vegetation of Smistan. Journ. As. Soc. Bengal. 1919 n.s. 15: 267-297.
- Arbeitegemeinschaft Afghanistan. Bibliographie der Afghanistan Literatur 1947-1967.

 1968 Hamburg. 189 p. (Bibliography of the Afghan literature: 1945-1967).
- Archibald, C. Reestablishing Siberian Cranes (Crus leucogeranus) as winter residents in 1975 Iran. Unpubl. research proposal submitted to the Iran Department of Environment, Division of Nature Conservation, Tehran.
- Arez, G.J. The climate of Afghanistan. Inst. of Geog., Fac. of Letters, Kabul University. 1969. 96 p.
- Aslanow, N.G. On the distribution of some Indian birds in Afghanistan. Zool. Nachr. 1953 Akad. Wiss. Usbek URSS (in Russian).
- Atchison, J.E. Consultants, Inc. A preliminary study to determine the prospects for establishing a paper manufacturing industry in Afghanistan. New York 2 Vols.
- Ayle, J.S. A mountain campaign on the borders of Afghanistan in 1863. Bentley, London. 1867 101 p.
- Baker, E.C.S. Notes on two collections of birds from Seistan. Rec. Indian Mus. 18(2): 1919 121-134.
- Balland, V. & Balland, D. Le géographie de l'Afghanistan, données nouvelles et transforma-1972 tions économiques récentes. Information G. (Paris) 2: 73-83, 3: 113-121. (The geography of Afghanistan, with new data and recent economic transformations.)
- Balland, V. & Leng, J. Les rapports géomorphologiques quaternaires et actuels du Bessin 1974 de Bamyan et de ses bordures montagneuses (Afghanistan central). Rev. G. Phys. et Géol. Dynamique (Paris), 2me sér. 16(3): 327-350.
- Bannikov, A.G., Darevsky, I.S. and Rustamov, A.K. Amphibians and reptiles of the USSR.

 1971 Moscow (in Russian).
- Barnes, H.E. Notes on the midification of certain species in the neighbourhood of Chaman, 1880 south Afghanistan. Stray Feathers. 9: 212-220.
- Barnes, H.E. A list of the birds observed in the neighbourhood of Chaman, south Afghanistan.

 1881 Stray Feathers. 9: 449-460.
- Barrat, C. Quelques traits caractéristiques de l'Afghanistan. Ann. G. (Paris) 81:444: 1972 206-229.
- Barth, F. Nomadism in the mountain and plateau areas of South West Asia. Arid Zone 1967 Research. 19

- Barth, F. Socio-economic changes and social problems in pastoral lands: Some concrete factors. Background paper no. 6 in: ecological guidelines for the use of natural resources in the Middle East and South West Asia. IUCN, Morges.
- Barus, V. & Tenora, F. Three interesting nematode species in Miniopterus schreibersi

 (Chiroptera) from Afghanistan. (Acta Univ. Agric. Brnc.) Sb. Uys. Sk. zeměd Brnc.

 (1): 95-101.
- Bellow, H.W. The races of Afghanistan. Thacker, Spink & Co. Caloutta. pp. 162.
- Béné, R. & Baldo, S. Etude microbiologique des sols halomorphes afghans. Bull. Assoc.

 des Diplomés de Microbiology Fac. pharm. Nancy. 98: 3-7. (A microbiological study of the Afghan halomorphic soils).
- Béranger Beauguesne, L. Le bois à travers le monde: Afghanistan. Revue int. Bois. 1956 23: 217-218, 139-140.
- Beveridge, A.S. Babur-Nama (memoirs of Babur). Translated from the Turki text of 1970 Zahiru'd-din Muhammad Babur Padshah Gazi. New Delhi, Oriental Books.
- Bharucha, F.R. Afghanistan, India and Pakistan, Plant Ecology. Arid Zone Res. 6: 19-39
- Biddulph, J. Tribes of the Hindu Kush. Graz, Akad. Druok u. Verlagsanstalt 164 1880 (repr. 1971)
- Biwas, 3. On the shrike Lanius tephronotus (Vigors), with remarks on the erythronotus and bioolor groups of Lanius schach Linné and their hybrids. J. Bombay Nat Hist. Soc. 49: 444-455.
- Blanford, W.T. Eastern Persia, zoology and geology. Macmillan, London. 518 pp. 1876
- Blanford, W.T. On the voles (Arvicola) of the Himalayas, Tibet and Afghanistan. J. Proc. 1881 Asiat. Soc. Bengal. 50(2): 88-117.
- Blyth, E. On the species of the genus Ovis. Proc. Zool. Soc. Lond. 8: 62-81.
- Blyth, E. An amended list of the species of genus Ovis. Proc. Zool. Soc. Lond. 1841 1(7): 195-197.
- Blyth, E. Notices and descriptions of various new or little known species of birds.

 1847 J. Asiat. Soc. Beng. 16: 428-476.
- Bobek, H. Die natürlichen Wälder und Gehölsfluren Irans. Bonner Geogr. Abh. 8: 1-62.

 1951 (The natural forests and wooded areas of Iran).
- Bobrinskiji, N.A., Kusnezov, B.A. & Kusjakin, A.P. Mammals of USSR. Moscow (in Russian).

Bogdanov, O.P. Reptiles of Turkmenia. Moscow (in Russian)

Bogdanov, O.P. Ecology of reptiles in Central Asia. Moscow (in Russian)

Bonvalet, G. Through the heart of Asia, over the Pamir to India. 2 Vols. Chapman and 1889 Hall Ltd., London.

Borisov, A.A. Climates of the USSR. Aldine Publishing Co., Chicago. xxi+225 pp 1965

Bornmüller, J. Aus der Flora Afghanistans. Bot. Jahrb. 66: 216-248.

Bowdler Sharpe, R. Scientific results of the Second Yarkand Mission, based upon collections of the late Ferdinand Stoliozka. Aves. London. xvii + 153 pp.

Brockle, S.W. Die Klimabedingungen im afghanischen Raum, Manuskript. (Climatic conditions in Afghanistan. Manuscript).

Breckle, S.W. Vegetation in alpine regions of Afghanistan in Plant Life of South-West 1971 Asia Botanical Soc. of Edinburgh. Aberdeen University Press. pp. 107-117.

Breckle, S.W. Kikroklimatische Messungen und Ökologische Beobachtungen in der alpinen 1973 Stufe des Afghanischen Hindu Kusch. Bot. Jb. 93: 25-55.

Brookle, S.W. & Frey, W. Die Vegetationsstufen im Zentralen Hindu Kusch. Afghanistan J. 1974 1(3): 78-80. (The Vegetation Composition in the Central Hindu Kush)

Breckle, S.W., Frey, W. & Hedge, I.C. Botanical literature of Afghanistan. Notes R. Bot 1969 Odn. Edinb. 29: 357-71.

Bretzl, H. Botanische Forschungen des Alexanderzuges. 412 p., Leipzig. 1903

Bristol University Expedition. The University of Bristol Expedition to the Band-i-Amir 1969 lakes, Afghanistan, 1968. Bristol. 30 p.

British Embassy. List of books on Central Asia in British Embassy Library, Kabul. 1974

Brooks, V. and Brooks, B. On the large sheep of the Tian-Shan and other Asiatic Argali.

1875 Proc. Zool. Soc. Lond. 509-526 pp.

Brown, L.H. East Africa. Chapter 6 in Flamingos Kear, J. and N. Duplaix-Hall (eds.).

1975 T. and A.D. Poyser, Berkhamsted, England.

Brück, G. Zur Herpetofauna Afghanistans. Vest. Asl. Zool. Spol. 32: 201-208. 1968 (On the herpetology of Afghanistan).

Brückle, K. Die Kinerallagerstätten von Ost-Afghanisten. Neues Jb. Kineralogie.

1933 (The ore deposits of eastern Afghanisten).

Brückle, K. Über die Geologie von Bedakhschan und Kataghan. Neues Jb. Miner. Geol., B. 1935 74: 360-401 (On the Geology of Bedakhshan and Qataghan)

Bundesanstalt für Bodenforschung. Geological map of Afghanistan. Hannover.

Burgess, R.L., Noktarzadeh, A. & Cornwallis, L. A preliminary bibliography of the natural 1966 history of Iran. Pahlavi Univ. Coll. of Arts and So., Bull No. 1, Shirax. 220 p.

Burnes, A. Travels into Bukhara; being the account of a journey from India to Cabool, 1934 Tartary and Persia. 3 Vols. London.

Burnes, A. Cabool. John Murray, London. 1843

Burton, P.J.K. Contenus stomacaux de limicoles collectés en Afghanistan. Alauda. 1971 39(2): 132-138.

Buttiker, W. Notizen über die Vogeljagd in Afghanistan. Z. Jagdwiss. 5: 95-105 1959 (Notes on the bird hunting in Afghanistan).

Calembert, J. & Strange, L.R.N. Ecological management of arid and semi-arid rangelands in
1978 Africa and the Near and Middle East (Emasar-Phase 3). Vol.6: Middle East, grassland education and training with special reference to Iran, Afghanistan and
Pakistan. UNEP/FAO, Rome.

Calpp, F.C. Geology of Afghanistan. Bull. Geol. Soc. Am. 50(2) 5: 1904. 1939

Carruthers, D. Severtzoff's sheep from Russian Turkestan. Field. 114: 623

Carruthers, D. Beyond the Caspian. A Naturalist in Central Asia. Oliver and Boyde, 1949 Edinburgh.

Champion, Sir H.G., Seth, S.K. & Khattak, G.M. Forest Types of Pakistan. The Pakistan 1966 Forest Institute, Peshawar.

Christison, A.F.P. Handbook of the Birds of Northern Baluchistan. Quetta. 1940

Christison, A.F.P. Notes on the birds of Chagai. Ibis XIV (S). 531-556 1941

Chuma, T. (ed.) Alingah Valley and Mir Samir, 1966. Afghanistan Expedition of the 1970 Chiba University, Chiba, Japan.

Clark, J.L. The Great Arc of the World Sheep. University of Oklahoma.

Clark, R.J., Clark, E.D., Anderson, S.C. & Leviton, A.E. Report on a collection of amphibians and reptiles from Afghanistan. Proc. California Acad. of Sc. 36(10): 279-316.

Collias, N.E. The analysis of socialisation in sheep and goats. Ecology, 37: 288-239

Conolly, E. Sketch of the physical geography of Seistan. J. Asiat. Soc. Bengal. 9(2): 1840 710-725

Corbet, G.B. The mammals of the Palaearotic Region: a taxonomic review. British Museum 1978 (Natural History), Cornell University Press, 314 pp. London & Ithaca.

Cotton, W.B. The Great Pamir or Maroo Polo Sheep (Ovis ammon pold). J. Bombay Nat. Hist. Soc. 1914 23, Misc. Notes, 770.

Cramp, S. Handbook of the birds of Europe, the Middle East and North Africa. The birds e.o. 1977, of the western Palaearotic. RSPB, Oxford University Press.

1980

Cumberland, G. Sport on the Pamirs. London 1895

Cumming, J.W.N. Breeding Seasons of Big Game. The Persian gazelle (Gazella subgutturosa)
1905a J. Bombay Nat. Hist. Soc. 16(3): Nisc. Notes No. 7,503

Cumming, J.W.N. Birds of Seistan, being a list of the birds shot or seen in Seistan by 1905b members of the Seistan Arbitration Mission, 1903-05. J. Bombay Nat. Hist. Soc. 16: 686-699

Curzon, G.N. The Pamirs and the source of the Oxus. Bull. R. Geogr. Soc. London. 83 p. 1896

Dada, H. & Pickett, L. 1968 Supplement to the bibliography of materials dealing with
1969 agriculture in Afghanistan. Gen. Publ. No. 1. Fac. of Agric. Kabul University
1-33 p.

Danford, G. & Alston, E.R. On the Namuals of Asia Minor. Part I. Proc. Zool Soc. Lond. 1877: 270-281; Part II Proc. Zool. Soc. Lond. 1880: 50-64

Davis, P.H., Harper, P.C. & Hedge, I.C. The Plant Life of South West Asia. The Botanical Society of Edinburgh. 335 p.

DeBlase, A.F. New distribution records of bats from Iran. Fuldrana Zool. 58: 9-14

De la Tour, G.D. Eine ökozoogeografische Revision der mittelasiatischen Halbesel, 1775 Equus hemionus Pallas. Säugetierkundliche Kitt. 23(2): 108-111

Dementiev, G. Theresa's Ground Finch, Montifringilla theresee. Ibis. 105: 107 1963

Demontiev, G.P. & Gladkov, N.A. The birds of the Soviet Union. Moscow (in Russian)

Desro, A., et al. On the geology of Central Badakhshan (NE Afghanistan). Q.J. Geol. S. 1964 London. 120(478): 127-151

Deutsche im Hindukusch. Deutsche Forschung., N.F. 1: 135

Dieterle, A. Vegetationskundliche Untersuchungen im Gebiete von Band-i-Amir (Zentral-1973 Afghanistan). (Research on the vegetation in the Band-i-Amir area) Dissues bot. 24 p.

Diver, M. Kabul to Kandahar. Davies, London. 192 p. 1935

Dollot, R. L'Afghanistan. Paris: Payot. 318 p. 1937

Dor, R. Lithogylphes du Wakhan et du Pamir. Afghanistan J. 3: 122-129 1976a

Dor R. Note sur le Yak au Pamir. Published in: Le Yak: son rôle dans la vie matérielle 1976b et culturelle des éleveurs d'Asie Centrale. Soc. d'Ethnozooteohnie, Paris. p. 126-132 (Remarks on the Yak in the Pamir)

Dor, R. & Naumann, C. Die Kirghisen des afghanischen Pamir. Graz. (The Kirghises of the Afghan Pamir)

Dückelmann, R. von. Die Holzgewächse und Waldformen von Afghanistan nach ihrer klimatischen 1951 Bedingtheit und geografischen Verteilung. in: Generalbericht d. Abt. Forstwesen Kgl. Dept. für Landwirtschaft (The trees and forest types of Afghanistan, according to climatic conditions and geographic distribution)

Dunmoor, The Earl of. The Pamirs. 2 Vols. John Murray, London 1894

Dupree, L. Note on the distribution of the Indian crested porcupine: Afghanistan.

1956 Nammalogy. 37/2: 299-300

Dupree, L. Shamshir Chor. Anthrop. Pap. Am. Mus. Nat. Hist. 46(2)

Dupree, L. Prehistoric research in Afghanistan. American Philosophical Society, 1972 Philadelphia. 84 p.

Dupree, L. Afghanistan. Princeton University Press, New Jersey. 760 pp. 1973

Dupree, L. Settlement and migration patterns in Afghanistan: A tentative statement.

1975a Nodern Asian Studies. 9(3): 397-413

Dupres, L. New Paleolithio Localities near Dashte-i-Nawar. Afghanistan J. :2 - 105

Dupree, L. & Albert, L. eds. Afghanistan in the 1970s. Praeger. 1974

Economist Intelligence Unit. Quarterly Economic Review of Pakistan, Bangladesh, Afghanistan.

1979, Spencer House, London.

1980

Edelberg, L. & Schuyler. Nuristan. 160 pp. 112 plates, 60 figures. Akademische 1978 Druck-u. Verlagsanstalt. Graz/Austria.

Egorov, O.V. Wild ungulates of Yakutia. Isdatel'stvo 'Nauka', Moscow. Translated 1967 1965 by Israel Programme for Scientific Translations, Jerusalem.

Electrowatt & Norconsult. Konar River Basin Development. Kasterplan studies, Annax 10.
Ninistry of Water and Power, Afghanistan.

Ellerman, J.R. Key to the rodents of South-West Asia in the British Museum Collection.

1948 Proc. Zool. Soc. Lond. 118: 765-816

Ellerman, J.R. The fauna of India, Mammalia (2nd ed.) Vol. 3: Rodentia. Calcutta. 1961

Ellerman, J.R. & Chaworth-Musters, J.L. A revision of the genus Mariones. Proc. Zool. 1947-48 Soc. Lond. 117:

Ellerman, J.R. & Morrison-Scott, T.C.S. Checklist of Palaearotic and Indian mammals, 1950 to 1946. 1966 2nd ed. London.

Eskilsson, E. Report on coordinating the development of Afghanistan's energy resources.

1966 UNDP, Kabul.

Etchecopar, R.D. L'extension de <u>Psittaoula krameri</u> (la perruche à collier rose) au Moyen
1969 Orient. Oiseau Revue Fr. Orn. 39: 178-181. (The extent of <u>Psittaoula krameri</u> (rose-ringed parakset))

Etemad, E. A Checklist of Iranian Bats. Bats of Iran (in Farsi). Teheran 1968

Fairservis, Walter A., Jr. Archaeological Studies in the Seisten Basin of Southwestern
1961 Afghanistan and Eastern Iran. Anthrop. Pap. Am. Nus. Nat. Hist. 129 p. 19 plates.

FAO Note sur la végétation des terrains de parcours de la région d'Andkhui, par Pabot, H.

1958 FAO Report to the Government of Afghanistan. 3 p. Kabul.

Rapport sur l'amélioration des pâturages naturels (de l'Afghanistan), par Pabot, H.

Report on the improvement of the natural pastures (of Afghanistan).

FAO General soil map of Afghanistan, by Subramanian, V.S., Nasirov, V. & Salem, M. Z.

1962a Report of the first soil correlation seminar for South and Central Asia, Tasjkent,
USSR, 14 September - 2 October 1962, p. 56-59. World Soils Resources Report No.4.

- FAO Generalized soil correlation seminar for South and Central Asia, by Subramanian, V. S.

 1962b Nasirov, V. & Salem, N.Z. Soil map of the world. FAO/UNESCO. Project World Soil

 Resources Reports 4: 56-59
- Survey of land and water resources, Afghanistan. Vol 1 general, Vol 2 geology, topo-1965 graphic surveys, Vol. 3 hydrology, Vol 4 soils and land classification, Vol 5 water control, irrigation, Vol 6 Irrigation development Hari Rud and Kabul Valleys. Rome, FAO/SF:9/AFC.
- Pasture development and range improvement through botanical and ecological studies, 1967 by Pabot, N. FAO/UNDP Report TA 2311, Rome.
- FAO The planning, execution and reconstruction of irrigation projects, by Horning, H.M. 1969a FAO/UNDP Report TA 2719. 44 p.
- FAO Improvement of sheep husbandry. Report to the Government of Afghanistan by Imankulov, 1969b H.I. FAO, Rome.
- Survey of irrigation possibilities in the Hari Rud and Upper Kabul River basins.

 1969 Series of Technical reports on geology, topography, hydrology, land tenure, soils,
 70, 71 agronomy, sociology, etc. and final report. UNDP/SF project 80/ Afg/ 10
- FAO Afghanistan, wildlife resources by G. Caughley. Project Report on wildlife, animal 1970, resources and wildlife conservation. FAO/UNDP report No. 2905, 15 p. 1971
- FAO Demonstration et formation en aménagement des forêts et des pâturages par L. Bostanoglu.

 1973a FAO/UNDP—SF/AFG/67/515, 193 p. (Demonstration and training in forest and range improvement)
- FAO Etude sur la classification écologique de la végétation ligneuse naturelle en 1973b Afghanistan by Nedialkov, S.T. Project UNDP/FAO/AFG/67/515. 46 p. (Study on ecological classification of the natural woody vegetation of Afghanistan)
- FAO Plan d'aménagement de la forêt de Tohapadara (région de Konar) par Nedialkov, S.T. 19730 FAO/AFU/67/515 (Management plan for the Tohapadara forest)
- FAO Progress Report No. 1: Conservation and Utilisation of Wildlife Resources by 1973d Petooz, R.G. FO:DP/AFG/72/005, Rome, 15 p.
- FAO Outline of hunting and tredding programmes: ourrent status and future requirements by 1973e Petooz, R.G. & King R.C. FO:DP/AFG/72/005, Kabul, 7 p.
- PAO Background information on fluctuations in animal population sizes with comments on the expanding rodent population in the north of Afghanistan by Petoos, R.G. FO:DP/AFG/72/005, Kabul, 4 p.
- FAO Formation et démonstration pour le développement des forêts et des pâtureges en

 1974 Afghanistan par F. Genty. FAO, Rome. projet AFO/67/515. (Training and demonstration for the development of the forests and pastures in Afghanistan).

- FAO National Parks, Wildlife Reserves and Sanctuaries in Afghanistan by R.G. Petooz & 1974b R.C. King. FO:DP/AFG/72/005, Kabul.
- FAO Report on the status of the Band-Amir National Park by Petooz, R.G. & T. Skogland.

 19740 FAO/UNDP, FO:DP/AFG/72/005, Kabul, mimeo, 18 p.
- FAO Range management and animal husbandry practices in Afghanistan. Demonstration and 1974d training in forest and range improvement by O.N. Alpay. FO:DP/AFG/67/515, Rome.
- FAO The Flamingos (Phosnicopterus ruber roseus) of Ab-i-Estada and Dashte Navar, Chazni
 1975a Province Afghanistan by R.G. Petocz & K. Habibi. FO:DP/AFG/72/005, Kabul,
 mimeo, 12 p.
- FAO The birds of Band-i-Amir, Dashte-i-Nawar and Ab-i-Estada by R.G. Petooz, K. Habibi 1975b & T. Skogland. FO:DP/AFG/72/005, Kabul, mimeo, 8 p.
- FAO Tentative list of bird species from the Central Highlands by R.G. Petooz, K. Habibi 1975c & T. Skogland. FO:DP/AFG/72/005, Kabul, mimeo, 8 p.
- FAO Ecology and behaviour of Marco Polo sheep (Ovis ammon polo) in the Pamir during 1975d winter by R.G. Petcoz & T. Skogland. FO:DP/AFC/72/005, Kabul, mimeo, 31 p.
- FAO Recommendations on the status of the Bactrian deer (Cervus elaphus bactrianus) in 1975e Ajar Valley reserve by K. Habibi. FAO/AFG/72/005, mimeo, 3 p.
- FAO The birds of Hamun-i-Puzak by R.G. Petocz, K. Habibi & W.F. Rodenburg. FO:DP/AFC/1976a 72/005, Kabul, mimeo, 7 p.
- FAO Ecological reconnaissance of the Hindu Kush Ibex (Capra ibex) in Ajar Valley,
 1976b Bamiyan Province, Afghanistan, by T. Skogland. FO:DP/AFC/72/005, Kabul, mimeo
 18 p.
- FAO Ecological and silvicultural study of the forestry vegetation of Kunar region by S.T. Nedialkov. FAO/UNDP: AFC/74/004, Assadabad, Kunar.
- FAO Brief on Iran trip, 16-26 January, 1976, by R.G. Petcoz. FAO/UNDP:FO:DP/AFG/74/016, 1976d Kabul, mimeo, 4 p.
- FAO Birds of the Hamun-i-Puzak by W.F. Rodenburgh, K. Habibi & C.C. Shak. Supplement I 1976e 7-10 December. FAO, 74/016, mimeo, 5 p.
- FAO Checklist of the birds of Afghanistan by K. Habibi. FO:DP/AFU/72/005, Kabul, mineo, 1976f 17 p.
- FAO Conservation and utilization of wildlife resources. Afghanistan. Interim Report. 1977a FO:DP/AFG/72/005, Rome.
- FAO Brief on a reconnaissance trip to Herat, Badghis and Chor Provinces, 17 May 3 June, 1977b by R.G. Petcoz, G. Shank, K. Habibi & N. Komak. FAO/UNDP Kabul: FO:DP/AFG/74/016, mimeo, 4 p.

- FAO The Mammals of Afghanistan, their distribution and status by K. Habibi. Field Working 19770 Document I. FO:DP/AFG/74/016, Kabul.
- FAO The Trade in Wild Animal Furs in Afghanistan by W.F. Rodenburg. Field Document No. 2. 1977d FO:DP/AFG/74/016, Kabul.
- FAO Management plan for Ab-i-Estada and Dashte Nawar flamingo and waterfowl sanctuaries 1977e by C.C. Shank & W.F. Rodenburg. Field Document No. 3. FO:DP/AFG/74/016, Kabul.
- FAO A preliminary management plan for the Ajar Valley wildlife reserve by C. Shank, 1977f R.G. Petooz & K. Habibi. FO:DP/AFG/74/016, Field Document No. 4, Kabul.
- FAO Establishment of forest nurseries and pilot plantations in Kunar by L. Bostanoglu. 1977g FAO/UNDP Assadabad, AFC/74/004 and AFC/77/001. Final Technical Report.
- FAO Report of the special survey of southwestern Afghanistan (Desert locust control) by 1978a K.R. Ehatia. FAO-AGP-SWA/DL/SS/13, 13 p.
- FAO Stock: starvation and surplus by M.H. Woodford. FO:DP/AFG Kabul. 1978b
- FAO Afghanistan, project findings and recommendations. National parks and utilization of 1978c wildlife resources. FO:DP/AFC/74/016. Terminal Report.
- FAO Report on the Afghan Pamir by R.C. Petocz. Part I: Ecological Reconnaissance.

 1978d Project Field Document No. 5. FO:DP/AFG/74/016, Kabul.
- FAO Report on the Afghan Pamir by R.G. Petooz, K. Habibi, A. Jamil & A. Wassey. Part II: 1978e Biology of Marco Polo sheep (Ovis ammon pold). Project Field Document No. 6. FO:DP/AFC/74/016, Kabul.
- FAO Report on the Afghan Pamir by R. Petocz. Part III: Management plan for the Big Pamir 1978f wildlife reserve. Project Field Document No. 7. FO:DP/AFG/74/016, Kabul.
- FAO A strategy for the establishment and development of Bande Amir National Park by C. Shank 1978g and J.Y. Larsson. Project Field Document No. 8. FO:DP/AFG/74/C16, Kabul.
- FAO Ecological reconnaissance of western Nuristan with recommendations for management by 1978h R.G. Petcoz and J.Y. Larsson. Project Field Document No. 9. FO:DP/AFG/74/016, Kabul.
- FAO A preliminary study of Lake Hashmat Khan with recommendations for management by A. 1978i Rahim & J.Y. Larsson. Field Document No. 10. FO:DP/AFG/74/016, Kabul.
- FAO Status of Alpine rangelands in Central Afghanistan with special reference to the Ajar 1978j Valley wildlife reserve by J.Y. Larsson. Field Document No. 1. FO:DP/AFG/78/007, Kabul.
- FAO Evaluation of the Kabul Green Belt by B.S. Kernan. Field Document No. 1. FO:DP/AFG/1979a 79/009, Kabul.

- 5. Suncus etruscus (Savi) pygmy white-toothed shrew Recorded from Kotal-e-Rabatak (Samangan, N.) (CN & JN).
- Suncus murinus (Linnaeus) house shrew

 S. m. griffithi (Horsfield). Lower Kabul Valley and Nuristan.

 Jalalabad is the westernmost record for the species, in houses and open vegetation. Kunar (JN), Jalalabad, Laghman (JH).

ORDER CHIROPTERA

Family Rhinopomatidae

- 1. Rhinopoma hardwickei Gray lesser mouse-tailed bat

 R. h. hardwickei Gray. Jalalabad Valley III-V/67 (JG), Nangarhar,

 Kunar (E), small caves (MO), Chaunchir (Kandahar) (KZ), (VA).

 R. h. bactrianus Satunin. Northern Afghanistan, but not yet collected.
- 2. Rhinopoma microphyllum (Brunnich) larger mouse-tailed bat

 R. m. microphyllum. Arid regions of the Jalalabad Valley (JG).

 Nangarhar, Kunar (E), small caves (MO), Girishk, Farah (SW), Sham Chir (SE), (VA), (KZ).
- 3. Rhinopoma muscatellum Thomas

 R. m. seianum Thomas. Recorded from Sham Chir (SE) (KZ).

Family Emballonuridae

Taphozous nudiventris Cretzschmar tomb bat
 T. n. kachhensis Dobson. Drier lowland areas, Lashkargah 990 m VIII/65 (MO), Chak Naur (Jalalabad Valley) IV/67 (JG).

Family Megadermatidae

 Megaderma lyra Geoffroy Indian false vampire
 M. 1. lyra Geoffroy. Oriental species, Jalalabad Valley III/67 (JG), Kunar River (Qachgar Cave) (VA).

Family Rhinolophidae

- 1. Rhinolophus blasii Peters Blasius' horseshoe bat Records from Laghman up to 2 000 m, Kabul, Parwan, Kandahar and Maimansh (JG), East Afghanistan (VA).
- 2. Rhinolophus clivosus Cretzschmar

 R. c. bocharicus Kastschenko and Akimov. One record from Maimanah (JG).
- 3. Rhinolophus ferrumeguinum (Schreber) greater horseshoe bat

 R. f. proximus Andersen. Recorded from Lalanda (20 km SSW of Kabul)

 V/67 (JG), (Lindberg), Kabul, Laghman (MO), Maimanah, Beltchiragh,
 Girishk (VA). (Southern specimen perhaps R. f. irani Cheesman).
- 4. Rhinolophus hipposideros (Bechstein) lesser horseshoe bat

 R. h. midas Andersen. Records up to 2 000 m from Jalalabad, Sarobi,

 Abdul Khel (E) II, IV/67 and Lalanda (20 km SSW of Kabul) V/67 (JG),

 Qalat (Bolan Cave) (VA).

- FAO The sheep breeds of Afghanistan, Iran and Turkey by B.C. Yaloin. FAO/UNEP. Project 1970b FP/1109-76-02(883), Rome.
- FAO Forest inventory field manual: Afghanistan by H.J. McCarthy. FO:DP/AFG/77/001, Kabul, 19790 40 p.
- FAO Forest inventory handbook: Afghanistan. Field Document No. 3. FO:DP/AFU/77/001, 1979d Kabul, 83 p.
- FAO National parks and wildlife management Afghanistan, project findings and recommenda-1980a tions. FO:DP/AFG/78/007, Terminal Report, Rome.
- FAO Watershed Management, Afghanistan, by M. Güloür. Field Dooument No. 7. FAO/UNDP: 1980b FO:DP/AFG/79/009, Rome.
- FAO Assistance to forestry and watershed development, Afghanistan. UNDP/FAO: FO:DP/AFG/1980c 79/009, Terminal Report, Rome, 29 p.
- FAO & IBPOR Consultative meeting on the orop genetic resources of Southwest Asia, 1979 Rome, 8-10 October. TF/REM/31(IBPOR).
- FAO/UNEP Technical consultation on animal genetic resources conservation and management.

 1980 Recommendations. Rome, 2-6 June, 4 p., mimeo.
- FAO & Unesco Soil map of the world, 1:5 000 000. Volume 7, South Asia. Unesco, Paris.
- Fayman, T. Preparation of an area development plan for the tourist region, Bamiyan, Bande 1976 Amir, Ajar Valley. UNDP, Kabul.
- Pelten, H. Bermerkungen zu Fledermäusen der Cattungen Rhinopoma und Taphozous. Senokenberg.

 1962 Biol. 43, No.(2):171-176. (Remarks on the bats of the genus Rhinopoma and Taphozous).
- Felten, H. Zur Taxonomie indo-australischer Fledermäuse der Cattung <u>Tadarida</u> Senok.

 1964 Senokenberg. Biol. 45, No.(1): 1-13. (On the taxonomy of the Indo-Australian bats of the genus <u>Tadarida</u>).
- Ferdinand, K. Nomed expension and commerce in Central Afghanistan. Folk. 4: 125-159.
- Ferdinand, K. Ethnographical notes on Chahar Aymaq, Hazara and Monghols. Acta Orientalia 1964 28: 175-203.
- Ferdinand, K. Nomadism in Afghanistan (with an appendix on milk production). Vieh-Wirt-1969 schaft und Hirten Kultur, Ethnographische Studien Bulapest Kiado. 127-160 p.
- Ferdinand, K. Aspects of the relations between nomads and settled populations in 1976 Afghanistan. 7th Intern. Congr. Anthropol. Ethnol. So. 10: 125-133, Noscow.
- Ferrier, J.P. Caravan journeys and wanderings in Persia, Afghanistan, Turkistan and 1976 Baluchistan. Karachi. O.U.P.

- Field, N.C. The Amu Daria, a study in resource geography. Geogr. Review. October. 1954 528-542 p.
- Finn, F. List of the birds collected by the Afghan-Baluch Boundary Commission of 1896.

 1896 J. Asiat. Soc. Beng. 65: 566-567.
- Finnoonsult Socio-economic survey of Bedakhshan. AFG/75/024, Field Document, 3 Vols., 1976 Kabul.
- Firouz, E. Environment Iran. Nat. Soc. for the Conserv. of Natural Resources and Human 1974 Environment, Tehran. 53 p. (4 maps, photographs).
- Firouz, E. & Marrington, F.A. Iran: Concepts of biotic community reservation. IUCN, 1975 Regional Meeting.
- Firouz, E., Massinger, J.D. & Ferquson, D.A. The wildlife parks and protected regions of 1970 Iran. Biol. Cons. 3(1): 37-45.
- Fischer, W.B. Afghanistan, physical and social geography. From: The Middle East and North 1978/9 Africa.
- Flohn, H. Zum Klima und Wasserhaushalt des Hindukuschesund der benachbarten Hochgebirge.

 1969 Erdkunde. 23: 205-215. (On olimate and water economy of the Hindu Kush and adjacent areas.)
- Floyer, E.A. Unexplored Baluchistan. Gosha-E-Adab, Quetta (Pakistan).
- Forster, U. & Bartsoh, C. Die Seen von Bande Amir, Dascht-i-Nawar, Ab-i-Istada und Hamun1970 i-Puzak (Zentral und Südwest Afghanistan). Science (Kabul):19-23. (The lakes
 of Bande Amir, Dashte Nawar, Ab-i-Estada and Hamun-i-Puzak (Central and Southwest
 Afghanistan.)
- Fox, C.S. The coalfields of Afghanistan. Report for the Geological Survey of India. 45 p. 1936
- Freise, F. Geografische Verbreitung und wirtschaftliche Entwicklung des Bergbaus in Vorder1907 und Kittelasien während des Altertums. Z. Prakt. Geol. 15: 73-117. (Geographical distribution and economic development of mining in the Near East and Central
 Asia during ancient times.)
- Freisleben, R. Anschauungen über die Abstammung der Kulturgersten auf Grund des Materials

 1941 der Deutschen Hindukusch Expedition 1935. Forschung und Fortschritte. 17: 140
 142. (On the origin of cultivated barley based on material collected by the

 German Hindu Kush Expedition 1935.)
- Freitag, H. Die natürliche Vegetation Afghanistans. (Beiträge zur Flora und Vegetation 1971 Afghanistans I.) Vegetatio 22: 286-344. (The natural vegetation of Afghanistan. Contributions to the flora and vegetation of Afghanistan I.)

- Freitag, H. Study on the natural vegetation of Afghanistan: on plant life in Southwest
 1971 Asia, P.H. Davis, P.C. Harper and I.C. Hedge (eds.), University Press, Aberdeen.
 p. 80-106.
- Freitag, H. Flora und Vegetation: Kraus W. 1972 Afghanistan: Natur, Geschichte und Kultur, 1972 Staat, Gesellschaft und Wirtschaft. Tübingen. (Flora and Vegetation)
- Frey, W. Zur Vegetation des zentralen afghanischen Hindukusch in: Zwischen Munjan und 1967 Bashgal. Göppingen. 1: 17-46. (On the vegetation of Hindu Kush in Central Afghanistan.)
- Frey, W. Beobachtungen im zentralen Hindukusch. In: Zwischen Munjan und Bashgal, Göppingen.

 2: 24-35. (Observations in Central Hindu Kush)
- Fulton, N.T. Notes on the birds of Chitral. J. Bombay Nat. Hist. Soc. 16: 44-64.
- Fulton, H.T. Additional notes on the birds of Chitral. J. Bombay Nat. Hist. Soc. 1905 16: 744.
- Furon, R. L'Afghanistan. Géographie, histoire, ethnographie, voyages. Avec carte et ill., 1926 133 p. Ethn. Nus. Paris.
- Furon, R. La géologie du plateau iranien (Perse, Afghanistan, Béloutschistan). Revue gén. 1937 Sci. pur. appl. 48(2): 36-43.
- Furon, R. La géologie du plateau iranien (Perse, Afghanistan, Béloutschistan). Mém. Mus. 1941 natn. nat., Paris. 7(2): 178-414.
- Furon, R. & Louis-Felicien, R. L'Afghanistan et les milieux paléontologiques mondiaux
 1954 Afghanistan. 9(4): 48-50. (Afghanistan and the World Paleontological Contexis)
- Furse, P. Iran and Afghanistan, 1964. Jl. R. Hort. Soc. 90: 462-475, 504-509; 1964 91: 18-26.
- Furse, P. & P. Afghanistan, 1966. Jl. R. hort. Soc. 93: 20-23, 92-97, 114-124. 1968
- Gaisler, J. The bats (Chiroptera) collected in Afghanistan by the Czechoslovak Expedition 1970a of 1965-1967. 4(6): 1-56, Prirodov. Pr. Cesk. Akad. Ved.
- Gaisler, J. Zoogeographical notes on the bat fauna of Afghanistan (Mammalia, Chiroptera)
 1970b 34(4): 284-288, Vest &sl. zeměd Mus, zool Spol.
- Gaisler, J. Systematic review and distinguishing characters of the bats (Chiropters) hiterto recorded in Afghanistan. Zool. Listy, 20(2): 97-110.
- Gaisler, J. Rodents in desert environment. Comparative ecological notes on Afghan rodents.

 1975 Konographie biol. 28: 59-74.

30

- Gaisler, J., Povolny, D., Sebek, Z. and Tenora, F. Faunal and ecological review of mammals 1967,68 occurring in the environs of Jalalabed, with notes on further discoveries of mammals in Afghanistan. I Insectivora, Rodentia. II Chiroptera, III Carnivora, Legomorpha. Zool. Listy. 16(4): 355-364; 17(1): 41-48; 17(2): 185-189.
- Ganss, O. On the geology of SE Afghanistan: I Geology of the area Ghazni-Gardez-Urgan,
 1970 II Geology of the area around Muhur and Ab-i-Estada. Bull. Afgh. Geol. Miner.
 Survey. 3: 1-203.
- Gansser, A. Geology of the Himalayas. Interscience Publishers, London. xv+ 289 pp. 1964
- Gee, E.P. The wildlife of India. Collins, London. 1964
- Geist, V. Mountain sheep. The University of Chicago Press, Chicago. 383 p. 1971
- Geographical Section. Asia, 1:1 000 000. World maps, sheets: J-41: Bukhara; J-42: 1942 Samarqand; J-43: Kashmir. General Staff, no. 2555. 2nd edition. London.
- Gilbert, O.L. A general account of high altitude vegetation in the Samir Valley. Environ1968a mental research in the Samir Valley of the Hindu Kush. Ed. A. James, Newcastle.
- Gilbert, O.L. Lichen ecology in the Hindu Kush. Environmental research in the Samir Valley 1968b of the Hindu Kush. Ed. A. James, Newcastle. 6.5 6.7
- Gillan, G.V.B. The distribution of the Great Pamir sheep (Ovis ammon poloi). J. Bombay 1934 Nat. Hist. Soc. 37(1): 216-217.
- Gilli, A. Neue Arten aus Afghanistan. Reprium nov. Spec. Regni veg. 56: 143-149; 1954-64 57: 94-101; 57: 218-231; 59: 162-178, 190-192; 62: 22-26, 193-209, 209-211; 63: 321-328; 66: 192-194; 68: 87-100; 69: 175. (New plant species from Afghanistan.)
- Gilli, A. Beiträge zur Flora Afghanistans. I Pteridophyta und Gymnospermae. II Mono-1958-64 ootyledines. III Salioales, Fagales, Urticales, Poleygonales, Centrospermae. IV Ranales und Rhoedales. V Rosales, Geraniales. Reprium nov. Spec. Regni veg. 61: 86-92; 64: 204; 68: 239-259; 69: 155-175.
- Gilli, A. Afghanische Pflanzengesellschaften. Vegetatio, 16: 307-375. (Afghan plant 1969 communities)
- Gladkov, N.A. & Nasimovich, A.A. Rare and disappearing animal species and their preservation in the USSR. In: Resources of the biosphere on the territory of USSR, Viniti, Noscow. 311-333 (in Russian).
- Godwin-Austen, K.N. Land and freshwater mollusca of India, including South Arabia, 1882-1914 Baluchistan, Afghanistan, Kashmir, Nepal, Burma, Pagu, Tenasserim, Malay Peninsula, Ceylon and other islands of the Indian Ocean. Vol. 1, prt 1-6; Vol. 2, prt. 7-12.

- Gorgas, N. Betrachtung zur Hirnschädelkapazität zentralasiatischer Wildsäugetiere und 1966 ihre Hausformen. Zool Anz. 176: 227-235. (Observations on the capacity of the oranium cerebrale of Central Asian wild mammals and their domestic forms)
- Gratzi, K. ed. Hindukusch, Österreichische Forschungsexpedition in den Wakhan. Graz, 1974 (2nd ed.) 148 p. (Hindu Kush, Austrian exploration expedition in the Wakhan)
- Griffith, W. Journals and travels in Assam, Burma, Bhutan, Afghanistan and the neighbouring 1947 countries. Calcutta. xxii + 529 p.
- Grote, H. Ubersicht über die Säugetierfauna des Pamir Gebietes. (Survey of the mammal 1951 fauna of the Pamir area) Zool. Gart. Frankf. 18: 142-145.
- Groun-Grachimailo, Gr. Le Pamir et sa faune lépidoptérologique in: Romanoff, Mémoires sur 1890 les lépidoptéres, Vol IV, St. Petersbourg. (Pamir and its butterfly fauna).
- Oroves, C.P. On the smaller gazelles of the genus Cazella. Z. Säugetierk. 34: 38-60
- Croves, C.P. & Mazak, V. On some taxonomic problems of Asiatic wild asses, with the description of a new subspecies (Perissodactyla; Equidae). Z. Säugetierk. 32(6): 321-335.
- Guggisberg, C.A.W. Wild cats of the world. David and Charles. London. 328 pp. 1975.
- Habibi, K. Sr. A short history of Afghanistan (in Dari). Vol. II Kabul Historical and Literature Soc. Afghanistan Academy.
- Haeckel, I. & Troll, W. Botanische Ergebnisse der Deutschen Hindukusch Expedition 1935.

 1938 Reprium nov. Spac. Regni veg. Beih. 108: 1-72. (Botanical results of the German Hindu Kush Expedition)
- Hanak, V. & Gaisler, J. Notes on the taxonomy and ecology of Nyotis longipes (Dobson, 1969 1873). Zool. Listy, 18(3): 195-206.
- Harrison, D.L. The mammals of Arabia. Vol. I., Insectivora-Chiroptera, Primates. 1964 Vol. II, Carnivora - Artiodactyla, Hyracpodas. London.
- Harrison, S.G. Edible pine-kernels; Afghanistan and Baluohistan. Kew Bull. 1951: 1952 372-373.
- Hassanyar, A.S. Environmental problems of Afghanistan. A National Report to the U.N. 1972 Conference on the Human Environment. Stockholm, Sweden. 17 pp.
- Hassanyar, A.S. Afghanistan forests. Geogr. Rev. Afghan., 13. 1-7 pp. 1973a
- Hassanyar, A.S. Problems of the biosphere in Afghanistan. Geogr. Rev. Afghan., 13. 1973b 83-88 pp.

- Hassanyar, A.S. Major life zones of Afghanistan. Kabul University Press, Kabul. 23pp. 1973c
- Hassanyar, A.S. Restoration of arid and semi-arid ecosystems in Afghanistan. Env. Cons. 1977 4(4).
- Hassinger, J.D. Introduction to the mammal survey of the 1965 street expedition to 1968 Afghanistan. Fieldiana, zool. 55: 1-81. 165 pp.
- Hassinger, J.D. A survey of the mammals of Afghanistan. Fieldiana, zool. 60: 195.
- Hassinger, J.D. Shrews of the <u>Crooidura zarudnyi pergrisea</u> group with description of a 1970b new subspecies. Fieldisna, zool. 58(2): 5-8.
- Hauser, G.F. Afghan alkali soil and its improvement. Soil Sci. 76: 75-367.
- Hay, R.G. Notes on the wild sheep of the Hindoo Koosh. J. Asiat. Soc. Bengal. 9(1): 440-1840 443.
- Hay, W.R. Band÷i-Amir. Geogrl. J. 87: 348-350. 1935
- Hayden, H.H. The geology of northern Afghanistan. Mem. geol. Surv. India. 39: 1-96.
- Hayon, J.C., Pelt, J.M. & Younos, M.Ch. Les formations végétales de la basse vallée du 1968 Koumar (Afghanistan Oriental). Compt. rend. Acad. So. Paris. 266, Ser. D: 2086-2089 (The vegetation types of the lower Kumar Valley, Eastern Afghanistan).
- Hayon, J.C., Pelt, J.M. & Younos, Ch. Les formations végétales de la moyenne vallée du

 1970 Kaboul et des massifs montagneux du Nuristan (Afghanistan). Vegetatio 20: 279-306.

 (The vegetation types of the middle part of the Kabul River Valley and of the mountain massifs of Nuristan)
- Hedge, I.C. Studies in the vegetation of Afghanistan. Notes R.B.G. Edinb. 23: 163-165, 1960-68 205-208; 26: 407-425; 27: 149-173; 28: 163-172, 89-161 (with Lemond, J.K.)
- Hedge, I.C. Notes on the giant asphodels in Afghanistan. Journ. Roy. Hort. Soc. 88: 402-1963 406.
- Hedge, I. & Wendelbo, P. Studies in the flora of Aghanistan. Arbok Univ. Bergen. 16: 1-56.
- Hedge, I. & Wendelbo, P. Some remarks on endemism in Afghanistan. Isr. J. Bot. 19(2/3): 1970 401-417.
- Hemmer, A., Schmidtler, J.F. & Böhme, W. Zur Systematik zentralasiatischer Grünkröten

 1978

 (Bufo viridio complex) (Amphibia, Sahenhia, Bufonidae). Zool. Abh. Nus. Turk.

 Dresden. 34(24): 349-384. (On the systematics of the Central Asian toad (Bufo viridis complex))

Heptner, W.G. Notes on the Gerbillidae (Mammalia, Clires) IX, Remarks on a new species of

1937

Meriones from Turkestan and on the systematic position of the gerbils belonging
to the Meriones persions group. Bull. Mosk. Cbshch. Isp. Prirody, Sect. Biol.

N.S. 46(4): 189-193.

Heptner, V.G. et al. The mammals of the Soviet Union (in Russian).

Herbordt, O. Die Erzlagerstätten Afghanistans. Internationale Bergwirtschaft, Vol. II, 1926 Leipzig. (The places with ore deposits of Afghanistan)

Herman, N.M. Le climat de l'Afghanistan. Monogra Mét. natn. 52: 55. 1965

Herman, N.N., Zillhardt, J. & Lalande, P. Recueil de données des stations météorologiques 1971 de l'Afghanistan. Kabul publications de l'institut de météorologie No. 2, 1-58. (Results of the meteorological stations of Afghanistan)

Heydemann, F. Beiträge zur Lepidopteren - Fauna Afghanistans. Z. wien. ent. Ges. 39: 1954 385-396, 412-428. (Contributions to the Lepidoptera - Fauna of Afghanistan)

Hill, J.E. Indo-Australian bats of the genus Tadarida. Mammalia. 25, 1.

Holdich, T.H. The origin of the Kafir of the Hindu Kush. Geogr. J. 7: 42-49. 1896

Holloway, C.L. Threatened deer of the world: research and conservation projects under the 1975 IUCN program. Deer 3(8): 428-433.

Horsfield, T. & Moore, F. A catalogue of birds in the Museum of the Hon. East India 1854 Company. London, W.H. Allen, 2 Vols.

Hué, F. & Etchecopar, R.D. Les ciseaux du proche et du moyen crient. Editions N. Boubée 1970 & Cie. Paris, 950 p.

Humlum, J. La géographie de l'Afghanistan. Copenhague, Gyldendal. 424 p. map supplement. 1959

Huntington, E. The basin of Eastern Persia and Seistan in: Exploration in Turkestan with an account of the Basin of Eastern Persia and Seistan. Carnegie Inst. Publ. No. 26: 219-317. 26 abb., Washington.

Hutton, I. The wild sheep of Afghanistan, 'koh-i-poombur' of the Afghans or bearded sheep 1842a or Pennant. Calcutta Journ. Nat. Hist. 2: 514-521.

Hutton, I. Borz of the Afghans - Wild goat of Afghanistan. The Pasang of Authors (Capra 1842b aegagrus). Calcutta Journ. Nat. Hist. 2: 523-535.

Hutton, I. Markhore or the snake eater of the Afghans. Wild goat of Afghanistan (Capra 18420 megaceros). Calcutta Journ. Nat. Hist. 2: 535-542.

- Hutton, T. Rough notes on the zoology of Candahar and the neighbouring districts (with footnotes by E. Blyth). J. Asiat. Soc. Bengal. 14: 340-354; 15: 135-170.
- Hutton, T. Rough notes on the ornithology of Candahar and its neighbourhood (with some additional information on the birds of Afghanistan by E. Blyth). J. Asiat. Soc. Bengal. 16: 775-794.
- Inskipp, T. & Wells, S. International trade in wildlife. An Earthsean Publication, London. 1979
- Irwin, Lieut. Memoir on the climate, soil, produce and husbandry of Afghanistan and neigh-1839 bouring countries. J. Asiat. Soc. Beng. 3(8,2): 1005-1015.
- Isakov, Yu. A. & Shevareva, T.P. Interrelationship of waterfowl breeding and wintering areas in the central Palearotic. Proceedings of a Technical Neeting on Wetland Conservation. IUCN Publ. New Series No. 12, Morges. 165-174 pp.
- IICN Ecological guidelines for the use of natural resources in the Middle East and South 1975 West Asia. IUCN Publ., New Series No. 34, Morges.
- IUCN Red Data Book, Volume 1, Mammalia, Revision by J. Thornback of 1972 edition.
- IUCN Red Data Book, Volume 2, Aves, Edited by W.B. King. 2nd revised edition. Morges.
- IUCN Red Data Book, Volume 3, Reptilia and Amphibia, compiled by R.E. Honegger. Morges.
- IUCN World Conservation Strategy. Gland, Switzerland.
- Ivanov. Avifauma des russischen Pamir Alai. Izdatelstvo Nauka. Leningrad. 1969
- Iven, H.E. Das Klima von Kabul (The Climate of Kabul). Breslau: F. Hirt, pp. 74-1924
- Jaffri, S.K.H. Taxonomic study on the <u>Capparidaceae</u> and <u>Cuciferae</u> of West Pakistan, 1954 Afghanistan and the Northwest Rimalaya. Edinburgh, Thesis.
- Janata, A. Die landwirtschaftliche Struktur Afghanistans. Bustan. 4: 36-40. (The economic 1963 structure of Afghanistan)
- Jentsch, Ch. Das Nomadentum in Afghanistan. Afghanische Studien. I, 236 p. (The nomadism 1973 in Afghanistan)
- Johnstone, C. Appendix 'D' Birds in No Tigers in the Hindu Kush by Transter, P. London. 1868
- Jones, S. Nuristan: Mountain Communities in the Hindu Kush. Afghan Studies. 1: 79-92.

- Jones, S. An annotated bibliography of Nuristan and the Kalash Kafirs of Chitral.

 1966 Munksgaard, Kobenhavn. 110 p.
- Jux, U. Frasne Riffbildner im Devon der Daste Nauwar (Afghanistan). N. Jb. geol. Paläont. 1969 11: 680-690. Stuttgart. (Reef builders in the Devon of Dashte Nawar).
- Jux, U. & Kempf. Stauseen durch Travertinabsatz im zentralen Afghanischen Hochgebrirge.

 1971 Z. Geomorph. Suppl. 12: 107-137. (Formation of reservoirs by travertin deposits in high mountains of central Afghanistan).
- Kästner, H. Bibliografie zur Geologie Afghanistans und ummittelbar angrenzender Gebiete 1971 Bull. Afgh. Geol. Miner. Survey. 6: 43 p.
- Kasy, F. Österreichische entomologische Expeditionen nach Persien und Afghanistan. Amln. 1965 naturh. Mus. Wien. 68: 653-666. (Austrian entomological expedition to Persia and Afghanistan)
- Kear, J. & Duplaix-Hall, N. Flamingos. The Wildfowl Trust. T and A.D. Poyser, Berkhamp-1975 sted, England.
- Kerstan, G. Die Waldverteilung und Verbreitung der Baumarten in Ost-Afghanistan und in 1937 Chitral. (The composition and distribution of treespecies in East Afghanistan and in Chitral) in: Berichte der Deutschen Hindukusch. Schr. dt. Forschungsgem. 141-167 pp.
- Keshtyar, M. Status of pasture development and management, vegetation survey and mapping in Afghanistan. C.A.Z.R.I. Jodhpur.
- Khan, M.I.R. & Beg, A.R. A phytogeographical exoursion in Afghanistan. Pak. Journ. For. 1968 18: 287-300.
- Kilbertus, C. & Pierre, J.F. Etude hydrologique, microbiologique et algologique des eaux 1970 du Kaboul (Afghanistan). Bull. Acad. Soc. Lorraines Sci. 9(3): 433-448 (A hydrological, microbiological and algae study of the waters of Kabul, Afghanistan)
- Kitamura, S. New species from Afghanistan collected by the Kyoto University Expedition, 1956-58 1955. Acta phytotax. geobot., Kyoto. 16: 131-140; 17: 33-44, 131-142
- Kitamura, S. Flora of Afghanistan. The Committee of the Kyoto University Scientific

 1960 Expedition to the Karakoram and the Hindu Kush, Vol. 2: 486. Kyoto University,

 Japan. Complement published in 1965.
- Kitamura, S. New species from West Pakistan and Afghanistan. Acta Phytotax. Geobot. 1963 19(4): 97-106.
- Kitamura, S. Plants of West Pakistan and Afghanistan. Results of the Kyoto University 1964 Expedition to the Karakoram and Hindu Kush. 3: 1-283.
- Klapperioh, J. Auf Forschungsreisen in Afghanistan. Ent. Bl. Biol. Syst. Käfer. 501 p. 1954 (On expeditions in Afghanistan)

...

Klemmer, K. Liste der rezenten Giftschlange: Marburg, Behringwerke. (List of recent 1963 poisonous snakes)

Klimburg, M. Afghanistan. Wien. 313 p. 1966

Klookenhoff, H. & Madel, G. Uber die Flamingos (Phoenic opterus ruber roseus) der ashte 1970 Nawar in Afghanistan. J. Orn., Lpz. 111: 78-84 (On the flamingos of Dashte Nawar in Afghanistan)

Koeie, M. & Reohinger, K.H. ed. with various authors. Symbolae Afghanicae. Siol. Skr. 1954-65 I 8(1): 1-80 (1954); II 8(2): 1-215 (1955); III 9(3): 1-208 (1957; IV 10(3): 1-191 (1958); V 13(4): 1-267 (1963); VI 14(4): 1-94 (1965)

Koelz, W. New birds from Asia, chiefly from India. Proc. Biol. Soc. Washington. 52: 61-82.

Koelz, W. A new rosefinoh from Afghanistan. Auk. 66: 209. 1949

Koelz, W. New subspecies of birds from Southwestern Asia. Am. Mus. Novit. 1452, 10 p. 1950

Koelz, W. New birds from Iran, Afghanistan and India. Contr. Inst. Reg. Explor. 1: 1-32.

Komroff, E. The travels of Marco Polo. Random House, New York. 1926 (ed)

Koning, F.J. Siberische witte Kraanvegels in Afghanistan. Vogeljaar. 1: 4-8 (Siberian 1972 cranes in Afghanistan)

Koning, F.J. & Dyksen, L.J. IWRB Mission to Pakistan and Afghanistan. Mimeogr. 1971

Koning, F.J. & Dyksen, L.J. IWRB - Mission to Afghanistan. Mimeogr. 1972

Koning, F.J. & Rooth, J. Greater flamingos in Asia. Chapter 4 in Flamingos. Kear, J. 1975 and Duplaix-Hall, N. (eds.). T. and A.D. Poyser, Berkhamsted, England.

Koning, F.J. & Walmsley, J.G. IWRB Mission to Afghanistan, February 1972. IMRB Bull. 1972 33: 39-41.

Kral, B. Notes on the herpetofauna of certain provinces of Afghanistan. Zool. Listy. 1969 18(1): 55-66.

Kraus, W. Afghanistan, Natur, Geschichte und Kultur, Staat, Gesellschaft und Wirtschaft.

1972 Tubingen. (Afghanistan, nature, history and culture, country, population and economy)

Rhinolophus lepidus Blyth

R. 1. monticola Andersen, 1905. Oriental species of eastern Afghanistan. Records from Jalalabad, Laghman, Parwan and Zabul (JG).

Family Hipposideridae

- Hipposideros fulvus Gray leaf-nosed bat
 - H. f. pallidus Andersen. Recorded from Laghman (JN) and Darunta in the Jalalabad Valley III/67 (JG).
- 2. Asellia tridens (Geoffroy) trident leaf-nosed bat A. t. tridens. Records from Seistan (VA), Dilaram, Farah and Kandahar (JG).

Family Vespertilionidae

Myotis blythi (Tomes)

lesser mouse-eared bat

M. b. oxygnathus (Monticelli). Scrub forest areas up to 2 700 m. Records from Ghazni, Nuristan (Baschgar Valley), Kabul, Maimanah, Qalat, and Beltchiragh (JG), Baschgar Valley (KZ).

2. Myotis emarginatus (Geoffroy)

notch-eared bat

- M. e. desertorum (Dobson). More arid regions. Records from Kabul (JG).
- Myotis formosus (Hodgson) 3. Hodgson's bat.

- M. f. formosus. Oriental species. Recorded in Kalat-us-Seraj (E) $\overline{V}I$, $\overline{V}I\overline{I}I/\overline{X}/65$ (MO):
- 4. Myotis longipes (Dobson) long-fingered bat Oriental species. Recorded from caves and tunnels in Jalalabad 650 m VII/65 and Pialeh Cave (E) 1 550 m VIII/61 (MO).
- 5. Myotis mystacinus (Kuhl) Whiskered bat M. m. meinertzhageni Thomas. One record from Kabul (JG).
- Pipistrellus babu Thomas Himalayan pipistrelle Oriental species. Records from Jalalabad 650 m and Kalat-us-Seraj (E) 900 m X/65 (MO).
- 7. Pipestrellus coromandra (Gray) Indian pipistrelle P. c. coromandra. Quite plentiful in the Kabul Valley near human habitations in Jalalabad III/65, III, IV/67, Dar-i-Nur (Kunar Valley) 2 000 m III/67 (JG).
- 8. Pipistrellus kuhli (Kuhl) Kuhl's pipistrelle. Steppe areas. Records from Kandahar, Dilaram (S), Khost (E) (MO), Jalalabad IV, V/67 (JG), Kandahar (Blyth, 1845).
- 9. Pipistrellus mimus Wroughton Indian pygmy pipistrelle P. m. glaucillus Wroughton. Oriental species of the wooded regions. Recorded from Kala-i-Shahi (Nangarhar) 850 m VIII/61 (MO).
- Pipistrellus pipistrellus (Schreber) common pipistrelle P. p. aladdin Thomas. Human habitations, northern Afghanistan. Kunduz IX/65, V/67 and Baghlan V/67 (JC).
 - P. p. pipistrellus. Human habitations, eastern Afghanistan. Records from Jalalabad IV/67 (JG, MO), Baschgar Valley (Nuristan) VII/53 (KZ).

Kuhnert, G. Vogelhaltung in Afghanistan. Gefied. Welt. 9: 173-176. (Birdkeeping in 1971 Afghanistan)

Kullmann, E. Die Säugetiere Afghanistans. I. Carnivora, Artiodaotyla, Primates. Soience, 1965 Kabul. 1-17 pp. (The mammals of Afghanistan)

Kullmann, E. Über Leoparden Afghanistans und ihre Parasiten. Freunde Köln Zoo. (On the 1967 leopards in Afghanistan and their parasites)

Kullmann, E. Expedition in die Heimat der Marco Polo Schafe. Freunde Edln. Zoo. 11: 107-1968 122. (Expedition into the homeland of the Marco Polo sheep)

Kullmann, E. Die Tierwelt Ostafghanistans in ihren geographischen Beziehungen. Freunde 1970 Köln. Zoo. 13(1): 3-25. (The fauna of eastern Afghanistan from a geographical aspect)

Kumler, M.L. Plant distribution on hills surrounding Kabul, Afghanistan. Transact. Illin. 1969 State Acad. Sci. 62(2): 141-153.

Kussmaul, F. Badaxsan und seine Tagiken. Tribus. 14: 11-99. (Badakhohan and its Tagiks) 1965

Kuz'akin, A.P. Letuoije mysi (bats). Moscow (in Russian) 1950

Kyoto University. Result of the Kyoto University Scientific Expedition to the Karakorum 1960-65 and Hindu Kush. 5 Vols. Kyoto, Japan.

Lalande, P. Généralités sur la végétation du Safed-Koh et de son prolongement occidental.

1967

Bull. Soc. Hist. Nat. Toulouse. 103: 297-304. (Generalities on the vegetation of Safed-Koh and its western extension)

Lalande, P. Observations sur qualques arbres: afghans. Bull. Soc. Hist. Nat. Toulouse. 1968a 104: 131-137. (Observations on some trees of Afghanistan)

Lalande, P. Btages à genévriers en Afghanistan. Compt. rend. Acad. Sc. Paris. 267D: 503-1968b 504. (The juiper formation in Afghanistan)

Lalande, P. Généralités sur la végétation du bassin du Kaboul en Afghanistan. Trav. Lab.

1968o for. Toulouse. V(I,3 art. III): 1-17. (Generalities on the vegetation of the Kabul basin in Afghanistan)

Lalande, P., Essone Selemani & Parenda, K. Report on the Mission done to Registan and to 1975 Seistan from 29.3.75 to 7.4.75. Mimeogr.

Lalande, P., Herman, N.M. & Zillhardt, J.Z. Cartes climatiques de l'Afghanistan. 2 Vols: 1973-74 texte, cartes. Aviation Civil et Tourisme, Publications de l'Institute de Meteorologie, No. 4, Kabul. (Climatological maps of Afghanistan)

Lenge, de la Camp, N. Die Weisen der Deutschen Hindukusch Expedition, 1935. Lendwirtsch.

1939 Jahrb. 88: 14-133. (The wheat species of the German Hind Kush Expedition, 1935)

Lapparent, A.F. Paysages biogeographiques de l'Afghanistan. Compt. rend. Soc. Biogeograph.

1970 No. 412 (non-publie) (Biogeographical landscapes of Afghanistan)

Lapparent, A.F., de Lavegne Sainte Lausanne, Z. & Bordet, P. Sur l'importance et l'extension 1965 du volcanisme récent de Nawar (Afghanistan). Bull. volcan. 28: 107-118, 4 abb, 3 tables. (On the importance and the extension of the recent volcanism of Nawar)

Lay, D.M. A study of the mammals of Iran resulting from the Street expedition of 1962-63.

1967 Fieldiana, Zool. 54: 1-282.

Lay, D.M., Anderson, J.A.W. & Hassinger, J.D. New records of small mammals from W. Pakistan 1970 and Iran. Mammalia. 34(1): 98-106.

Leviton, A.E. Report on a collection of reptiles from Afghanistan. Proc. Calif. Acad. 1959 Soi. 29/12, 445-463.

Leviton, A.E. & Anderson, S.A. Further remarks on the amphibians and reptiles of Afghanistan. 1961a Wasmann, J. of Biology. 19: 269-276.

Leviton, A.E. & Anderson, S.A. Third contribution to the herpetology of Afghanistan. Proc. 1961b Calif. Acad. of Sc., 4th Ser. 31(12): 329-339.

Leviton, A.E. & Anderson, S.C. The amphibians and reptiles of Afghanistan, a checklist and 1970 key to the herpetofauma. Proc. Calif. Acad. Sci., 4th Ser. 38(10): 163-206.

Lincheviski, I.A. The principal regularities in vegetation distribution of Afghanistan. 1946 Bot. Zh. Kfyiv. 183-218 pp.

Linchevski, I.A. & Prozorovski, A.V. The vegetation of Afghanistan. Bot. Zhurn. 29: 114-1944 123 (in Russian).

Linchevski, I.A. & Prozorovsky, A.V. Basic principles of the distribution of the vegeta-1949 tion of Afghanistan. Kew Bull. 1949(2): 179-214.

Lindberg, K. Le lao dit de Chiva (Afghanistan) (Le pays des Choghnis et des marmottes).

1961 Aota geogr., Paris. 37: 4-10, 17.

Lindberg, K. Recherches biospéleologiques en Afghanistan. Acta Univ. lund. 57(1), 58(2). 1962 (Biospeleogolical research in Afghanistan)

Lindsay, H.N. Indian shrews. Scientific results from the mammal survey no. 48. J. Bombay 1929 nat. Hist. Soc. 33: 326-340.

18hrl, H. & Thielcke, C. Zur Brutbiologie, Okologie und Systematik einiger Waldwögel
1969 Afghanistans. Bonn. zool. Beitr. 20: 85-98. (On the breeding biology,
ecology and systematics of some forest birds of Afghanistan)

Lumsden, H.B. The mission to Kandahar. Caloutta. 1860

Lydekker, R. Wild oxen, sheep and goats of all lands. Rowland Ward, London. 1898

Madel, G. Hummelbeobschtungen im Wakhan. Soience (Kabul) 5: 21-28. (Observations on 1968 the bumble-bees in Wakhan)

Narien, D. Notes from the Walter Koelz Collections. V: Notes on some Asiatic Meropidae,
1950-51

J. Bombay nat. Hist. Soc. 49: 151-164; VII: Notes on the bird family Prunellidae
in Southern Eurasia, Am. Nus. Novit. 1428, 28 p.; IX: Notes on some Asiatic
Sturnidae, J. Bombay nat. Hist. Soc. 49: 471-487; X: Notes on some pheasants
from Southwestern Asia, with remarks on moult. Am. Nus. Novit. 1518, 25 p.

McArthur, I.D. & Harrington, G.N. A grazing ecosystem in Western Afghanistan. Proc. 1st 1978 Internat. Rangelands Congress, Denver, Colorado.

McArthur, I.D., Sayed, S. & Nawim, M. Rangeland livestock production in Western Afghanistan.

1979 J. Arid Environments. 2: 163-179.

McClure, B. A review of range research in Afghanistan. Kabul, mimeo, 43 p. 1979

McClure, M.E. Migration and survival of the birds of Asia. US Army Component, Seata, 1974 Bangkok, Thailand.

McDonald, W. Marco Polo Sheep. Karatasi Yanye Habari. 15(1): 10-14.

Meher-Homji, V.N., Gupta, R.K. & Freitag, H. Bibliography on "Plant Ecology" in Afghanistan.

1973 Excerpta Botanica Sectio B Sociologica. 12: 310-315.

Meinertzhagen, R. On the birds of Northern Afghanistan. Ibis. XIV(2): 480-520, 671-717.

Meinertzhagen, R. Notes on Afghan Birds. Ibis XV: 347. 1939

Meklenburtzev, R.N. The Arkhar of the Pamir Ovis poloi poloi Blyth. Biology and cyto-1948 genetical-economical importance. Bull. Sc. Hist. Nat. de Moscow. 53(5): 65-84.

Meyer-Oehme, D. Die Säugetiere Afghanistans (Teil III). Science (Kabul) 1965: 42-56.

1965 (The mammals of Afghanistan)

Meyer-Oehme, D. Zur Kenntnis der Chiropteren-Fauna Afghanistan. Bonn. zool. Beitr. 1968 19(1/2): 97-103. (On the knowledge of the bat-fauna of Afghanistan)

Nichels, A.A. The Kabul, Kunduz and Helmand valleys and the national economy of Afghanistan.

1959 Nat. Acad. Sci., National Research Council, Washington.

Mirwald, P. & Roemer, H. Beobachtungen im Wakhan. Erdkunde. 21: 48-57. (Observations in Wakhan 1967

Misonne, X. Analyse Zoogsographique des Mammiféres de l'Iran. Ném. Inst. r. Soi. nat. 1959 Belg. Nem-Deux, Ser., 59 p. (Zoogsographical analysis of the mammals of Iran)

- Misra, S.D. Natural regions of Afghanistan. J.G. (Stellenbosch) 3(1): 27-31.
- Morden, W. Across Asia's snows and deserts. Putmans, London. 414 pp. 1927
- Murray, J.A. The zoology of Balu chistan and Southern Afghanistan. Bombay Education Society Press.
- Murray, J.A. The zoology of Baluchistan and Southern Afghanistan. Ann. Mag. nat. Soi., 1887 India. 1(1-3): 49-68, 105-131.
- Kuss, H. Über Flora und Vegetation des Wakhan und 'Grossen Pamir'. in Grosser Pamir,

 1978 Akademische. Druck u. Verlagsanstalt Graz. (On the flora and vegetation of
 Wakhan and the Big Pamir)
- Musson, C. Narrative of various journeys in Baluchistan, Afghanistan and the Punjab.
- Nadler, C.F., Korobitsina, K.V., Hoffmann, R.S. & Vorontsov, N.N. Cytogenetic differentiation, geographic distribution and domestication in Palaearctic sheep (Ovis). Z. Sängetierk. 38: 109-125.
- Nadler, C.F., Lay, D.M. & Hassinger, J.D. Cytogenetic analysis of wild sheep populations in Northern Iran. 10: 137-152.
- Nalivkin, I.D. Obzor geologii Pamira i Badakchana. Moscow-Leningrad: Transactions of 1932 the United Geological and Prospecting Service of the USSR. 104 (Geological survey of the Pamirs and Badakhshan)
- Nasimovitch, A.A. The role of the snow cover regime in the life of the ungulates of the 1955 USSR (in Russian). Izv. imp. Ahad. Nauk. 12: 681-710.
- Nasimovich, A.A. Rare and endangered species of mammals and their protection in the USSR.

 1970 Engl. transl. Henry Field (ed.). Field Research Projects 16, Kiami, Florida.

 10 p.
- Naumann, C. Ein ehemaliges Wildyak-Vorkommen im Afghanischen Pamir. Bonn. zool. Beitr. 1973 24: 254-269. (Former occurrence of wild yaks in the Afghan Pamir)
- Naumann, C. Tentative checklist of Afghanistan's birds. Nimec. 21 pp. 1974

THERESENDED

- Naumann, C.M. Zur Tierwelt des 'Grossen Pamir'. in: Grosser Pamir, Akademische Druck 1978 u. Verlagsanstalt Graz. (On the fauma of the Big Pamir)
- Naumann, C. & Niethammer, G. Zur Säugetierfauna des Afghanischen Pamir und des Wakhan.

 1973 Bonn. zool. Beitr. 21: 237-248. (On the mammal fauna of the Afghan Pamir and Wakhan)

- Naumann, C. & Niethammer, J. Neumachweise von Säugetieren aus dem nördlichen Afghanistan.

 1975 Säugetierkundliche Kitt. 22(4): 295-298. (New mammal species from northern Afghanistan)
- Naumann, C. & Nogge, C. Die Grossäuger Afghanistans. Z. Kölner Zoo. 16:79-93.

 (The large mammals of Afghanistan)
- Nauroz, M.K. & Naumann, C.M. Bemerkungen zur Verbreitung des Markhors, Capra falooneri
 1975 (Wagner, 1839) in Afghanistan. Säugetierkundliche Mitt. 23(2) 81-85. (Remarks
 on the distribution of the markhor in Afghanistan)
- Nedyalkov, S.T. Ecological areas of the forest tree and bush vegetation in Afghanistan. 1978a Corskostopenska Nanka (Sofia) 15(2): 71-89.
- Nedyalkov, S.T. Ecological areas of forest tree and shrub vegetation in Afghanistan. 1978b Abstract No. 1401, 40(5) 1979.
- Neubauer, H.F. Die Pflanzengesellschaften Afghanistans. Ber. an das Kgl: Afghan.Ministerium 1951 für Landwirtschaft vom 19.10.1951. (The plant communities of Afghanistan)
- Neubauer, H.F. Die Hälder Afghanistans. Angew. Pfl. Soziol. Festschr. E. Aichinger. 1954a 1: 494-503. (The forests of Afghanistan)
- Neubauer, H.F. Versuch einer Kennzeichnung der Vegetationsverhältnisse Afghanistans.

 1954b AnnIn. naturh. Mus. Wien. 60: 77-113. (Attempt on the characterisation of the relationships in vegetation in Afghanistan)
- Neubauer, H.F.: Bemerkungen über das Vorkommen wilder Obstsorten in Nuristan. Angew. Bot. 1954c 28: 81-88. (Remarks on the existence of primitive fruit varieties in Muristan.)
- Newhauser, H.N. & De Blase, A. Notes on bats new to the faunal lists of Afghanistan and 1974 Iran. Fieldiana, Zool. 62(5): 85-96.
- Nicolson, N. The Himalayas. Time Life Books, Amsterdam. 1975
- Niethammer, G. Über die Kehltaschen des Rotflügelgimpels, Rhodopechys sanguinea. J. Orn., 1966 Lpz. 107(3)(4): 278-282. (On the throat sack of the crimson-winged finch (Rhodopechys sanguinea))
- Niethammer, G. On the breeding biology of <u>Montifringilla theresae</u>. Ibis 109: 117-118.
- Niethammer, G. Königshühner. Vogel-Kosmos, Stuttgart. 3: 86-90 (Phasianidae: Tetrao-1968 gallus)
- Niethammer, G. Vogelleben am Ab-i-Estada (Afghanistan). Vogelwarte. 26: 221-227. 1971 (Bird life at the Ab-i-Estada (Afghanistan)).
- Niethammer, G. Störche über Afghanistan. Z. Köln. Zoo. 15(2): 47-54. (Storks over 1972 Afghanistan)

- Niethammer, G. Zur Vogelwelt des Afghanistan Pamir und des Darwaz. Bonn. zool. Beitr. 1973 24: 270-284. (On the birds of Afghan Pamir and Darwaz)
- Niethammer, G. & J. Neumachweis für Afghanistans Vogelwelt. J. Orn., Lpz. 108: 76-80. 1967a (New records of Afghan birds)
- Niethammer, G. & J. Zur Variabilität der Kehlzeichnung beim Steinmarder, Martes foina 1967b (Erxleben, 1777). Z.f. Säugetierkunde. 32(3): 185-187. (On the variability of the throat marking of the stone marten)
- Niethammer, G. & J. Hoohgebirgs-Vogelzug in Afghanistan. Bonn. zool. Beitr. 18: 501-507.

 1967c (Bird migration in the high mountains of Afghanistan)
- Niethammer, J. Die Säugetiere Afghanistans (Teil II) Insectivora Lagomorpha, Rodentia.

 1965 Science, Kabul. 1965: 18-41. (The mammals of Afghanistan (Part II): Insectivora, Lagomorpha, Rodentia)
- Niethammer, J. Der Zug von Kranichen (<u>Gruidae</u>) bei Kabul, Afghanistan. Vogelwarte. 1966a 23: 308-309. (The migration of cranes (<u>Gruidae</u>) near Kabul)
- Niethammer, J. Zur Ernährung des Sumpfluchses, Felis chaus, in Afghanistan. Z. Säugetierk. 1966b 31: 393-394. (Feeding habits of the Jungle-Cat, Felis chaus, in Afghanistan)
- Niethammer, J. Störche in Afghanistan. Vogelwarte. 24: 42-44. (Storks in Afghanistan) 1967a
- Niethammer, J. Zwei Jahre Vogelbeachtungen an stehenden Gewässern bei Kabul in Afghanistan.

 1967b J. Orno. Lpz. 108: 119-164. (Two years of bird watching at standing waters near Kabul, Afghanistan)
- Niethammer, J. Ein Pfeifhase, Ochotona rufescens (Gray, 1842), ohne Stiftzähne. Säugetierk.

 1967c Hitt. 16(2): 160-162. (An Afghan pika without front teeth)
- Niethammer, J. Die Flughörnchen (Petauristinae) Afghanistans. Bonn. Zool. Beitr. 1/2:14. 1967d 18: 1-14. (The flying squirrels (Petauristinae) of Afghanistan)
- Niethammer, J. Pelztierfelle im Basar von Kabul, Afghanistan. Pelzgewerbe. 18(1): 7-9.
 1967e (Furs in the bazar of Kabul, Afghanistan)
- Niethammer, J. Zur Taxonomie der Ohrenigel in Afghanistan (Gattung Hemieohinus).

 1969a Z. Sängetierk. 34(5): 257-274. (On the taxonomy of the hedgehogs in Afghanistan (genus Hemieohinus))
- Niethammer, J. Die Waldmaus, Apodemus sylvaticus, in Afghanistan. Säugetierk. Mitt. 1969b 17: 121-128. (The wood mouse, Apodemus sylvaticus in Afghanistan)
- Niethammer, J. Die Flamingos am Ab-i-Estada in Afghanistan. Natur. Nus. Frankf. 100: 1970a 201-210. (Flamingos at the Ab-i-Estada in Afghanistan).
- Niethammer, J. Die Wühlmäuse (Microtinae) Afghanistans. Bonn. Zool. Beitr. 21: 1-24.
 1970b (The voles and lemmings (Microtinae) of Afghanistan)

- Niethammer, J. Das Mauswiesel (Mustela nivalis) in Afghanistan. Bonn. Zool. Beitr. 1973a 24: 1-6. (The weasel (Mustela nivalis) in Afghanistan)
- Niethammer, J. Zur Kenntnis der Igel (Erinaceidae) Afghanistans. Z. Säugetierk. 28(5): 1973b 271-276. (The hedgehogs (Erinaceidae) of Afghanistan)
- Niethammer, J. & Martens, J. Die Gattungen Rattus und Maxomys in Afghanistan und Nepal.

 1975 Z. Säugetierk. 40(6): 325-355. (The genera Rattus and Maxomys in Afghanistan and Nepal)
- Nogge, C. Afghanistan the Ab-i-Estada: A vanishing breeding place of flamingos. I.W.R.B. 1969 Bull. 31: 28-30.
- Nogge, G. Afghanistan's unique bird paradise. Kabul Times. p. 2.
- Nogge, C. Observations of breeding places of flamingos in Afghanistan. I.W.R.B. Bull. 1972b 34: 30-39.
- Nogge, G. Kabul Zoo. Outdoorman. 3: 82-36. Karachi. 1972c
- Nogge, G. Ornithologische Beobachtungen im Afghanischen Pamir. Bonn. Zool. Beitr. 1973a 24: 254-269. (Ornithological observations in the Afghan Pamir)
- Nogge, G. Vogeljagd im Hindukusch. Natur. Mus. Frankf. 103: 276-279. (Bird hunting in 1973b the Hindu Kush)
- Nogge, G. Beobachtungen an den Flamingobrutplätzen Afghanistans. J. Orn. Lpz. 115: 142-151. 1974a (Observations on the flamingo breeding places of Afghanistan)
- Nogge, C. Sieben Jahre Kabul Zoo Afghanistan. Z. Köln. Zoo. 17: 105-109. (Seven years in 1974b Kabul Zoo)
- Nogge, G. & Niethammer, J. Die Vögel in den Basars von Kabul und Charikar. Afghanistan J. 1: 151-154. (The birds in the Kabul and Charikar markets)
- Novikov, G.A. Carnivopous mammals of the fauna of the USSR. Translated by Israel 1962 Programme for Scientific Translation. Jerusalem.
- Nuri, J.N. Beiträge zur Kenntnis der Steppenpflanzen Afghanistans. Die morphologische

 1961 Plastizität von Leotuca orientalis und Artemisia fortiana. Diss. Würzburg,
 63 pp. (Contribution to the knowledge of the steppeplants of Afghanistan. The
 morphological plasticity of Leotuca orientalis and Artemisia fortiana)
- Oedekoven, K. Forest history of the Near East. Unasylva, 17: 13-21. 1964.

- Ognev, S.I. Mammals of the USSR and adjacent countries, in 9 vols. Israel Programme for 1962-67 Scientific Translation (1928-1950, in Russian). Vol. I Inscotivora and Chiroptera, 1962; Vol. 2 Carnivora (Fissipedia), 1962; Vol. 3 Carnivora (Fissipedia and Pinnipedia), 1962; Vol. 4 Rodents (Duplucidentata and Pinnipedia), 1966; Vol. 5 Rodents (Sciuridae, Marmota), 1963; Vol. 6 Rodents (Dipodidae), 1963; Vol. 7 Rodents (Microtines), 1964.
- Olufsen, 0. Through the unknown Pamirs. Heinemann, London, 238 pp. 1904
- Orkel, C. Ist Rissa tridactyla für Afghanistan wirklich nachgewiesen? J. Orn., Lpz. 113: 447-448. (Is Rissa tridactyla really observed in Afghanistan?)
- Pabot, H. Les grandes régions phytogéographiques et écologiques de l'Iran. Coll. Conserv.

 1961 Restaur. Sols. Lutte contre l'érosion. 340-349 pp. (The great vegetation
 and ecological regions of Iran)
- Paludan, K. On the birds of Afghanistan. Vidensk. Meddr. dansk naturh. Foren. 122: 1-332...
- Paludan, K. Notes on the Afghan mammals. Made in 1948-49 while participating in 3rd Danish 1948-49 Expedition to C. Asia, unpubl.
- Pelt, J.M., Hayon, J.C. & Younos, M.C. Plantes médicinales et drogues de l'Afghanistan.

 1965 Bull. Soc. Pharm. Nancy. 66: 16-61. (Medicinal and drug plants of Afghanistan)
- Pelt, J.K., Hayon, J.C. & Younos, M.C. Sur la flore et la végétation d'une zone halophile 1968a steppique en bordure de l'Amou-Darya (Afghanistan). Compt. rend. Acad. Sc. Paris. 267D: 505-508. (On the flora and vegetation of a halophytic steppe zone along the Amu Darya (Afghanistan))
- Pelt, J.H., Hayon, J.C. & Younos, N.C. Sur la flore et la végétation des bords du lac 1968b Ab-i-Estada. Compt. rend. Acad. Sc. Paris. 267D: 1279-1282. (On the flora and vegetation of the banks of the Ab-i-Estada lake)
- Pelt, J.M., Hayon, J.C. & Younos, M.C. Caractères écologiques et floristiques de deux 1969 halophiles d'Afghanistan. Vegetatio. 20: 307-328. (Ecological and floristical characteristics of two halophyllics of Afghanistan)
- Pelt, J.M., Hayon, J.C. & Younos, M.C. La végétation d'une cuvette halophile du bassin de 1970a Bamian (Afghanistan Central). Compt, rend. Acad. Sci. Paris. 271D: 320-323. (The vegetation of a halophil basin in the Bamian Basin (Central Afghanistan))
- Pelt, J.M., Hayon, J.C. & Younos, N.C. La végétation de la Valléé d'Madjar (Afghanistan 1970b Central). Bull. Soc. bot. France. 117: 297-306. (The vegetation of the Ajar Valley (Central Afghanistan))
- Perrin, N. L'Afghanistan, ou description géographique, du pays théatre de la guerre,
 1842 accompagnée de détails sur les tribus de contrées, leurs moeurs, leurs usages,
 etc., etc. Paris. (Afghanistan, or a geographical description of a country which
 is a theatre of war, accompanied by details of the tribes, their habits, etc.)

- Peterson, R., Mountfort, G. & Hollom, P.A.D. A fieldguide to the birds of Britain and 1974 Europe. London, Collins, 344 p.
- Petooz, R.G. The exploration and conservation of wild ungulates and their habitat in the 1971 Afghan Pamir. Kimeograph, Kabul, 17 p.
- Petocz, R.G. Report of the Laghman markhor survey. Afghan Tourist Organization, 17 p. 1972 .
- Petocz, R.G. The Bactrain deer (Cervus elaphus bactrianus). A report of the March 1973

 field survey in Northern Afghanistan. Kabul, Afghan Tourist Organization, Mimeo,

 8 p.
- Petocz, R.G. Kabul markhor (<u>Capra falconeri megaceros</u>) and urial (<u>Ovis orientalis oycloceros</u>)

 1973b in the Kohe Safi region of Kapsia Province: report of the April 1973 field survey.

 Afghan Tourist Organization, Kabul. Mimeo, 8 p.
- Petocz, R.C. Marco Polo sheep (Ovis ammon poloi) of the Afghan Pamirs: a report of biolo-1973c gical investigations in 1972-73. ATO Report, 26 pp. Kabul.
- Petocz, R.G. Reorganization and improvement of the Maroo Polo sheep hunting programme.

 1973d Afghan Tourist Organization. Mimeo, 17 p.
- Petrak, F. Beiträge zur Pilzflora von Afghanistan. Sydowia. 7: 64-97; 16: 331-349. (Contributions to the fungus-flora of Afghanistan)
- Petter, F. Eléments d'une révision des lièvres européens et asiatiques du sous-genre Lepus.

 1961a Z. Säugetierk. 26: 1-11. (Elements for a revision of the European and Asian hares of the sub-genus Lepus)
- Petter, F. Répartition géographique et écologie des rongeurs désertique de la région 1961b paléarctique. Mammalia 25, Special nr., 1-222. (Geographical and ecological distribution of desert rodents in the Palaearctic region)
- Pfeffer, P. Le mouflon de Corse (Ovis ammon musimon Sohreber, 1782); position systématique, écologie et ethologie comparées. Mammalia 31, suppl. 262 p. (The mouflon of Corsica (Ovis ammon musimon Sohreber, 1782); systematic position, ecology and behaviour compared)
- Pias, J. Formations superficielles et sols d'Afghanistan. Office de la Recherche Scien-1976 tifique et Technique d'Outre-Ner, France, Paris. (Superficial formations and soils)
- Pickett, L.C. Ma-Yar, M.Q. & Saleh, Z. Bibliography of materials dealing with agriculture in Afghanistan. Technical Bull. No. 8 Fac. of Agric. Kabul Univ. 316 p.
- Pilger, A. Die zeitlich-tektonische Entwicklung der Iranischen Gebirge. Clausthaler Geol.
 1971 Abh. 8: 1-27. (The historic tectonic development of the Iranian mountains)
- Pilger, A. & Rösler, A. The contemporaneous tectonic events of the Indian Ocean and neighbouring areas. Abh. Braunschweigischen wissensch. Gesell. 26: 67-98.

- Pimms, S.L. Report on the birds seen on the Hindu Kush agricultural and ecological project.

 1968 St. Catherine's College, Oxford. 11 p. ron.
- Pocook, R.I. Two local races of the Asiatic wild ass. J. Bombay nat. Hist. Soc. 47(2): 1947 143-144.
- Pocook, R.I. Catalogue of the genus <u>Felis</u>. London, 190 pp, 8 figs. 1951
- Podlech, D. Several articles on new plant species in Afghanistan. Mitt. Bot. Staatssamml. 1967-69 Munchen. 6: 547-591; 7: 101-117; 6: 541-546; 7 and 8.
- Poelt, J. & Wirth, V. Flechten aus dem nordöstlichen Afghanistan gesammelt von H. Roemer 1968 im Rahmen der Deutschen Wakhan-Expedition 1964. Mitt. Bot. Staatssamml. München. 7: 219-261. (Michens from northeastern Afghanistan, collected by H. Roemer in connection with the German Makhan Expedition of 1964)
- Poncins, E.D. Shooting Ovis (sic) poloi on the Pamirs. J. Bombay nat. Hist. Soc. 10: 53-1895 62.
- Potapov, R.L. On the biology of the Pamir Leucosticte (<u>Leucostichte arctoa pamirensis</u>). 1963 Zool. Zh. 42: 716-727.
- Povolny, D. The discovery of the bear <u>Selenarctos</u> thibetanus in Afghanistan. Zool. Listy. 1966 15(4): 305-316.
- Povolny, D. Beiträge zur Kenntnis der Fauna Afghanistan. Reisebericht um Charakteristik des Sammelgebietes. Cas. morav. zemsk. Eus. 52(suppl.): 12-14, 21-29. (Contribution to the knowledge of the Afghan fauna)
- Prater, S.H. The book of Indian animals. 2 ed. Bombay Natural History Society. 1965
- Pritchard, P.C. Encyclopedia of turtles. T.F.H. Publications. Inc. Ltd. 1979
- Puget, A. Contribution à l'étude des oiseaux du nord-est de l'Afghanistan. Thèses Toulouse
 1969 Univ. p. 487. (Contributions to the study of the birds of northeastern Afghanistan)
- Puget, A. <u>Tetraogallus himalayensis himalayensis</u> Gray in Afghanistan. Alauda. 38: 169-173. 1970a
- Puget, A. Observations sur les Psittacides vivant en Afghanistan. Alaula. 38: 306-309. (Observations on the Psittacidae in nature in Afghanistan)
- Puget, A. Observations sur l'avifaume nidificatrice de l'étang de Binchesar (Chamane Cala 1971 Hachmatran) dans la proche banlieue de Kaboul en Afghanistan. Alauda. 39: 139-144. (Observations on the breeding avifauma of the Kole Hashmat Khan lake, in the Kabul environment, Afghanistan)

- 11. Nyctalus leisleri (Kuhl) lesser noctule Recorded from Ismael Khel (E), 1 340 m V/63 (MO).
- 12. Nyctalus montanus (Barrett-Hamilton) mountain noctule Collected in Ismael Khel (E) 1 340 m (MO) and Jalalabad III/67 (JG).
- 13. Eptesicus bottae (Peters)

 E. b. ognevi Bobrinskii. Records from Faizabad (NE) and probably
 Kabul (JG).
- 14. Eptesicus nasutus (Dobson) Sind bat E. n. nasutus. Jalalabad III, IV/67 (JG).
- 15. Eptesicus serotinus (Schreber) common serotine

 E. s. pashtomus Gaisler. Roost in fissures of buildings. Recorded from Jalalabad IV/67 (JG).

 E. s. turcomanus (Eversmann). Recorded from Faizabad (NE) (Kuzakin, 1950) and Kunduz V/67 (JG).
- 16. <u>Vespertilio murinus</u> Linnaeus. particoloured bat <u>V. m. murinus</u>. Recorded from Afghanistan (Corbett, 1978).
- 17. Otonycteris hemprichi Peters Hemprich's long-eared bat

 O. h. cinerea Satunin. Upland steppes and mountains. Record from

 Neuabad-Bagrami (Kabul Province) VII/63 (NO).
- 18. Scotophilus heathi (Horsfield)

 S. h. heathi. Oriental species, often near human habitation.

 Recorded from Jalalabad IX/64, VII/65, III, IV/67 (JG).
- 19. Barbastella leucomelas (Cretzschmar) Asian barbastelle

 B. 1. darjelingensis (Hodgson). Forested mountains. Recorded from Wazir Bagh 1 240 m (Paktya) VII/65 and Kalat-us-Seraj (E) IX/65 (MO).
- 20. Plecotus austriacus (Fischer) grey long-eared bat

 P. a. wardi Thomas. Woodlands up to 2 700 m. Recorded from Kabul
 and Bamian II/65 (JG), Kabul VI/53 (KZ), Parwan (Lindberg), Qoul
 Kharideh (75 km W of Kabul) (VA).
- 21. Miniopterus schreibersi (Kuhl) long-winged bat

 M. s. fuliginosus (Hodgson). Recorded from Jalalabad (MO), II/66,

 III, V/67 (JG).
 - $\underline{\text{M. s. pallidus Thomas.}}$ Arid mountainous regions. Chamchir Cave

Family Molossidae

- 1. Tadarida aegyptiaca (Geoffroy) Egyptian free-tailed bat
 T. a. sindiaca Wroughton. Recorded from Kabul VII/62, IX/65 (MO).
- 2. Tadarida teniotes (Rafinesque). European free-tailed bat
 T. t. teniotes. Recorded from Kabul VI/62 (MO).

- Rako, A. & Cizek. The problem of breeding and winter feeding of sheep in Kandahar, 1977 Afghanistan. Ninistry of Agriculture, Kabul.
- Rathjens, C. Verbreitung, Mitzung und Zerstörung der Wälder und Gehölzfluren in Afghanistan.

 1968-69 Jb. Südasien Inst. Univ. Heidelberg. 3: 18. (Distribution, use and destruction of the forests and treestands in Afghanistan)
- Rathjens, C. Fragen der horizontalen und vertikalen Landschaftsgliederung im Hochgebirgs1972 system des Hindu Kusch in: Geoeco of the high-mountain regions of Eurasia. 205222, Publ: Vtroll, Wiesbaden. (Questions on the horizontal and vertical landscape
 formations in the high mountain system of the Hindu Kush)
- Rathjens, C. Klimatische Jahreszeiten in Afghanistan. Afghanistan J. 1: 13-18. (Climato-1974 logical seasons in Afghanistan)
- Raven, P.M. The relationship between Mediterranean Flora in: Plant life of SW Asia.

 1971 Botanical Society of Edinburgh.
- Rechinger, K.H. Various articles on new species. Ost. Bot. Zeitschr.: 95: 422-427;
 1949 97: 221-268; 100: 437-477; 104: 173-174. Ann. Nathist. Hofmus. Wein 58: 44-61;
 66: 45-50; Bot. Jahrb. 75: 342-364; Anz. Ost. Akad. Wiss. math.-nat. Klasse
 7: 58-64.
- Rechinger, K.H. Notizen zur Orient Flora. Anz. Ost. Akad Wiss. Math.-nat. Klasse. nos. 1961-68 11-13, 1961-104-108 1968. (Notes on the oriental flora)
- Rechinger, K.H. Bericht über eine botanische Expedition nach Afghanistan. Anz. öst. Akad. 1963a Wiss. 100: 71-72. (Report on a botanical expedition to Afghanistan)
- Rechinger, K.H. Flora Iranica. Graz. 1963b
- Rechinger, K.H. & Koie, M. Symbolae Afghanicae. Biol. Skr. 1 Labiatae 8(1), 1954; 1954—63 2 Compositae 8(2), 1955; 3 Leguminosae 9(3), 1957; Cruciferae, Rubiaceae, Sorophulariaceae 10(3), 1959; 5 Div. 13(4), 1963; 6 Desor. of plants coll. by 3rd Danish Exp. to Central Asia 10(3), 1958, 191 p.
- Rechinger, K.H. & Riedl, H. Zur Flora von Nord-Ost-Afghanistan. Annln naturh. Mus. Wien. 1961-62. 65: 29-37. (On the flora of northeastern Afghanistan)
- Reeb, F. Contribution a l'étude de l'avifaume et des migrations en Afghanistan. Alauda.

 45: 293-333. (Contributions to the study of the Avifauma and the migrations in Afghanistan)
- Reiner, E. Some remarks on the knowledge of the geography of Afghanistan. C. Rev. 1962 Afghanistan. 1: 4-8.
- Reining, W.F. Beiträge sur Faunistik des Pamir-Gebietes in: Wissenschaftliche Ergebnisse 1932 der Alai - Pamir Expedition Teil III. 2 Vols., Berlin. (Contribution to the fauna of the Pamir area)

- Richter, H. Zur Systematik und Verbreitung Afghanischer Grooiduren. Zool. Abh. Staat.

 1970a Nus. Tierk. Dresden. 31(5): 269-279. (On the systematics and distribution of Afghan Grooidura species)
- Richter, H. Zur Taxonomie und Verbreiting der palaearktischen Grooiduren (Mamm., Insecti1970b vora, Sorioidae). Zool. Abh. Mus. Tierk. Dresden. 31: 293-304. (On the taxonomy
 and distribution of the palaearotic crocidura)
- Riedl. H. Bemerkungen über neue oder kritische Boraginaceen der asiatischen Flors. Ost. 1963 Bot. Zeitschr. 110: 511-542. (Remarks on new or oritical Boraginaceae of the Asiatic flora)
- Riemer, E. Botanische Ergebnisse der Deutschen Hindukush Expedition 1935. Lichenes.

 1938 Feddes Rep. Beiheft. 108: 13-25. (Botanical results of the German Hindu Kush expedition, 1935. Lichenes)
- Roberts, T. A note on Capra falconeri (Wagner, 1839) Z. Säugetierk. 34: 238-249.
- Roberts, T.J. The mammals of Pakistan. Ernest Benn Ltd. London and Tonbridge. 1977
- Robertson, G.S. The Kafirs of the Hindu Kush. London, Lawrence and Bullen. Reprinted 1896 1974, Oxford University Press.
- Robertson, G.S. Kafiristan and its people. J. Anthrop. I. 27: 75-89. 1897
- Robinson, J.A. Notes on the nomad tribes of eastern Afghanistan. Government of India 1935 Publication.
- Roemer, H. Vegetation in Wakhan. In Deutsche Wakhan Expedition, 1964. Traunstein. 1965 41-43 pp.
- Roshkova, O. De speciebus afghanicis generis Psylliostachys (Jaub. et Sp.) Nevsky. 1965 Nov. sist. vyss. rast. 210-219 p.
- Rosset, L.F. Phénomènes volcaniques anciens dans le nord de l'Afghanistan. J. 8(4): 27-41.

 1953 (Ancient volcanic phenomena in the Afghan North)
- Royal Afghan Government. A survey of progress. Ministry of Planning, Kabul. 1958
- Rozanov', A.N. The soil of Afghanistan. Soviet Soil Soi. (in Russian) 3,4: 199-208.
- Salem, A. Mollusos from prehistorio sites in Afghanistan. Trans. Amer. Philos. Soc. 1972 62(4): 57-65.
- Salem, A. Some mollusos from Afghanistan. Fieldiana, Zool. New Ser. 1: 1-89.

Salem, N.Z. & Mole, F.D. Soil geography and factors of soil formation in Afghanistan. Soil 1969 Soi. 107(4): 289-295.

Santapau, H., Futchally, Z. & Damels, J.C. Recovery ringed of birds. J. Bombay nat. Hist. 1969 Soc. 66: 147-182.

Savage, C. Wildfowl survey in southwest Asia. Progress in 1967. 19: 144-147 1968

Schaller, G.B. & Kahn, S.A. Distribution and status of markhor (Capra falconeri) Biol. 1975 Conserv. 7(3): 185-198.

Schaller, C.B. & Kirza, Z.B. On the behavior of the Kashmir markhor (Capra falconeri 1971 oashmiriensis). Hammalia. 35(4): 548-566.

Scheibe, A. Deutsche im Hindukusch. Berlin Scully, J. (Germans in the Hindu Kush)

Scheibe, A. Die Landbauverhältnisse in Nuristan. Deutsche Forschung. N.F. 1: 98-140.
1937b (The agricultural circumstances in Nuristan)

Scheibe, A. Ergebnisse der Deutschen Hindukush-Expedition 1935. Vorbemerkungen.

1939 Landwirtsch. Jahrb. 88: 12-14. (Results of the German Hindu Kush expedition 1935)

Schneider, P. Der Yak (Bos grunniens) als Haustier. III Bericht der Biologischen Wakhan 1968 Exkursion 1968. Science Kabul 5: 1-4. (The yak as domestic animal)

Schnieder, P. & Djalal, A.S. Erstnachweis einer Weichschildkröte, <u>Trionyx gangetious</u> 1970 Cuvier, 1825 in Afghanistan. Bonn. Zool. Beitr. 21(3/4): 269-273. (First record of a <u>Trionyx gangetious</u> Cuvier, 1825 in Afghanistan)

Schurmann, H.F. The mongols of Afghanistan. Mouton and Co., The Hague, The Netherlands.

Schliz, E. & Gehlhoff, W. Die Brutverbreitung des Weisstorchs im Vorderen und Hittleren 1967 Orient. Vogelwarte, 29: 48-63. (The breeding distribution of white storks in the Near and Middle East)

Schweinfurth, U. Über klimatische Trockentäler im Himalaya. Erdkunde. 10: 297-302. 1956 (On climatically dry valleys in the Himalayas)

Schweinfurt, U. Die horizontale und vertikale Verbreitung der Vegetation im Himalaya.

1957 Bonn. geogr. Abh. 20: 1-373. (The horizontal and vertical distribution of the vegetation in the Himalayas)

Scott, D.A. Iran. Chapter 3 on Flamingos. Kear, J. and N. Duplaix-Hall (Eds.). T. and 1975a A.D. Poyser, Berkhamsted, England.

Scott, D.A. Mid-winter wildfowl genuses in Iran: 1353(74/75). Department of Environment, 1975b Ornithological Unit. Nimeogr. 25 p.

- Scott, D.A. Hid-winter wildfowl counts in the wetlands of the Seisten besin. Iran Dept.

 1975c of Envir., Division of Parks & Wildlife, Ornithology Unit, special report
 W.2.54.4. Mimeo. 7 p.
- Scott, D.A. & Others. The birds of Iran. Department of the Environment. Tehran, Iran. 1975
- Scully, J. On some mammals from Kandahar. Ann. Mag. nat. Hist. ser. 5,8.
- Scully, J. On the mammals and birds collected by Capt. C.E. Yake of the Afghan Boundary 1887 Commission. J. Asiat. Soc. Beng. 56(2).
- Senarolans de Granoy, R. & Kostka, R. Grosser Pamir, Österreichisches Forschungs1978 unternehmen 1975 in den Wakhan-Pamir, Afghanistan. Akademische Druck und
 Verlagsanstalt, Graz, Austria. (Big Pamir, the Austrian research expedition
 1975 in the Wakhan-Pamir)
- Serfaty, A. & Vachon, M. Quelques remarques sur la biologie d'un scorpion d'Afghanistan. Bull. Mus. Hist. nat., Paris. 22: 215-218. (Some remarks on the biology of a scorpion of Afghanistan)
- Severtzov, N.A. On the birds of the Pamir range. Ibis. 5(1): 48-63.
- Severtzov, N.A. Faunes des Vertébrés du Turquestan. Les Oiseaux. Z. Ges. Orn. 4: 1-141.
- Severtzov, N.A. & Menzbier, N.A. Ornithology of Turkestan and the adjacent countries.

 1894 (in Russian) Moscow.
- Seward, A.C. Mesozoic plants from Afghanistan and Afghan-Turkestan. Palaeont. Ind. n.s. 1912 4(4): 1-57.
- Shahrani, M.F.K. Kirghiz pastoral nomads of the Afghan Pamirs: A study in ecological and intra-cultural adaption. PhD thesis, University Microfilms American Arbor, Michigan, Order No. 77-620.
- Sharpe, R.B. The zoology of the Afghan Delimination Commission. Trans. Linn. Soc. 1989 Lond. 5, Zool. Birds: 66-93.
- Sharpe, R.B. Soientific results of the Second Yarkand Mission, based upon the collections of the late Ferdinand Stoliczka, Aves. XVII + 153 p. London.
- Shaw, R.B. Visit to high Tartary, Yarhand and Kashgar and return journey over the 1871 Karakorum Pass. John Murray, London.
- Shor, J.B. After you, Maroo Polo. McGrew-Hill Book Co., New York. 1955
- Short, R.V. The introduction of new species of animals for the purpose of domestication.

 1976 Symp. zool. Soc. Lond. 40: 321-333.

Shumakov, E. Aoridoidea of Afghanistan and Iran. Trudy russk. Ent. Obshoh. 49: 1-248 (in 1963 Russian)

Siddiqui, M.S. Checklist of mammals of Pakistan with particular reference to the mammalian ollection in the British Kuseum (Natural History), London. 7(2): 240-250.

Singer, A. Tribal migrations on the Iran-Afghan border. Asian Affairs. 60(4): 160-165.

Singer, A. Problems of pastoralism in the Afghan Pamirs. J. Asian Affairs v. 63 (new 1976 series v. VII) pt. II: 156-160.

Sirdar Ikbal Ali Shah. Afghanistan of the Afghans. London, Deamond Press Ltd. 272 p. 1928

Smith, E.C. Some additional information on birds of Afghanistan. Ardea. 62: 226-235.

Smith, H.H., et al. Area handbook for Afghanistan. 4th edition, Washington US Government 1973 Printing Office, DAPAM 550-565.

Smith, N.A. Contribution to the herpetofauma of Afghanistan. Ann. and Mag. Nat. Hist. ser. 1940 11(5): 382-384.

Snoy, P. Die Kafiren, Formen der Wirtschaft und geistigen Kultur. Diss., Frankfurt, 1962 277 p. (The Kafirs, economy and culture).

Snoy, P. Nuristan und Humjan. Tribus. 14: 101-148.

Société d'Etudes et d'Informations économiques. La structure économique de l'Afghanistan.

1932 Paris. (The economic structure of Afghanistan)

Sokolov, I.I. (ed.) Mammalian fauna of USSR. I Insectivora, Chiroptera and Lagomorpha; 1959,63 II Cetacea, Carnivora, Pinnipedia and Artiodactyla; III Perissodactyla and Artiodactyla. Acad. Sci. USSR, Moscow (in Russian).

Stenz, E. The climate of Afghanistan: Its aridity, dryness and divisions. Geogri. Rev. 1946a 37: 672.

Stenz, E. The climate of Afghanistan, its aridity, dryness and divisions. Polish Institute 1946b of Arts and Sciences in America, N.Y. 15 pp.

Stenz, E. Le climat de Kabul. Afghanistan. 3(1): 12-19.

Stenz, E. Precipitation, evaporation and aridity in Afghanistan. Acta geophys. pol. 1957 5: 245-266.

Sterling, T. They fled to Kabul in search of law. Ceres. 4(2): 35-38.

St. John, O.S. On the birds of southern Afghanistan and Kelat. Ibis 6(1): 145-180.

Strand, R.F. The changing herding economy of Kom Muristani. Afghanistan J. 2(4) 1975

Strisemann, E. & Portenko, L.A. Atlas der Verbreitung palearktischer Vögel. Berlin. 1967 (Atlas of the distribution of palearctic birds)

Surdler, M.W. Some demographic factors regulating the formation of flocks and camps among the Brahui of Baluchistan. In: Irons, M. & Dyson-Hudson, N. (Eds.), International Studies in Sociology and Social Anthropology. Vol. 13: 69-75, Leiden Brill.

Survey of India Offices. Southern Asia Series. Afghanistan, 1: 2 000 000. Hind Reg. no. 1932 1080. Calcutta. 1st edition.

Survey of India Offices. Afghanistan. Quarter Inch. Wap 1: 253 440. Calcutta. 1941

Survey of India Offices. Southern Asia Series. Beloutchistan, 1: 2 000 000. Hind Reg. 1944 no. 1035. Revised 2nd edition 1929, reprint 1944.

Swinhoe, C. On the birds of Southern Afghanistan. Ibis. 4(6): 95-126.

Synnot, T.J. Report on forestry study for H.L.D.C. Afghanistan. Second Afghan Livestock 1978 Project, Herat. 42 p.

Taber, R.D. & Hoffmann, R.S. Behavioral adaptations of mammals to mountain environments.

1963 Proc. 6th Congr. Zool. 3:54.

Talbot, L.M. A look at threatened species: Report on some animals of the Middle East and 1960 southern Asia, which are threatened with extermination. Oryx. 5: 153-293.

Tapper, R. Nomadism in Afghanistan: asset or anachronism in: L. Dupree & L. Albert (eds.)

1974 Afghanistan in the 1970's. Praeger.

Tashhir, A. Insect fauna of Afghanistan. 3 Coleoptera; 4 Lepidoptera. Indian J. Ent. 1947-49. 3: 33-52; 20: 202-223.

Tate, G.P. Seistan. Government printing, Calcutta. Parts I to III, pp. 271, 1910.

1910 Part IV, the people of Seistan, 1912, pp. 273-374, index.

Terenter, P.V. & Chernov, S.A. Key to amphibians and reptiles. Moscow. Eng. translation, 1949 1965, by IPST. Jerusalem.

Thiollay, J.N. Précisions nouvelles sur le statut de certains rapaces en Afghanistan.

1978 Alauda. 46: 98-100. (New precisions on the status of certain raptors in Afghanistan)

Third Danish Expedition to Central Asia, Zoological Results. in: Vidensk. Neddr. dansk. naturk. Foren. Plecoptera, 112: 133-138; Thysanura, 12: 139-155; Hemiptera, 112: 225-234; Clouata, Ephemeroptera, Neuroptera, 112: 235-241; Pseudos-corpionides, 114: 245-250; Neloides, 115: 305-311; Tioks, 115: 169-174; Carabidae, 115: 175-188; Xylocopine bees, 116: 267-269; Cerambicidae, 117: 271-274; Scarabaeidae, 117: 275-287; Curculionidae, 117: 289-304; Solifugae, 118: 115-140; Lepidoptera (Rhopalocera), 118: 141-192; Araignees, 120: 81-120; Scorpionidae, 120: 121-187.

Thomas, O. A collection of mammals from Persia and Armenia. Proc. zool. Soc. Lond. 2: 520. 1905

Thomas, O. Two new Asiatic voles. Ann. Mag. nat. Hist. Ser. 8.9.

Tong, M. Report on the birds seen on the Hindu Kush. Roneotype. not published.

Transter, P. No tigers in the Hindu Kush. London. 1868

Troll, C. Die klimatische und vegetations geographische Cliederung des Himalaya-Systems.

1967 Khumbu Himal (Munchen) 1: 353-388. (The climatological and vegetation geographic levels in the Himalayas)

Tsalkin, V.L. Hountain sheep of Europe and Asia. Publ. Hosk. Obshch. Isp. Prirody. Zool. 1951 Ser. 27(42), 343 p., 101 figs., 4 pls.

U.S.S.R., Cartographic Institute. Afghanistan, 1959. 1 population; 2 economy. Ministry of Interior, Noscow (in Russian).

Udvardy, H.D. A classification of the biogeographical provinces of the world. IUCH, 1975 Gland.

Ueno, N. Insect fauma of Afghanistan and Hindu Kush. Results of the Kyoto University 1960-65 Scientific Expedition to the Karakorum and Hindu Kush. Vol. 5, Kyoto, Japan.

Ueno, S. & Nakamura, K. The anurans collected by the Kyoto University Pamir Hindu Kush

1960 Expedition. Results of the Kyoto University Scientific Expedition Karakorum and
Hindu Kush. Kyoto, Japan.

United Nations. Afghanistan, 1: 2 000 000, Map no. 279, Rev. 2. New York 1954

United Nations Economic Commission for Asia and Far East. Geological map of Asia and the 1961 Far East, 1: 5 000 000.

United States Aeronautical Chart Service. World Aeronautical Chart, 1: 1 000 000.

1946
335: Tarkand River, 337: Kugi-Tang Tan Range, 430: Hari River, 431: Khyber Pass,
441: Sulaiman Range, 442: Margo Desert. Lambert Conformal Conic Projection.

Standard Parallels 30° and 45°. Washington, D.C. Revised 1951.

- United States Air Force, Operational Navigation Chart, 1: 1 000 000. Aeronautical chart 1960-69 and information center. United States Air Force, St. Louis. Sheets: ONO-G6 (central, north), ONC-H7 (south-west), ONC-H8 (south-east).
- United States Department of Commerce. Afghanistan as a source of crude drugs and essential 1948 oils. Washington, D.C. Industr. Ref. Serv. 4(2): 133-134.
- Vasic, V.F. Observations ornithologiques en Afghanistan. Alauda. 42: 259-280. (Ornitholo-1974 gioal observations in Afghanistan)
- Vaurie, C. Notes from the Walter Koelz Collections. 1) Notes on some Ploceidae from western 1949-51

 Asia. Am. Mus. Nov. 1406, 41 p. 2) Notes on some Asiatic finches. Am. Mus. Nov. 1424, 63 p. 3) Notes on the bird genus Oenanthe in Persia, Afghanistan and India. Am. Mus. Nov. 1425, 47 p. 4) Notes on some Asiatic bitmice. Am. Mus. Nov. 1459, 66 p. 6) Notes on some Asiatic nuthatches and creepers. Am. Mus. Nov. 1472, 39 p. 8) Notes on the wrens and dippers of western Asia and India. Am. Mus. Nov. 1485, 19 p. 11) A study of Asiatic larks. Bull. Am. Mus. Nat. Hist. 97, 431-526 pp. 12) Notes on some Asiatic swallows. Am. Mus. Nov. 1529, 47 p.
- Vaurie, C. The birds of the palearctic fauna. 2 Vols., London. 1959,65
- Vavilov, N.I. & Bukinich, D.D. Report on the expedition to Afghanistan of the Institute of 1924 Applied Botany. Leningrad (in Russian).
- Vavilov, N.K. & Bukinich, D.D. Agricultural Afghanistan. Trudy prikl. Bot. Cenet. Selek. 1929 Suppl., 33 tbl., 1-610 pp.
- Viellard, J. Données biogéographiques sur l'avifaume d'Asie Occidentale. 1. Afghanistan 1969 (Première partie) Alauma. 37: 273-300. (Biogeographical data on the Avifauma of Western Asia. 1 Afghanistan)
- Vigne, G.T. A personal narrative of a visit to Ghuzni, Kabul and Afghanistan and of a 1843 residence at the court of Dost Mohomed with notices of Runjit Sing, Khiva and the Russian expedition. London, Routledge.
- Vinogradov, B.S. A third new species of dwarf jerboa. Ann. Mag. nat. Hist. Ser. 10.1. 1928
- Vinogradov, B.S. & Argiropula, A.I. Fauna of the USSR mammals. Key to rodents. Moscow.

 1941 Translated from Russian by IPST Jerusalem, 1968.
- Virgo, K.J. & Clarke, I.D. Satellites over the Hindu Kush. G. Mag. 50(6): 401-406.
- Volk, O.H. Vegetationseindrücke in Afghanistan. 3: 210-212. (Vegetation impressions in 1951 Afghanistan)
- Volk, O.H. Klima und Pflanzenverbreitung in Afghanistan. 5-6: 422-433. (Climate and 1954 plant distribution in Afghanistan)

Volk, O.H. Landwirtschaftliche Probleme Afghanistans. Mitt. Inst. Auslandsbez. 4(9-10): 1954 233-236. (Agricultural problems of Afghanistan)

Volk, 0.H. Afghanische Drogen. Planta med. 3: 171-178. (Afghan herbs)

Voous, K.N. List of recent holarotic bird species. Ibis. 115: 612-638; 119: 223-250, 1973.77 376-406.

Vos, A. de, Eilean, C. & Haghighat, C. A master plan for Mohammad Reza Shah National Park.

1977 Iran, FO:DP/IRA/76/001 Field Document. Department of Environment, Iran.

Halker, E.P. Mammals of the world. 3 Vols., Baltimore.

Haller, H. Die Vegetation Osteuropas, Nord-und Zentral Asiens. Stuttgart. (The vegeta-1974 tion of East Europe, North and Central Asia)

Ward, R. Records of big game. 7th ed. London. 1914

Wardlaw-Ramsey, R.G. Ornithological notes from Afghanistan. Ibis IV(3): 444-449; 1879.80 IV(4): 45-71 On the birds of the Hariab district. Kurram.

Weippert, D. Zur Geologie des Gebietes Doab-Saighan-Hajar (Nord Afghanistan). Beih. geol. 1964 Jb. 70: 153-184. map: 1: 120 000 (On the geology of Doab-Saighan-Hajar area)

Wendelbo, P. Plants from Tirich Mir. A contribution to the flora of Hindu Kush. Nytt. 1952 Mag. Bot. 1: 1-77.

Herdelbo, P. Studies in Primalaceae IV: the genus Dionysia in Afghanistan. Arbok Univ. 1963 Bergen. 19: 1-28.

Werdelbo, P. On the genus Eremurus in SW Asia. Arbok. Univ. Bergen. 5: 1-45.

Wendelbo, P. Trekk av Afghanistans plantegeografi. Acta Univ. Gothob., Botanica Gothob. 1966a 5: 1-20. (Nove in Afghan plant geography)

Wendelbo, P. The genus Allium. A lily group discussion. The Lily Year Book, London. 1966b 1967: 1-15.

Wendelbo, P. New subgenera, sections and species of Allium. Bot. Not. 122: 25-27. 1969a

Wendelbo, P. Two new species of Iris from Afghanistan (studies in the flora of Afghanistan 1969b X) Bot. not. 122: 204-206.

Wendelbo, P. & Furse, P. Eremurus of South-West Asia. The Lily Year Book, London. 1969: 1968 56-69.

Wendelbo, P. & Hedge, J. Studies in the flora of Afghanistan. Arbok Univ. Bergen 1963 18: 56.

Went, F.W. & Ramesh Babu, V. Plant life and desertification. Evn. Conserv. Vol. 5(4). 1978

Wermuth, M. Liste der rezenten Amphibien und Reptilien. Vo. 80 Gekkonidae, etc. Vol. 86 1965,67 Agamidae. Berlin.

Wettstein, O. von. Lacertilia aus Afghanistan. Contribution à l'étude de la faune de 1960 Afghanistan. Zool. Anz. 165: 58-63. (Lacertilia from Afghanistan)

Wettstein-Westersheimb, O. von. Drei seltene Echsen aus Stiwest-Asien. Zool. Anz. 1960 165: 190-193. (Three rare lizards from Southwest Asia).

Whistler, H. Materials for the Ornithology of Afghanistan. J. Bombay nat. Hist. Soc. 1944-45 44: 505-519; 45: 61-72, 106-122, 280-302, 462-485.

Whistler, H. Popular handbook of Indian birds. Edinburgh and London. 1949

Mhitehead, C.H.T. On the birds of Kohat and Kurram, northern India. With an introduction by Major H.A.F. Margrath. Ibis. 1909: 90-134, 214-284, 620-623. Reprinted in: J. Bombay nat. Hist. Soc. 20: 169-197, 776-799, 954-980.

Wilber, D.N. Annotated bibliography of Afghanistan. Human relations area files, New Haven. 1956 220 pp.

Wilber, D.N. Afghanistan, its people, its society, its culture. Human Relations Area 1962 Files Press, New Haven. 320 pp.

Wileshire, E.P. New species and forms of Lepidoptera from Afghanistan and Iraq. J. Bombay 1958 nat. Hist. Soc. 55(2): 288-337.

Wilson, Sir A.T. A bibliography of Persia. Clarendon, Oxford. 253 pp. 1930

Mood, J. Source of the river Oxus. Blackwood and Sons, London. 1841

Nood, J. A journey to the source of the river Oxus by Capt. John Wood. 2nd ed. John Nord. Hurray, London. 280 pp. map.

Wood, Brian, Madge, S.C. & Waller, C.S. Description moult and measurements of Montifringilla 1978 theresae. Bull. Br. Orn. Club. 18(2): 55-59.

Horld Bank. Afghanistan - The journey to economic development. Vol. 1 & 2. Report No. 1978 17778-AF, 2030/IBRD/-/4. Internal Working Document. 255 p.

Yamashita, K. Cultivated plants and their relatives. Kyoto University, Scientific 1960-65 Expedition to the Karakorum and the Hindu Kush. Vol. 1, Kyoto, Japan.

ORDER PRIMATES

Family Cercophithecidae

1. Macaca mulatta (Zimmermann)

rhesus macaque

Montane forests of monsoonal Afghanistan. Recorded from the Pech Valley, Laghman, Nuristan (KH), Paktia, Bashgal Valley (Ellermann and Morrison Scott, 1951).

ORDER LAGOMORPHA

Family Ochotonidae

1. Ochotona rutila (Severtzov)

Turkestan red pika

Recorded from northern Afghanistan, east to the Pamirs, over 2 000 m (Corbett, 1978), also from Khwahan (Darwaz) (CN & JN).

2. Ochotona roylei (Ogilby)

large-eared pika

(including <u>0. macrotis</u> (Günther)). Montane steppes of the Pamirs and Nuristan between <u>2 000</u> and 4 800 m elevation. Recorded from Wakhan, and Big Pamir, between 4 000 and 4 800 m (CN), Big and Small Pamir (RP), southern Badakhshan (JH), western Nuristan (RP), Stiewe and Weran Passes 3 350 m (KP). Sight records from the Salang Pass (Madge, 1970) probably refer to 0. rufescens).

3. Ochotona rufescens (Gray)

Afghan pika

O. r. rufescens. Montane steppes of the Hindu Kush Mountains, west to Herat, between 1 000 and 3 500 m altitude. Reported from Shibar Pass and around the Sawzak Pass (JH), Gardez, Dashte Nawar, Unai Valley, Paghman, Bande Amir, Salang and Herat (JN), Paghman (KZ).

Family Leporidae

1. Lepus capensis Linnaeus

Cape hare

L. c. tibetanus Waterhouse. Common throughout Afghanistan except in the monsoonal east where L. c. ruficaudatus Geoffroy is found. Recorded from the Pamir and Wakhan (CN), Sarobi, Kabul, Istalif, Dashte Nawar (JN), Herat, Seistan and Sabzak Pass (KH).

ORDER RODENTIA

Family Sciuridae

- 1. Petaurista petaurista (Pallas) common giant flying squirrel

 P. p. albiventer (Gray). Evergreen oak and conifer forest. One record from Pech river, 1 000 m Nuristan (JN).
- 2. Hylopetes fimbriatus (Gray) smaller Kashmir flying squirrel

 H. f. baberi (Blyth). Locally abundant in evergreen monsoonal oak and conifer forest between 1 600-3 500 m elevation. Records from Pech river 1 000 m (JN), Nyrow, Kohistan (Khost?) (Blyth, 1847), Peiwar Kotal (Blanford, 1881).

Yate, C.E. Northern Afghanistan. London. 419 p. 1888

Yate, C.E. Khursan and Seistan. London 1900

Yule, Col. Sir H. ed. The book of Sir Marco Polo. 3rd edition. Murray, London, 1: 462; 1929 2: 662.

Zarudny, N.A. Materials for the study of the birds of the Pamir and surrounding countries.

1926 Byull. mosk. Obshch. Ispyt Prir. 35: 233-287 (in Russian).

Zimmermann, K. Zur Fauna Afghanistan. Z. Säugetierk. 20: 189-191. (On the fauna of 1955 Afghanistan)

Zimmermenn, K. Fledermäuse aus Afghanistan. Z. Säugetierk. 21: 195-196. (Bats from 1956 Afghanistan)

Zohary, M. On the geobotanical structure of Iran. Bull. Res. Conn. Israel. 110, suppl. 1963

Appendix 10

BIBLIOGRAPHY - SUBJECTS

I. GEOGRAPHY

General

Natural Sites

- 1. Ab-i-Estada
- 2. Dashte Nawar
- 3. Seistan, Hamun-i-Puzak
- 4. Registan and Kandahar
- 5. Hindu Kush
- 6. Pamir and Wakhan
- 7. Nuristan, Laghman, Safed Koh, etc.
- 8. Kole Hashmat Khan, Kabul, Qargah, etc.
- 9. Lower Kabul Valley, Jalalabad Valley
- 10. Ajar Valley
- 11. Bande Amir, Bamian
- 12. Amu Darya Valley, Darqad and West Badakhshan
- 13. Turkestan, Herat

II. DEMOGRAPHY

III. PHYSICAL ENVIRONMENT

Climate

Geology and Soils

Archaeology

IV. GENETIC RESOURCES

V. ENVIRONMENT

General

Irigation

Forestry

Range Management

Boonomy

National Parks, including Legislation and Tourism

Hunting and Fur Trade

VI. BIBLIOGRAPHY

VII. MAPS

VIII. VEGETATION

Afghanistan

General

IX. FAUNA

Mammals

- 1. Afghanistan
- 2. Regional (general)
- 3. Regional
- 4. Ungulates
- 5. Carnivores
- 6. Small Nammals

Birds

- 1. Afghanistan
- 2. Regional
- 3. Haterfowl
- 4. Migration and Bird-ringing

Reptiles and Amphibians

Lower Animals

I. GEOGRAPHY

General

Afghanistan Journal; Akram, N., 1951; Balland, V. & D. Balland, 1972; Barrat, G., 1972; Conolly, E., 1840; Dollot, R., 1937; Dupree, L. 1973; Dupree, L. & L. Albert eds., 1974; Edelberg, L. & Sohuyler, 1978; Field, N.C., 1954; Furon, R., 1926; Humlum, J., 1959; Klimberg, M., 1966; Kraus, W., 1972; Misra, S.D., 1967; Perrin, N., 1842; Reiner, E., 1962; Sirdar Ikbal Ali Shah, 1928; Smith, H.H. et al, 1973; Tate, G.P., 1910; Wilber, D.N., 1962.

Natural Sites

1. Ab-1-Estada

Akhtar, S.A., 1947; FAO, 1965; FAO, 1975a; FAO, 1975b; FAO, 1977a; FAO, 1977e; FAO, 1978c; Forstner, U. & G. Bartsch, 1970; Ganss, O., 1970; Koning, P.J. & J. Rooth, 1975; Musson, C., 1974; Niethammer, G., 1971; Niethammer, J., 1970a; Nogge, G., 1969; Nogge, G., 1972; Nogge, C., 1974; Paludan, K., 1959; Pelt, J.M. et al, 1968; Pelt, J.M. et al, 1969; Savage, C., 1968; Vieillard, J., 1969.

2. Dashte Nawar

Dupree, L., 1975b; FAO, 1975a; FAO, 1975b; FAO, 1977a; FAO, 1977e; FAO, 1978c; Forstner, U. & G. Bartsch, 1970; Jux, U., 1969; Klockenhoff, H. & G. Hadel, 1970; Koning, F.J. & J. Rooth, 1975; Lapparant, A.F., Z. de Lavegne Sainte Lausanne & P. Bordet, 1965; Niethammer, G., 1967; Nogge, C., 1972; Nogge, C., 1974.

3. Seistan, Hamun-i-Puzak

Annandale, N., & H.G. Carter, 1919; Atchison, J.E. Consultants, Inc., 1974; Baker, E.C.S., 1919; Concily, E., 1840; Cumming, J.W.N., 1905a; Fairservis, W.A., Jr., 1961; FAO, 1976a; FAO, 1976d; FAO, 1976e; FAO, 1978c; Fforstner, U. & G. Bartsch, 1970; Hassinger, J.D., 1970; Huntington, E., 1905; Lalande, P., Essone Selemani & K. Parenda, 1975; Michels, A.A. 1959; Paludan, K., 1959; Scott, D.A., 1975; Tate, G.P., 1910; Yates, C.E., 1900.

4. Registan and Kandahar

Gaisler, J., 1975; Hassinger, J.D., 1970; Hutton, I., 1845; Hutton, T., 1847; Lalande, P., Essone Selemani & K. Parenda, 1975; Murray, J.A., ND; Paludan, K., 1959; St. John, O.B., 1889; Swinhoe, C. 1882.

5. Hindu Kush

Akram, N., 1967; Biddulph, J., 1880; Breckle, S.W., 1973; Breckle, S.H., & W. Frey, 1974; Deutsche im Hindukusch, 1937; FAO, 1975o; FAO, 1977b; PAO, 1978j; Ferdinand, K., 1964; Flohn, N. 1969; Freisleben, R., 1941; Gansser, A., 1964; Gilbert, O.L., 1968s; Gilbert, O.L., 1968b; Hacokel, I., ed., 1935;

Hacokel, I. & W. Trool, 1938; Hassinger, J.D., 1970; Hay, R.G., 1840;
Hayden, H.H., 1911; Kitamura, S., 1960; Kitamura, S., 1964; Kyoto University,
1960-65; Lange, de la Camp, M., 1939; Memertzhagen, R., 1939; Paludan, K.,
1959; Petooz, R.G., 1973b; Pimme, S.L., 1968; Riemer, E., 1938; Soheibe,
A., 1937; Scheibe, A., 1939; Tong, M. 1968; Tsalkin, V.I., 1951; Ueno, M.,
1960-65; Virgo, K.J. & I.D. Clarke, 1978; Wendelbo, P., 1952.

6. Pamir and Wakhan

Adams, L., 1858; Bonvalet, C., 1889; Bowdler Sharpe, R., 1891; Brooke, V. & Brooke, B., 1875; Cotton, W.B., 1914; Cumberland, G., 1895; Curron, G.N., 1896; Dor, R., 1976a; Dor, R., 1976b; Dor, R. & C. Naumann, 1978; Dunmoor, The Earl of, 1894; Egorov, O.V. 1965; FAO, 1975d; FAO, 1977a; FAO, 1978o; FAO, 1978d; FAO, 1978e; FAO, 1978f; Cansser, A., 1964; Cillan, C.V.B., 1934; Gratzi, K., ed., 1970; Grote, H., 1951; Groun-Grshimailo, Gr., 1890; Ivanov, 1969; Komroff, E., 1926; Kullmann, E., 1968; Kussmaul, F., 1965; Lindberg, K., 1961; Madel, G., 1968; McDonald, W., 1968; Meklenburtzer, R.N., 1948; Mirwald, P. & Roemer, H., 1967; Muss, M., 1978; Nalivkin, I.D., 1932; Naumann, C., 1973; Naumann, C.M., 1978; Naumann, C. & G. Niethammer, 1973; Naumann, C. & J. Niethammer, 1975; Niethammer, G., 1973; Nogge, G., 1973; Olufsen, O., 1904; Petocz, R.G., 1971; Petocz, R.G., 1973o; Petocz, R.G., 1973d; Poelt, J. & V. Wirth, 1968; Ponoins, E.D., 1895; Potapov, R.L., 1963; Reinig, W.F., 1932; Roemer, H., 1965; Schneider, P. 1968; Schweinfurth, U., 1956; Schweinfurth, U., 1957; Senarclans de Grancy, R. & R. Kostka, 1978; Severtsov, N.A., 1883; Shahrani, M.N.M., 1976; Shaw, R.B., 1871; Shor, J.B., 1955; Singer, A., 1976; Tsalkin, V.L., 1951; Wendelbo, P., 1952; Wood, J., 1841; Wood, J., 1872; Yule, Col. Sir H., ed, 1929; Ueno, S. & K. Nakamura, 1960; Zarudny, N.A., 1926.

7. Nuristan, Laghman, Safed Koh, etc.

Ahlemann, H., 1970; Artohison, J.E.T., 1879; Breckle, S.W. & W. Frey, 1974; Chuma, T., ed., 1970; Christison, A.F.P., 1941; Duokelmann, R. von, 1951; Edelberg, L. & Schuyler, 1978; Electrowatt & Norconsult; FAO, 1976c; FAO, 1977a; FAO, 1977g; FAO, 1978c; FAO, 1978h; Habibi, K., Sr., 1970; Hayon, J.C., J.M. Pelt, & M.C. Younos, 1968; Hayon, J.C., J.M. Pelt, & M.C. Younos, 1970; Holdich, T.H. 1896; Holloway, C.L., 1975; Jones, S., 1966; Kerstan, G., 1937; Lalande, P., 1967; Löhrl, H. & G. Thieloke, 1969; Nauroz, M.K. & C.M. Naumann, 1975; Neubauer, H.F., 1954o; Paludan, K., 1959; Puget, A., 1970b; Petcoz, R.G., 1972; Robertson, G.S., 1896; Robinson, J.A., 1935; Robertson, G.S., 1897; Scheibe, A., 1939; Snoy, P., 1965; Vieillard, J., 1969; Wardlaw-Ramsay, R.G., 1879-80; Whitehead, C.H.T., 1909.

8. Kole Hashmat Khan, Kabul, Qargah, etc.

Burnes, A., 1843; FAO, 1965; FAO, 1978o; FAO, 1978i; Hayon, J.C., J.M. Pelt & Ch. Younos, 1970; Iven, H.E., 1924; Kilbertus, G. & J.F. Pierre, 1970; Kumler, M.L., 1969; Lalande, P., 1968; Niethammer, J. 1967b; Puget, A., 1971; Stenz, E., 1948.

9. Lower Kabul Valley, Jalalabed Valley

Gaisler, J. et al, 1967; Viellard, J., 1969.

10. Ajar Valley

FAO, 19750; FAO, 1975e; FAO, 1976b; FAO, 1977a, FAO, 1977f; FAO, 1978o; FAO, 1978j; Faymann, T., 1976; Holloway, C.L., 1975; Pelt, J.M., J.C. Hayon & M. Ch. Younos, 1970; Scheide, A., 1937; Heippert, D., 1964.

47

11. Bande Amir, Bemian

Balland, V. & J. Leng, 1974; Bristol University Expedition, 1969; FAO, 1974c; FAO, 1975b; FAO, 1975c; FAO, 1977a; FAO, 1978c; FAO, 1978g; Faymann, T., 1976; Fürstner, U. & G. Bartsch, 1970; Hay, W.R., 1935; Hayon, J.C., G. Kilbertus & J.M. Pelt, 1970; Jux, U. & Kempf, 1971; Pelt, J.M., J.C. Hayon & H. Ch. Younos, 1970; Scheide, A., 1937; Wood, Brian, Madge S.S. & Haller, C.S., 1978; Dieterle, A., 1973.

12. Amu Darya Valley, Darged and West Badakhshan

Brückle, K., 1935; Desro, A. et al, 1964; Field, N.C., 1954; Holloway, C.L., 1975; Naumann, C. & J. Niethammer, 1975; Nauroz, K.K. & C.M. Raumann, 1975; Paludan, K., 1959; Pelt, J.M., J.C. Hayon & M. Ch. Younos, 1968; Pelt, J.M., J.C. Hayon, & M.Ch. Younos, 1969; Petocz, R.G., 1973a.

13. Turkestan, Herat

Aitchison, J.E.T., 1888; Bogdanov, O.P., 1962; De La Tour, G.D., 1975; FAO, 1958; FAO, 1977b; Ferrier, J.P. 1976; Groves, C.P. & V. Kazak, 1967; Hassinger, J.D., 1970; Heptner, M.G., 1937; NcArthur, I.D. & Harrington, G.N., 1978; McArthur, I.D., S. Sayed & M. Nawin, 1979; Meinertzhagen, R., 1939; Paludan, K., 1959; Rechinger, K.H. & H. Riedl, 1961, 62; Severtzov, N.A., 1887; Severtzov, N.A. & N.A. Menzbier, 1894; Seward, A.C., 1912; Synnott, T.J., 1978; Yate, C.E., 1888.

II. DEHOGRAPHY

Afghanistan Cartographic Institute, 1959; Barth, F., 1967; Bellew, H.W., 1880; Biddulph, J., 1880; Dor, R. & C. Naumann, 1978; Dupree, L., 1973; Dupree, L., 1975a; Dupree, L. & L. Albert, eds., 1974; Edelberg, L. & Sohuyler, 1978; Ferdinand, K., 1962; Ferdinand, K., 1964; Ferdinand, K., 1969; Ferdinand, K., 1976; Holdich, T.H., 1896; Jentsch, C., 1973; Jones, S., 1978; Kussmaul, F., 1965; Robertson, G.S., 1896; Robertson, G.S., 1897; Robinson, J.A., 1935; Schurmann, H.F., 1962; Senarolans de Grancy, R. & R. Kostka, 1978; Shahrani, M.N.M., 1976; Singer, A., 1973; Sirdar Ikbal Ali Shah, 1928; Smith, H.H. & others, 1973; Snoy, P., 1962; Sterling, T., 1971; Surdler, W.W., 1969; Tapper, R., 1974; Wilbur, D.N., 1962.

III. PHYSICAL ENVIRONMENT

Climate

Afghanistan, Civil Aviation and Tourism Authority, General Meteorological Directorate, Arnez, G.J., 1969; Borisov, A.A., 1965; Breokle, S.W., 1970; Breokle, S.W., 1973; Edelberg, L. & Schuyler, 1978; Flohn, H., 1969; Herman, N.M., 1965; Herman, N.M., J. Zillhardt & P. Lalande, 1971; Irwin, Lieut., 1839; Iven, H.E., 1924; Lalande, P., N.M. Herman & J.Z. Zillhard, 1973-74; Lincheviski, I.A., 1946; Rathjens, C., 1974; Schweinfurth, U., 1956; Stenz, E., 1946; Stenz, E., 1948; Stenz, E., 1957; Troll, C., 1967; Volk, C.H., 1954.

1,2

Geology and Soils

Balland, V. & J. Leng, 1974; Béné, R. & S. Baldo, 1965; Blandford, W.T., 1876; Brückle, K., 1933; Brückle, K., 1935; Bundesanetalt für Bodenforschung, 1969; Clapp, F.G., 1939; Desro, A., et al, 1964; FAO, 1962b; FAO, 1965; FAO/UNDP, 1969, 70, 71; FAO/Unesco, 1977; Fox, C.S., 1936; Freise, F., 1907; Furon, R., 1937; Furon, R., 1941; Furon, R. & Louis-Felicien, R., 1954; Ganss, O., 1970; Gansser, A., 1964; Hauser, G.F.; Hayden, H.H., 1911; Herbordt, O., 1926; Humlum, J., 1959; Huntington, E., 1905; Irwin, Lieut., 1839; Jux, U., 1969; Kästner, H., 1971; Lapparent, A.F., Z. de Lavegne Sainte Lausanne & P. Bordet, 1965; Kisra, D.S., 1967; Nalivkin, I.D., 1932; Pias, J., 1976; Pilger, A., 1971; Pilger, A. & A. Rösler, 1976; Rathjens, C., 1972; Rosset, L.F., 1953; Rozanov', A.N., 1945; Salem, N.Z. & F.D. Mole, 1969; Seward, A.C., 1912; United Nations Economic Commission for Asia and Far East, 1961; Weippert, D.

Archaeology

Dor, R., 1976; Dupree, L., 1958; Dupree, L., 1972; Dupree, L., 1975b; Fairservis, Ealter A., Jr., 1961; FAO, 1978d.

IV. CENETIC RESOURCES

FAO & IRPGR, 1979; FAO/UNEP, 1980; Freisleben, R., 1941; Gorgas, N., 1966; Harrison, S.G., 1952; Imankulov, M.I., 1969; Lange, de la Camp, N., 1939; Mehlenburtzev, R.W., 1940; Nadler, C.F. et al, 1971; Nadler, C.F. et al, 1973; Neubauer, H.F., 19540; Pelt, J.W., J.C. Hayon & N.C. Youngs, 1965; Short, R.V., 1976; Vairlov, N.I. & D.D. Bukinich, 1929; Yamashita, K., 1960-65; Yalcin, B.C., 1979.

V. ENVIRONMENT

Ceneral

Atchison, J.E. Consultants, Inc., 1974; Calembert, J., & L.R.N. Strange, 1978; FAO; 1977a; FAO, 1978a; FAO, 1978o; FAO, 1980a; Firouz, E., 1974; Ferdinand, K., 1969; Caisler, J., 1975; Hassanyar, Amir S., 1972; Hassanyar, A.S., 1973; IUCN, 1975; Kilbertus, C. & J.F. Purre, 1970; McArthur, I.D. & Harrington, C.N., 1978; Rathjens, C., 1968-69; Went, F.W. & V. Ramesh Babu, 1978.

Irrigation

FAO, 1980b; FAO/UNDP, 1969, 70, 71; Horning, H.M., 1969; Humlum, J., 1959.

Forestry

Beranger-Beanguesne, L., 1956; FAO, 1973a; FAO, 1973o; FAO, 1974a; FAO, 1976o; FAO, 1979a; FAO, 1979o; FAO, 1979d; FAO, 1980a; FAO, 1980c; Rathjens, C., 1968-69; Synrot, T.J., 1978.

Range Management

Barth, F., 1967; Bulbul, S.P.D. & V.I. Hasalov, 1964; Calembert, J., & L.R.N. Strange, 1978; FAO, 1973a; FAO, 1974a; FAO, 1974d; FAO, 1977; FAO, 1980b; FAO, 1980c; Ferdinand, K., 1962; Ferdinand, K., 1976; Imankulov, N.I., 1969; Jentsch, C., 1973; Lapparent, A.F., 1970; Keshtyar, M., 1970; McArthur, I.D. & G.N. Harrington, 1978; McArthur, I.D., Sayed, S. & H. Navem, 1979; McClure, B., 1979; Nuri, J.M., 1961; Pabot, H., 1959; Pabot, N., 1967; Rako, A. & J. Cizek, 1977; Shahrani, N.M.N., 1976; Strand, R.F., 1975; Synnot, T.J., 1978; Tapper, R., 1974; Woodford, M.H., 1978; Yalcin, B.C., 1979.

Economy

Afghanistan, 1959; Afghanistan, Central Statistic Office, 1978; Afghanistan, Ministry of Planning, 1979; Aitchison, J.E.T., 1890; Barth, F., 1975; Bulbul, S.P.D. & V.I. Masalov, 1964; Economist Intelligence Unit, 1979, 80; Electrowatt & Norconsult; Eskilsson, E., 1966; FAO, 1969a; Ferdinand, K., 1962; Fox, C.S., 1936; Inskipp, T. & S. Wells, 1979; Irwin, Lieut., 1839; Janata, A., 1963; Krans, W., 1972; EcArthur, I.D., S. Sayad & M. Nawin, 1979; Michels, A.A., 1959; Pelt, J.M., J.C. Hayon & M.C. Younos, 1965; Royal Afghan Government, 1958; Scheibe, A., 1937; Smith, H.H. & others, 1973; Société d'Etules et d'Informations économiques, 1932; Sterling, T., 1971; Strand, R.F., 1975; United States Department of Commerce, 1946; Vavilov, N.I., & D.D. Bukinich, 1929; Volk, O.H., 1954; Volk, O.H., 1955; Woodford, M.H., 1978; World Bank, 1978.

National Parks, including Legislation and Tourism

Afghanisten, Central Statistic Office, 1975; FAO, 1970, 71; FAO, 1973d; FAO, 1974b; FAO, 1974c; FAO, 1977a; FAO, 1977e; FAO, 1977f; FAO, 1978c; FAO, 1978f; FAO, 1978s; FAO, 1978h; FAO, 1980a; Faymann, T., 1976; IUCN, 1975; Vos, A. de, C. Eilean & G. Haghighat, 1977.

Hunting and Fur Trade

Ahlemann, H., 1970; Elittiker, W., 1959; Inskipp, T. & S. Wells, 1979; FAO, 1973d; FAO, 1973e; FAO, 1974b; FAO, 1976b; FAO, 1977a; FAO, 1977d; FAO, 1977f; FAO, 1978e; FAO, 1978e,f; FAO, 1980a; Niethammer, G., 1972; Niethammer, J., 1967e; Nogge, G., 1973; Nogge, G. & J. Niethammer; Petooz, R.G., 1973d; Ponoins, E.D., 1895.

VI BIBLIOGRAPHY

Akram, M., 1947; Arbeitsgemeinschaft Afghanistan, 1968; Breckle, S.W., W. Frey & I.C. Hadge; 1969; British Embassy, 1974; Burgess, R.L., A. Moktarzadeh & L. Cornwallis, 1966; Dada, H. & Pickett, L., 1969; Jones, S., 1966; Kästner, H., 1971; Meher-Homji, V.N., R.K. Gupta & H. Freitag, 1973; Pickett, L.C., Ma-Yar, N.Q. & Z. Saleh, 1968; Wilber, D.N., 1956; Wilson, Sir A.T., 1930.

VII KAPS

Afghanistan; Afghanistan Cartographic Institute, 1968; Bundesanstalt fur Bodenforschung, 1969; FAO, 1965; FAO & Unesoo, 1977; Geographical Section, 1942; Gratzi, K., ed, 1970; Survey of India Offices, 1932; Survey of India Offices, 1941; Survey of India Offices, 1944; United Nations, 1954; USSR, Cartographic Institute, 1959; United States Aeronautical Chart and Information Service, 1951; United States Air Force, Operational Navigation Chart, 1960-69.

VIII VECETATION

Afghanistan

Adye, Col. J., 1867; Apachanjanc, O.E., 1958; Agachanjanc, O.E., 1960; Agachanjanc, O.E., 1961; Agachanjanc, O.E., 1964; Aitchison, J.E.T., 1379; Aitchison, J.E.T., 1881-82; Aitchison, J.E.T., 1880; Aitchison, J.E.T., 1888; Aitchison, J.E.T., 1889b; Aitchison, J.E.T., 1890; Akram, M. 1967; Annandale. N. & H.G. Carter, 1919; Béranger-Beauguesne, L., 1956; Bharucha, F.R., 1955; Bornmuller, J., 1934; Breckle, S.W., 1971; Breckle, S.W., 1973; Breckle, S.W. & W. Frey, 1974; Breckle, S.W., W. Frey & I.C. Hedge, 1969; Bretzl, H., 1903; Burnes, Sir A., 1843; Chuma, T., ed, 1970; Dieterle, A., 1973; Diver, H., 1935; Duokelmann, R. von, 1951; Edelberg, L. & Schuyler, 1978; FAO, 1958; FAO, 1959; FAO, 1976c; FAO, 1977e; FAO, 1977f; FAO, 1978d; FAO, 1978h; FAO, 1978i; FAO, 1978j; Ferrier, J.P., 1976; Freisleben, R., 1941; Freitag, K., 1971a; Freitag, H., 1971b; Freitag, H., 1972; Frey, W., 1967; Frey, W., 1968; Gilbert, O.L., 1968a; Gilbert, O.H., 1968b; Gilli, A., 1954-64; Gilli, A., 1961-62; Gilli, A., 1962-64; Gilli, A., 1969; Gratzi, K., ed., 1970; Haeckel, T. & W. Troll, 1938; Hassanyar, A.S., 1973a; Hassanyar, A.S., 1973c; Hassanyar, A.S., 1977; Hassinger, J.D., 1968; Hayon, J.C., G. Kilbertus & J.M. Pelt, 1970; Hayon, J.C., Pelt, J.M., & N.C. Younos, 1968; Hayon, J.C., J.M. Pelt & N.C. Younos, 1970; Hedge, I.C., 1960-68; Hedge, I.C., 1963; Hedge, I. & P. Wendelbo, 1963; Hedge, I. & P. Wendelbo, 1970; Humlum, J., 1959; Jaffri, S.N.H., 1954; Kersten, G., 1937; Keshtyar, M., 1970; Khan, M.I.R. & A.R. Beg, 1968; Kitamura, S., 1956-58; Kitamura, S., 1963; Kitamura, S., 1960; Kitamura, S., 1964; Koeie, N. & K.H. Rechinger, 1954-65; Kumler, M.L., 1969; Kyoto University, 1960-65; Lalande, P., 1967; Lalande, P., 1968a; Lalande, P., 1968b; Lalande, P., 1968o; Lapparent, A.F., 1970; Lange, de la Camp, N., 1939; Linchevski, I.A., 1946; Linchevski, I.A. & A.V. Prozorovski, 1944; Linchevsky, I.A. & A.V. Prozorovsky, 1949; McClure, B., 1979; Meher-Homji, V.M., R.K. Gupta & H. Freitag, 1973; Muss, M. 1978; Nedyalhov, S.T., 1978a; Nedyalkov, S.T., 1978b; Neubauer, H.F., 1951; Neubauer, H.F., 1954a; Neubauer, H.F., 1954b; Newbauer, H.F., 1954c; Nuri, J.M., 1961; Paludan, K., 1959; Pelt, J.M., Hayon, J.C. & M.C. Younos, 1968; Pelt, J.N., Hayon, J.C. & M.C. Younos, 1969;

Pelt, J.K., Hayon, J.C. & M.C. Younos, 1970a; Pelt, J.M., Hayon, J.C. & M.C. Younos, 1970b; Podleoh, D., 1967-69; Petrak, F., 1963; Rathjens, C., 1974; Poelt, J. & V. Wirth, 1968; Rathjens, C., 1968-69; Rathjens, C., 1972; Rechinger, K.H., 1949; Rechinger, K.H., 1963; Rechinger, K.H. & M. Koie, 1954-63; Rechinger, K.H. & H. Riedl, 1961-62; Riemer, E., 1938; Roemer, H., 1965; Roshkova, O., 1965; Senarolans de Granoy, R. & R. Kostka, 1978; Tate, G.P., 1910; Vavilov, N.I. & Senarolans de Granoy, R. & R. Kostka, 1978; Tate, G.P., 1910; Vavilov, N.I. & D.D. Bukinich, 1924; Volk, O.H., 1951; Volk, O.H., 1954; Wendelbo, P., 1966a; Wendelbo, P., 1966b; Wendelbo, P., 1969; Wendelbo, P., 1964; Wendelbo, P., 1966a; Wendelbo, P., 1968; Wendelbo, P., 1969; Wendelbo, P., & J. Hedge, 1963; Tate, C.E., 1888.

::

General

Bobek, H., 1951; Champion, Sir H.G., S.K. Seth & G.M. Khattak, 1966; Davis, P.H., P.C. Harper & I.C. Hedge, 1971; Floyer, E.A., 1977; Worden, W., 1927; Oedekoven, K., 1964; Pabot, H., 1961; Raven, P.M., 1971; Rechinger, K.H., 1961-68; Rechinger, K.H., 1965; Riedl, H., 1963; Schweinfurth, U., 1957; Troll, C., 1967; Rechinger, K.H., 1974; Mendelbo, P., 1969; Wendelbo, P. & P. Furse, 1968; Went, F.W. & V. Ramesh Babu, 1978; Zohary, H., 1963.

IX FAUNA

Hammals

1. Afghanistan

Aitchison, J.E.T., 1889a; Adye, Col. J: Sitana, 1867; Blandford, W.T., 1876; Burnes, Sir A., 1843; Dunmoor, The Earl of, 1894; Edelberg, L. & Schuyler, 1978; FAO, 1973d; Ferrier, J.P., 1976; Gaisler, G., 1975; Gaisler, J., D. Povolny, Z. Sebek & F. Tenora, 1968; Humlum, J., 1959; Kullmann, E., 1970; Kyoto University, 1960-65; Naumann, C.M., 1978; Povolny, D., 1967; Reinig, W.F., 1932; Senarolans de Grancy, R. & R. Kostka, 1978; Sharpe, R.B., 1889; Tate, G.P., 1910; Third Danish Expedition to Central Asia; Yate, G.E., 1888.

2. Regional (general)

Adlen, V., 1959; Ahlemann, H., 1970; Barnus, V. & F. Tenora, 1967; Blan Blanford, W.T., 1881; Brooke, V., Brooke, B., 1875; Cotton, W.B., 1914; Cumming, J.W.N., 1905b; De La Tour, G.D., 1975; Dupree, L., 1956; Dupree, L., 1958; Dupree, L., 1973; FAO, 1970,71; FAO, 1973f; FAO, 1975d; FAO, 1975e; FAO, 1976b; FAO, 1976f; FAO, 1977a; FAO, 1978e; Ferrier, J.P., 1976; Gaisler, J., 1970a; Gaisler, J., 1970b; Gaisler, J., 1971; Gaisler, J., 1975; Gaisler, J., D. Povolny, Z. Sebek & F. Tenora, 1967; Gillan, G.V.B., 1934; Grote, H., 1951; Groves, C.P. & V. Mazak, 1967; Hanak, V. & J. Gaisler, 1969; Hassinger, J.D., 1968; Hassinger, J.D., 1970a; Hassinger, J.D., 1970b; Hay, R.G., 1840; Hutton, I., 1842a; Hutton, I., 1842b; Hutton, I., 1842o; Hutton, I., 1845; Johnstone, Cavin, 1968; Kullmann, E., 1965; Kullmann, E., 1967; Kullmann, E., 1968; MoDonald, M., 1968; Heyer-Oehme, D., 1965; Neyer-Oehme, D., 1968; Hurray, J.A.; Murray, J.A., 1887; Meklenburtzer, R.N., 1948; Naumann, C., 1973; Naumann, C. &

- 3. <u>Spermophilopsis _ ytodactylus</u> (Lichtenstein) long-clawed ground squirrel
 - S. 1. bactrianus Scully. Northern and northwestern Afghanistan in loess, steppe habitats below 1 000 m elevation. Records from Shibergan, Karam Kol (30 km W of Kunduz) (JN), Kunduz, Mazar-i-Sharif, Maimanah, Herat (JH), Khamiab (Scully, 1888).
- 4. Marmota caudata (Jacquemont)

long-tailed marmot

- M. c. aurea (Blanford). Wakhan, Pamir.
- M. c. dichrous (Anderson). Hindu Kush, north of Kabul. Common throughout sub-alpine and alpine biotopes of Afghanistan, between 2 500 and 4 000 m elevation. Plentiful at Dashte Nawar (JN), and Pamir and valley heads in Nuristan (RP).
- 5. Spermophilus fulvus Lichenstein Large-toothed souslik
 (Synonym Citellus fulvus). Northwestern and southeastern Afghanistan, sandy steppes and alpine meadows between 1 000-3 500 m elevation.
 Records from 30 km W of Herat, Dashte Nawar, Ab-i-Estada, Mukur-Ghazni (JN).

Family Cricetidae

1. Calomyscus bailwardi Thomas

mouse-like hamster

- C. b. musteri Ellerman. Widely distributed between 400 and 3 500 m elevation in the arid mountains and surrounding steppes but not in Wakhan. As abundant in evergreen oak forest as in the barren hills north of Kandahar and the Jalalabad Valley.
- 2. Cricetulus migratorius (Pallas) migratory hamster

 Throughout the country between 400 and 3 600 m elevation; common in dry rocky and open areas.
- 3. Alticola roylei (Gray).

Royle's mountain vole

- A. r. roylei. Hindu Kush and Pamir mountains between 2 000 and 4 300 m elevation, typical for areas with rocky slopes. Recorded from Wakhan (JH), Big Pamir (CN), Paghman, N. Salang, W. Pansjir (JN), Shibar Pass (C) and Ishkashem (N) (JH), Do Schak (W Salang, C) (KZ).
- 4. Pitymys afghanus (Thomas)

Afghan vole

- P. a. afghanus. Common in eastern, central and northwestern
 Afghanistan on high montane meadows along water courses. Recorded
 from Gulran (NW) (Thomas, 1912), Herat, Kabul, Shibar Pass, Dashte Nawar,
 Unai Pass (C), Salang (JN), Shibar Pass (C) and Sabzak Pass (W) (JH).
- 5. Pitymys juldaschi (Severtzov) Pamir vole
 Wakhan, Pamir and Kohe Baba (C) montane meadows above 2 000 m elevation.
 Recorded from Panjao (C) (KP), Qala Panja (Wakhan) and Tohbai (Big
 Pamir), Aksu (Small Pamir) (RP) (CN), Faizabad (NE) (JH).
- 6. Microtus socialis (Pallas) social vole

 Montane valleys. Kabul (JN), Afghan Turkestan (Scully, 1887), (JH),
 Ghilzai (C) (KP).
- 7. <u>Microtus transcaspicus</u> Satunin Transcaspian vole

 Montane meadows of the Hindu Kush above 2 000 m. Paghman, Unai Pass
 (C) (JN), Shibar Pass (C) (JH), (Chaworth-Musters).

G. Niethammer, 1973; Naumann, C. & J. Niethammer, 1975; Naumann, C. & G. Nogge, 1973; Nauroz, M.K. & C.M. Naumann, 1975; Neuhauser, H.N. & A. De Blase, 1974; Niethammer, G. & J., 1967b; Niethammer, J., 1965; Niethammer, J., 1966b; Niethammer, J., 1967o; Niethammer, J., 1967d; Niethammer, J., 1967e; Niethammer, J., 1969a; Niethammer, J., 1969b; Niethammer, J., 1970b; Niethammer, J., 1973a; Niethammer, J., 1973b; Niethammer, J. & Martens, 1975; Nogge, G., 1972; Nogge, G., 1974; Paludan, K., 1948-49; Petocz, R.G., 1971; Petocz, R.G., 1972; Petocz, R.G., 1973a; Petocz, R.G., 1973b; Petocz, R.G., 1973o; Petocz, R.G., 1973d; Ponoins, E.D., 1895; Povolny, D., 1966; Richter, H., 1970; Schneider, P., 1968; Soully, J., 1881; Soully, J., 1887; Senarclans de Grancy, R. & R. Kostka, 1978; Sharpe, R.B., 1889; Transter, P., 1868; Zimmermann, K., 1955; Zimmermann, K., 1956.

3. Regional

Adams, L., 1858; Aleem, A., 1976; Ali, S., 1949; Blyth, E., 1840; Blyth, E., 1841; Bobrinskiji, N.A., B.A. Kusnezov & A.P. Kusjakin, 1965; Burgess, R.L., A. Moktarzadeh & L. Cornwallis, 1966; Carruthers, D., 1909; Carruthers, D., 1949; Clark, J.L., 1964; Collias, N.E., 1956; Corbet, G.B., 1978; Danford, C. & E.R. Alston, 1877; De Blase, A.F., 1971; Ellerman, J.R., 1948; Ellerman, J.R., 1961; Ellerman, J.R. & L.L. Chaworth-Musters, 1947-48; Ellerman, J.R. & T.C.S. Morrison-Scott, 1951; Etemad, E., 1968; Felten, H., 1962; Felten, H., 1964; Geist, V., 1971; Gladkov, N.A. & A.A. Nasimovich. 1968; Corges, M., 1966; Groves, C.P., 1969; Guggisberg, C.A.W., 1975; Harrison, D.L., 1964; Heptner, M.G., 1937; Heptner, V.G., 1961; Hill, J.E., 1961; Holloway, C.L., 1975; Inskipp, T. & S. Wells, 1979; IUCN, 1978; Kus'alin, A.P., 1950; Lay, D.M., 1967; Lay, D.M., J.A.W. Anderson & J.D. Hassinger, 1970; Lindsay, H.M., 1929; Lydekker, R., 1898; Misonne, X., 1959; Nadler, C.F. et al, 1971; Nadler, C.F. et al, 1973; Nasimovitoh, A. A., 1955; Nasimovich, A.A., 1970; Omev, S.I., 1962-67; Petter, F., 1961a; Petter, F., 1961b; Pfeffer, P., 1967; Pocock, R.I., 1947; Pocock, R.I., 1951; Prater, S.H., 1965; Richter, H., 1970; Roberts, T.J., 1969; Roberts, T.J., 1977; Schaller, G.B. & S.A. Khan, 1975; Schaller, G.B. & Z.B. Mirza, 1971; Short, R.V., 1976; Siddiqui, M.S., 1961; Scholov, I.I. ed., 1963; Taber, R.D. & R.S. Hoffmann, 1963; Talbot, L.M., 1960; Thomas, 0., 1905; Thomas, O., 1912; Tsalkin, V.I., 1951; Vinogradov, B.S., 1928; Vinogradov, B.S. & A.I. Argiropula, 1941; Walker, E.P., 1964; Hard, R., 1914.

4. Ungulates

Ahlemann, H., 1970; Blyth, E., 1840; Blyth, E., 1941; Brooks, V., Brooke, B., 1875; Carruthers, D., 1909; Carruthers, D., 1949; Cotton, W.B., 1914; Cumming, J.W.N., 1905b; De La Tour, G.D., 1975; Dor, R., 1976; FAO, 1975d; FAO, 1975e; FAO, 1976b; FAO, 1977a; FAO, 1977b; FAO, 1978e; Gillan, G.V. B., 1934; Groves, C.P., 1969; Groves, C.P. & V. Mazak, 1967; Hay, R.G., 1840; Holloway, C.L., 1975; Hutton, I., 1842a; Hutton, I., 1842b; Hutton, I., 1842c; Kullmann, E., 1965; Kullmann, E., 1968; Lydekker, R., 1898; McDonald, W., 1968; Mcklenburtser, R.N., 1948; Madler, C.F. et al, 1971; Madler, C.F. et al, 1973; Naurann, C., 1973; Nauroz, M.K. & C.M. Naumann, 1975; Petooz, R.G., 1971; Petooz, R.G., 1972; Petooz, R.G., 1973a;

Petooz, R.G., 1973b; Petocz, R.G., 1973o; Pocock, R.I., 1947; Ponoins, E. D., 1895; Roberts, T., 1969; Schaller, G.B. & S.A. Khan, 1975; Schaller, G.B. & Z.B. Mirza, 1971; Short, R.V., 1976.

5. Carnivores

Abbott, J., 1884; FAO, 1973f; Gaisler, J., D. Povolny, Z. Sebek & F. Tenora, 1968; Guggisberg, C.A.W., 1975; Kullmann, E., 1965; Kullmann, E., 1967; Naumann, C. & J. Niethammer, 1975; Niethammer, G. & J. 1967b; Niethammer, J., 1966b; Pocock, R.I., 1951; Povolny, D., 1966.

6. Small Hammals

Adlen, V., 1959; Barus, V. & F. Tenora, 1967; Blanford, W.T., 1881; Ellerman, J.R. & J.L. Chaworth-Musters, 1947-48; Felten, H., 1962; Felten, H., 1964; Gaisler, J., 1970a; Gaisler, J., 1970b; Gaisler, J., 1971; Gaisler, J., 1975; Gaisler, J., D. Povolny, Z. Sebek & F. Tenora, 1967; Hanak, V. & J. Gaisler, 1969; Hassinger, J.D., 1970; Heptner, W.G., 1937; Hill, J.E., 1961; Klapperich, J., 1954; Lindberg, K., 1961; Meyer-Oehme, D., 1965; Meyer-Oehme, D., 1968; Naumann, C. & J. Niethammer, 1975; Neuhauser, H.N. & A. De Blase, 1974; Niethammer, J., 1965; Niethammer, J., 1967c; Niethammer, J., 1967d; Niethammer, J., 1969a; Niethammer, J., 1969b; Niethammer, J., 1970b; Niethammer, J., 1973a; Niethammer, J., 1973b; Niethammer, J. & J. Martens, 1975; Petocz, R.G., 1973; Petter, F., 1961; Riohter, H., 1970; Thomas, O., 1912; Zimmermann, K., 1956.

Birds

1. Afghanistan

Abbott, J., 1884; Akhtar, S.A., 1946; Akhtar, S.A., 1947; Akhtar, S.A., 1955; Aslanow, M.G., 1953; Baker, E.C.S., 1919; Barnes, H.E., 1680; Barnes, H.E., 1881; Blyth, E., 1847; Bowdler Sharpe, R., 1891; Burton, P.J.K., 1971; Buttiker, W., 1959; Carruthers, D., 1949; Christison, A.F.P., 1941; Cumming, J.H.N., 1905a; Dementiev, G., 1963; FAO, 1975a; FAO, 1975b; FAO, 1975c; FAO, 1976a; FAO, 1976e; FAO, 1976f; FAO, 1977a; FAO, 1977e; Finn, F., 1896; Griffith, H., 1947; Horsfield, T. & F. Moore, 1854; Hue, F. & R.D. Stoheoopar, 1970; Hutton, T., 1847; Johnstone, C., 1968; Kerstan, G. 1937; Klookenhoff, H. & Madel, G., 1970; Koelz, M., 1939; Koelz, 1949; Koelz, H., 1950; Koelz, H., 1954; Konig, F.J., 1972; Koning, F.J. & L.J. Dyksen, 1971; Koning, F.J. & L.J. Dyksen, 1972; Koning, F.J. & J. Rooth, 1975; Koning, F.J. & J.C. Walmsley, 1972; Kuhnert, G., 1971; Löhrl, H. & G. Thieloke, 1969; Marien, D., 1950-51; Meinertzhagen, R., 1938; Heinertzhagen, R., 1939; Hurray, J.A., 1887; Husson, C., 1974; Naumann, C., 1974; Niethammer, G., 1966; Niethammer, G., 1967; Niethammer, G., 1968; Niethammer, G., 1971; Niethammer, G., 1972; Niethammer, C., 1973; Niethammer, C. & J., 1967a; Niethammer, C. & J., 1967c; Niethammer, J., 1966a; Niethammer, J., 1967a; Niethammer, J., 1967b; Niethammer, J., 1970e; Nogge, G., 1969; Nogge, G., 1972; Mogge, G., 1972;

Nogge, G., 1973a; Nogge, G., 1973b; Nogge, G., 1974; Nogge, G. & J.
Niethammer; Orkel, G., 1972; Paludan, K., 1959; Pimms, S.L., 1968;
Potapov, R.L., 1963; Puget, A., 1969; Puget, A., 1970a; Puget, A., 1970b;
Puget, A., 1971; Reeb, F., 1977; Savage, C., 1968; Scheide, A., 1937;
Scott, D.A., 1975; Scully, J., 1887; Severtsov, N.A., 1883; Sharpe, R.B.,
1889; Sharpe, R.B., 1891; Smith, E.C., 1974; St.John, O.S., 1889; Swinhoe,
C., 1882; Thiollay, J.M., 1978; Tong, N., 1968; Vasio, V.F., 1974; Vaurie,
C., 1949-51; Viellard, J., 1969; Vigne, G.T., 1843; Wardlaw-Ramsey, R.G.,
1879-80; Whistler, H., 1944-45; Mood, B., Madge, S.C. & C.S. Waller, 1978;
Yates, C.E., 1900; Zarudny, N.A., 1926.

2. Regional

Ali, Salim, 1972; Ali, S. & S. D. Ripley, 1970; Archibald, G., 1975; Biswas, B., 1950; Burgess, R.L., A. Waktarzadeh & L. Cornwallis, 1966; Chirstison, A.F.P., 1940; Cramp, S., e.o., 1977, 80; Dementiev, G.P. & N.A. Gladkov, 1951-56; Etcheoopar, R.D., 1969; FAO, 1976d; Fulton, N.T., 1904; Fulton, H.T., 1905; Futehally, Z., J.C. Daniels & P.V. Bole, 1971; Inskipp, T. & S. Wells, 1979; Isalov, Yu. A. & T.P. Shevareva, 1967; IUCN, 1978, 79; Ivanov, 1969; Kear, J. & N. Duplaix-Hall, 1975; McClure, M.E., 1974; Peterson, R., Nountfort, G. & P.A.D. Hollum, 1974; Santapau, H., Z. Futehally & J.C. Daniels, 1969; Sohliz, E. & W. Gehlhoff, 1967; Scott, D.A., 1975; Scott, D.A. et al, 1975; Severtzov, N.A., 1887; Severtzov, N.A. & M.A. Menzbier, 1894; Stresemann, E. & L.A. Portenko, 1967; Vaurie, C., 1965; Voous, K.M., 1973, 77; Whistler, H., 1949; Whitehead, C.H.T., 1909.

3. Waterfowl

Akhtar, S.A., 1947; Brown, L.H., 1975; Burton, P.J.K., 1971; FAO, 1975b; FAO, 1976a; FAO, 1976d; FAO, 1976e; FAO, 1977e; Isakov, Yu. A. & T.P. Shevareva, 1967; Kear, J. & N. Duplaix-Hall, 1975; Klookenhoff, H. & Madel, G., 1970; Koning, F.J. & L.J. Dyksen, 1972; Koning, F.J. & L.J. Dyksen, 1971; Koning, F.J. & J. Rooth, 1975; Niethammer, G., 1972; Niethammer, J., 1970a; Nogge, G., 1969; Nogge, G., 1972; Nogge, G., 1974; Scott, D.A., 1975; Scott, D.A., 1975.

4. <u>Kigration and Bird-ringing</u>

Akhtar, S.A., 1955; Futehally, Z., I.C. Daniels & P.V. Bole, 1971; Huß, F. & R.D. Etchecopar, 1970; Isakov, Yu. A. & T.P. Shevareva, 1967; McClure, M.E., 1974; Niethammer, G., 1972; Niethammer, G. & J., 1967; Niethammer, J., 1966a; Niethammer, J., 1967a; Paludan, K., 1959; Reeb, F., 1977; Santapau, H., Z. Futehally & J.C. Daniels, 1969; Savage, C., 1968; Whistler, H., 1944-45.

Reptiles and Amphibians

Alcock, A. & F. Finn, 1897; Anderson, S.C. & A.E. Leviton, 1969; Bennikov, A.G., I.S. Darevsky, & A.K. Rustamov, 1971; Bodganov, O.P., 1962; Bogdanov, P.O., 1965;

Brück, G., 1968; Clark, R.J., E.D. Clark, S.C. Anderson & A.E. Leviton, 1969; Hemmer, A., J.F. Schmidtler & H. Böhme, 1978; TUCN, 1979; Klemmer, K., 1963; Kral, B., 1969; Leviton, A.E., 1959; Leviton, A.E. & S.A. Anderson, 1961a; Loviton, A.E. & S.A. Anderson, 1961a; Loviton, A.E. & S.A. Anderson, 1970; Pritchard, P.C.H., 1979; Schneider, P. & A.S. Djalal, 1970; Smith, M.A., 1940; Terentev, P.V. & S.A. Chernov, 1949; Ueno, S. & K. Nakamura, 1960; Wermuth, H., 1965, 67; Wettstein, O. von, 1960; Wettstein-Westersheimb, O. von, 1960.

Lower Animals

Barus, V. & F. Tenora, 1967; FAO, 1978a; Godwin-Austen, H.H., 1882-1914; Groun-Grehimailo, Gr., 1890; Heydemann, F., 1954; Kasy, F., 1965; Nadel, G., 1968; Salem, A., 1972; Salem, A., 1979; Schumakov, E., 1963; Serfaty, A. & M. Vachon; Ueno, H., 1960-65; Tashhir, A., 1947, 49; Third Danish Expedition to Central Asia; Vileshire, E.P., 1958.

8. Ellobius fuscocapillus (Blyth)

southern mole-vole

Dry stony deserts of central and northern Afghanistan between 500 and 2 600 m. Reported from Herat Province (Scully, 1887), Ghazni, Qargah, Herat, Kala Safid (C) (JN), Murghab (Scully, 1887), Paghman, Ghazni, Herat (JH).

9. Ellobius talpinus (Pallas)

northern mole-vole

E. t. talpinus. Northern loess plains below 1 300 m. Common in Mazar-i-Sharif, Maimanah (JH), Balkh, Lala Guzar (Taloqan), and Iman Sahib (Kunduz) (JN).

10. Gerbillus cheesmani Thomas

Cheesman's gerbil

G. c. aquilus Schlitter and Setzer. In sandy habitats of southern Afghanistan, below 1 000 m elevation. Abundant in Spin Baldak (SW) sand dunes, Kandahar, Girishk (S) (JH), Bakwa and Registan deserts (KH).

11. Gerbillus nanus Blanford

Baluchistan gerbil

Below 1 150 m elevation in southern sandy steppes and semi-deserts. Reported from Farah, Kang (Seistan) and Dilaram (JN), Kandahar and Qala-i-Kang (SW) (JH).

12. Tatera indica (Hardwicke)

Indian gerbil

Subtropical species of sandy steppes of the south and east. Reported from Herat, Kandahar, Girishk (S), Qala-i-Kang (SW), Jalalabad (JH), Somerkhel, Meterlam (E) (JG), Farah, Kala-Bust, Kandahar (JN).

13. Meriones crassus Sundevall

Sundevall's jird

Southern desert below 1 000 m. Gatai, Kandahar (JN), Kandahar, Girishk (S) Qala Bust and Kang (S) (JH), Bakwa, Registan and Margo deserts (KH).

14. Meriones hurrianae (Jerdon)

Indian desert gerbil

Southern Afghanistan (Corbett). Questionable reports, but probably between Ghazni and Kandahar (JH).

15. Meriones libycus Lichtenstein

Libyan gerbil

Common below 2 300 m elevation throughout all the steppes and deserts, including the Jalalabad Valley (JH).

16. Meriones meridianus (Pallas)

midday gerbil

Northern loess steppes of Afghanistan. Recorded from Kunduz (JH, JG) and below 700 m in the wetlands of Amu Darya river.

17. Meriones persicus (Blanford)

Persian jird

Mountains and foothills between 700 and 2 500 m elevation of dry central, southern and western Afghanistan. Recorded from Kabul Valley: Dare Nu, Meterlam, Darunta and Somerkhel (JG), Northern Salang, Qargah, 25 km SW of Kabul, Herat, 70 km N of Dilaram (JN), Kabul-Paghman, Shibar Pass (C), Herat, Ghazni, Kandahar (JH).

18. Meriones zarudnyi Heptner

Zarudny's jird

Northern loess steppes of Afghanistan. One record from Maimanah (JH).

19. Rhombomys opimus (Lichtenstein)

great jird

Numerous in clay and loess biotopes of northern and northwestern Afghanistan. Reported from Kunduz, Mazar-i-Sharif, Herat, Bala Murghab (NW), Sabzak Pass (NW) (JH), Balkh (MO), 60 km NW of Pul-i-Kumri, 50 km E of Bala Murghab (JN).

Family Muridae

1. Apodemus sylvaticus (Linnaeus)

field mouse

A. s. sylvaticus: Widespread in the Hindu Kush, Nuristan and Wakhan mountains on places with dense or high shade vegetation and water-courses between 2 000 and 3 000 m elevation.

2. Rattus rattoides (Hodgson)

Turkestan rat

R. r. turkestanicus (Satunin)

In central, northern and northeastern Afghanistan, semicommensal. Seasonal migration from fields in summer to villages and towns in winter. In Jalalabad valley only in the rocky areas (JN & Martens).

3. Rattus rattus (Linnaeus)

house rat

R. r. povolnyi Niethammer and Martens. In or near buildings in Jalalabad Valley and Nuristan, up to 1 200 m elevation (JN & Martens).

4. Mus musculus Linnaeus

house mouse

Commensal to man, up to 3 000 m elevation throughout Afghanistan.

5. Nesokia indica (Gray)

bandicoot rat

N. i. indica. Synanthrope, along watercourses below 2 000 m elevation throughout Afghanistan. Also Jalalabad Valley (JH).

Family Gliridae

1. Dryomys nitedula (Pallas)

forest dormouse

D. n. angelus (Thomas). Along watercourses of central and northern Afghanistan between 1 000 and 2 800 m. The animals are harmful for the irrigation works. Recorded from Paghman, Salang, Herat (Chaworth-Musters, 1939, JH), Qargah (JN), Bagrabad (S) (KP).

Family Dipotidae

1. Jaculus blanfordi (Murray)

greater three-toed jerboa

Steppe and desert regions of southern Afghanistan below 1 000 m elevation. Recorded from Kandahar, Girishk (S), Kang (S) (JH), Kandahar (JN).

2. Allactaga elater (Lichtenstein)

small five-toed jerboa

A. e. indica Gray. Steppe and desert regions between 400 and 2 500 m throughout Afghanistan. Recorded from Herat, Logar river, Ghazni, Kandahar, Girishk (S), Dilaram (S) (JN), 25 km NW of Kunduz (N) (JC), Kunduz, Mazar-i-Sharif, Herat, Ghazni, Kandahar (JH).

3. Allactaga euphratica Thomas

Euphrates jerboa

(Synonym A. williamsi Thomas)

A. e. caprimulga Ellermann. Dry montane regions of the Hindu Kush Mountains between 1 800 and 3 200 m elevation. Recorded from Bamian, Bande Amir, Dashte Nawar, Qargah, Logar, Paghman, Kabul (JN), Paghman (JH), Shibar Pass (C) (Ellerman, 1948) (JH).

4. Allactaga hotsoni Thomas

Hotson's jerboa

Bakwa and Artom deserts of southern Afghanistan. Recorded from Dilaram (S), Kang (S) (JH), Girishk (JN) (JH). Corbett however put these records under \underline{A} . euphratica.

Salpingotus michaelis Fitzgibbon

dwarf jerboa

Not yet recorded for Afghanistan, but from Nushki (Pakistan), 30 km SW of border. This species is perhaps synonymous with S. thomasi Vinogradov, type specimen from Afghanistan-Baluchistan border (JH).

Family Hystricidae

1. Hystrix indica Kerr

Indian crested porcupine.

Throughout Afghanistan below 2 600 m, along the rivers. Recorded from Kandahar, Mukur (E), Kabul, Lataband (E) (JN), Gusalek (E), Obeh (W) (KP), Jalalabad and Darre Nure Valleys (JG), Shahjoy (SE) (KH), Paghman, Shibar Pass (C), Faizabad (NE), Kandahar (JH).

ORDER CARNIVORA

Family Canidae

1. Canis aureus Linnaeus

jacka1

Widespread in the arid lowlands and mountainous steppes up to an elevation of 3 500 m. Recorded from all over Afghanistan, frequently near urbanizations. Pursued in great numbers for its skin.

2. Canis lupus Linnaeus

wolf

- C. 1. lupus. Widely distributed in all types of terrain, especially mountains. Common in the Pamirs (JN), widespread in Central Highlands, Nuristan forests, Ghazni (KH) between 1 000 and 4 600 m elevation. Pursued in great numbers for its skin. Main centra are in Mazar-i-Sharif, Maimanah and Khanabad (Ro).
- 3. <u>Vulpes</u> cana Blanford

Afghan fox

Confined to the mountainous steppes. A very rare species. One record from Kandahar (Blanford, 1881). Rodenburg reports 4 furs in Kabul Bazar.

4. Vulpes corsac (Linnaeus)

corsac fox

Dry steppes and deserts of northern Afghanistan, below 1 000 m elevation. Common and much sought after for its skin. Main centra Mazar-i-Sharif and Maimanah (Ro).

5. Vulpes rueppelli (Schinz)

sand fox

Extremely rare desert species. Probably one Afghan record from Dashte Barang, near Afghanistan-Baluchistan border (JA).

6. Vulpes vulpes (Linnaeus)

red fox

V. v. vulpes. Widely distributed throughout the country, from southern deserts to Pamir tundras, up to 4 500 m elevation. Most important Afghan fur animal (Ro).

Cuon alpinus (Pallas)

red dog, dhole

Not yet recorded for Afghanistan but occur in neighbouring Pamir (Novikov) and probably Pakistan.

Family Ursidae

1. Selenarctos thibetanus (Cuvier)

Asiatic black bear

Rare in the monsoonal forests of east Afghanistan above 1 500 m elevation. Recorded from Darunta (W of Jalalabad) (Povolny), common in the Pech-Parun Valley (E) (KP).

2. Ursus arctos Linnaeus

brown bear

U. a. isabellinus Horsfield. Alpine valleys of the Pamir and Nuristan between 2 500 and 5 000 m altitude. Strong hunting pressure (RP). In Big Pamir common (CN & JN).

Family Mustellidae

1. Mustela erminea Linnaeus

ermine, stoat

M. e. erminea. Only reported from the alpine zones of Badakhshan, Zebak, Tolibai Valley (KH) and Small Pamir (CN & JN).

2. Mustela nivalis Linnaeus

weasel

M. n. nivalis. Paghman Mountains between 1 500 and 3 500 m elevation and probably also in Pamir and Hindu Kush (JN). Recorded from Paghman, Shibar Pass (JN), 72 km NW of Kabul (JH), Weran Valley (KP).

3. Vormela peregusna (Güldenstädt)

marbled polecat

V. p. peregusna. Steppes throughout the country; uncommon. Recorded from Kandahar (Hutton), (KH), (JN), Dashte Laili (Shiberghan-Tashkurghan, N) (CN), Alinghar river (Laghman) (JG), Ab-i-Estada, Kabul (RP).

4. Martes flavigula (Boddaert)

yellow-throated marten

M. f. flavigula. Forest dwelling species of monsoonal Afghanistan.

Reported from Pech-Parun Valley (Nuristan) (KP), Kabul furshops (Ro).

5. Martes foina (Erxleben)

stone marten

M. f. foina: Montane steppes. Recorded from Qala-i-Pandja (Wakhan) (JN), Shibar Pass (C), Badakhshan (JH), Barak (Badakhshan) (CN), Ajar Valley (KH), Waygal (E) skin (RP), Kabul furshops with large numbers from Mazar-i-Sharif, Kunduz and Khanabad (Ro).

6. Mellivora capensis (Schreber)

honey badger

M. c. indica (Kerr). Drier parts of Afghanistan. Very rare. Novikov gives Hari Rud, Murghab and Amu Darya river valleys in Turkestan. Roberts, 1977 gives southern plains.

7. Meles meles (Linnaeus)

badger

Steppes of northern Afghanistan; extremely rare. Kabul fur bazar, one skin from Badakhshan (JN) and one from Nuristan (Rodenburg). A zoo animal was probably taken in Behsud (C) (CN).

8. Lutra lutra (Linnaeus)

otter

Widespread along rivers and watercourses throughout the country. Heavily hunted for their skins. Recorded from Hari Rud (near Herat), Kantiway river (near Chaman, E), Bajagul Valley (Laghman) (RP), Lindai Sin, Gusalek (E) (CN) (KP), Anjuman Pass (Pansjir), Juwain (Farah) (JH).

Family Viverridae

- 1. Paguma larvata (Hamilton-Smith) masked palm-civet Reported from Nuristan (JN in Roberts).
- 2. Herpestes auropunctatus (Hodgson) small Indian mongoose

 H. a. pallipes Blyth. Semi-deserts of Kandahar, Seistan and Eastern
 Afghanistan. Reported from Laghman and Jalalabad (JG), Herat,
 Kandahar, Girishk (C) (JH), Kandahar, Seistan, Jalalabad (KN),
 Kandahar 1953 (KZ).
- 3. Herpestes edwardsi Geoffrey Indian grey mongoose
 H. e. ferrugineus Blanford. Recorded from Afghanistan (Kullman)
 but not yet collected.

Family Hyaenidae

1. Hyaena hyaena Linnaeus

striped hyaena

 $\frac{\text{H. h. hyaena.}}{1\,500}$ Widely distributed in semi-deserts and steppes below $\frac{1}{1\,500}$ m, mainly in the southern part but probably sporadically north (JH). Also reported from lower Kabul River Valley (JG).

Family Felidae

1. Felis bengalensis Kerr

leopard cat

- F. b. trevelyani Pocock. Forested mountains of eastern Afghanistan. Kunar and Pech Valleys (E) (KP), Dare Nur (E) (CN).
- 2. Felis caracal Schreber

caracal

- F. c. schmitzi Matschie. Very rare animal of the steppes of Turkestan. Recorded from Sare Pul (Jozjan Province) and Herat (pelt) (No).
- 3. Felis chaus Guldenstädt

jungle cat

Common in riverine reed beds and swamps but also in steppes. Recorded from Maimanah (Scully, 1887), Hari Rud near Herat, Girishk, Helmand river, Kandahar (JH), Nangarhar (E) (JN), Meterlam (Laghman) (CN). Abundant on the Mazar-i-Sharif, Kunduz, Herat and Kabul fur markets (Ro). (Ro).

4. Felis lynx Linnaeus

1.ynx

- F. 1. isabellina Blyth. Alpine slopes above treeline and upland valleys, up to 4 500 m elevation in the Hindu Kush, Nuristan, Wakhan and the Pamirs. Recorded from Wakhan, Big and Small Pamirs (RP), Boroghil Valley (Sarghad, Pamir), Ghorban Valley (C), Shibar Pass (C) (CN), Upper Nuristan (KH).
- 5. Felis manul Pallas Pallas's cat

 Montane steppes between 1 500 and 3 500 m elevation. Recorded from

 Hakban Corridor and Zabak Vallay Salara Page and Parishin Vallay

Montane steppes between 1,500 and 3,500 m elevation. Recorded from Wakhan Corridor and Zebak Valley, Salang Pass and Panjshir Valley (JN).

6. Felis silvestris Schreber

wild cat

(conspecific with F. lybica Forster). F. s. ornata Gray. Desert cat of the southern and eastern desert areas of Afghanistan.
F. s. caudata Gray. Turkestan (Corbet). Recorded from Central Mountains, Shibar Pass (C), Herat (KH), Bamian (JG), Waygal Valley (E) (Kabul Zoo info.). Much sought after for its pelt and common in the Mazar-i-Sharif, Maimanah, Andkhui and Herat fur bazars (Ro).

Appendix 1

CONVENTION ON INTERNATIONAL TRADE IN ENDANGERED SPECIES OF WILD FAUNA AND FLORA (CITES)

The Convention states that for species threatened with extinction, and so included in Schedule I, an export permit from the country of origin and an import permit from the country of importation are required. These permits are issued by national management authorities, provided that removal from its country of origin will not be detrimental to the survival of the species and that the motive is not primarily commercial.

For Afghanistan the following are included in Schedule I of the Convention:

Mammals:

Selenarctos thibetanus
Ursus arctos isabellinus
Lutra lutra
Acinonyx jubatus
Felis bengalensis bengalensis
Felis caracal
Felis marmorata
Panthera pardus
Panthera tigris
Panthera uncia
Moschus moschiferus
Capra falconeri jerdoni
Capra falconeri megaceros
Ovis orientalis vignei

Birds:

Aquila heliaca Haliacetus albicilla Falco peregrinoides babylonicus Lophophorus impejanus Grus leucogeranus

Reptiles:

Varanus bengalensis Varanus greseus

Species not yet threatened with extinction but which could become endangered if trade is not controlled are included in Schedule II of the Convention. An export permit from the country of origin is required but not an import permit. For Afghanistan the following species are included in Schedule II:

7. Panthera pardus (Linnaeus)

leopard

P. p. pardus. Restricted but still occurring in the eastern and central mountain ranges. Reported from Khost Fereng (C) and Salang Mountains (C), Lataband Pass (E) (KH), Wakhan and Darqad (NE) (RP).

8. Panthera tigris (Linnaeus)

tiger

P. t. virgata Illiger. Turanian tiger, probably extinct for Afghanistan. Last reliable report from 1967 Darqad river (Takhar Province) (RP). Formerly common along the Amu Darya and Murghab rivers but during the last 50 years only recorded from the Imam Sahib and Darqad river islands (KH).

9. Panthera uncia (Schreber)

snow leopard

Arid alpine regions above treeline. Reported from Wakhan, ramirs and Nuristan (JN) (RP). Heavily hunted for its skin (Ro). Unsubstantiated report from Ajar Valley (KH).

10. Acinonyx jubatus (Schreber)

cheetah

Formerly widely scattered throughout the lower steppes up to 1 000 m elevation but now believed to be extinct (KH). Roberts, 1977 reported skins in Pakistan probably originally from Afghan Turkestan, Paludan bought a skin from Farah (1948) and Nogge from Herat (1971).

ORDER PERISSODACTYLA

Family Equidae

1. Equus hemionus Pallas

onager, wild ass

E. h. onager Boddaert. Southern Afghanistan and E. h. kulan Groves and Mazak. North-western Afghanistan. Formerly abundant on the steppes of southern and western Afghanistan but unlimited hunting eliminated most of them. Recently only recorded from the steppeland in the Herat-Badghis Provinces near the Soviet border (RP).

ORDER ARTIODACTYLA

Family Suidae

1. Sus scrofa Linnaeus

wild boar

S. s. cristata Wagner. Common in the marshes, swamps and riverbanks throughout the country, up to 2 000 m elevation.

Family Camelidae

Camelus ferus Przewalski

Bactrian camel

As a wild animal now confined to the western Gobi Desert but formerly throughout the dry steppes of Turkestan (Corbett) probably also in northern Afghanistan.

Family Moschidae

1. Moschus moschiferus Linnaeus

musk deer

Reported from the Parun Valley (E) (KP), Nisheigram and Maktosho (KH) and Kamu (E) (CN), all in the forested parts of Nuristan, up to 3 000 m elevation. It is not yet collected from Afghanistan.

Family Cervidae

1. Cervus elaphus Linnaeus

red deer

C. e. bactrianus Lydekker. Riverine woodland of northern Afghanistan. Endangered, at present restricted to the Amu Darya River wetlands of Darqad (Takhar) and Iman Sahib (Kunduz) (RP). Introduced in the Ajar Valley.

Family Bovidae

Bos mutus (Przewalski)

wild yak

Formerly as a wild species the yak probably occurred in the Pamirs.

Gazella dorcas (Linnaeus)

common gazelle, Indian gazelle

C. d. bennetti (Sykes). Pakistan. G. d. fuscifrons Blanford, Iranian Seistan. Not yet recorded for Afghanistan but possibly occurring in southern or south-western Afghanistan.

1. Gazella subgutturosa (Güldenstädt)

goitered gazelle

G. s. subgutturosa. Formerly widely distributed in the steppes and semi-deserts below 2 000 m but hunting eliminated most of the population. Recently recorded from Girishk, Kalat (NW) (JH), Registan and Margo deserts (CN), Khulm plain and Hairatan Desert (NW) (RP), Hamadan Desert (W) (Herat Livestock Company), Girishk, 25 km W Dilaram, 50 km NW Kandahar, N Farah (W), Tirpul-Islam Qala (W) (KP).

Nemorhaedes goral (Hardwicke)

goral

Not yet established in Afghanistan but hearsay reports from Nangarhar and Kunar fit with the goral. It occurs in contiguous parts of Pakistan.

2. <u>Capra aegagrus Erxleben</u>

wild goat

- C. a. blythi Hume. Widely distributed in mountainous southwest and south Afghanistan (JH), Uruzgan Mountains and headwaters of the Helmand river, north of Gulran in the Farah and Sharak in the Ghor Provinces, up to 3 500 m elevation (KH), S of Farah, Kabul (KP).
- 3. Capra falconeri (Wagner)

markhor

- C. f. cashmiriensis Lydekker. Headwaters of Alingar and Alishang rivers in Kunar and Laghman Provinces, still viable populations (RP), Kamdesh (JH). C. f. heptneri Zalkin. Darwaz Peninsula of Takhar Province (Roberts), C. f. jerdoni Hume. Springhar forests of Paktia, endangered; C. f. megaceros Hutton. From Kohe Safi to Lataband and Tange Gharu Passes in the Kabul Province (PP), endangered (IUCN Red Data Book). The markhor occur in alpine and subalpine habitat and conifer forests between 1 000 and 4 000 m elevation (KH).
- 4. Capra ibex Linnaeus

ibex

 $\frac{\text{C. i. }}{2}$ siberica (Pallas). Widespread in all Afghan mountains between $\frac{\text{C. i. }}{2}$ and 5 000 m elevation.

Pseudois nayaur Hodgson

blue sheep

Not yet reported from Afghanistan but from neighbouring Pakistan where it occurs in the Hunza district, bordering the Pamir, in the same general area as that from which the Marco Polo sheep migrate in the summer.

5. Ovis ammon Linnaeus

Marco Polo sheep

O. a. poloi Blyth. Alpine valleys of the Pamir between 3 000 and 5 000 m elevation. The overall population number is thought to be 2 500 for the Big and Small Pamir and an unknown additional number in the Waghjir Valley of the Small Pamir (RP).

6. Ovis orientalis Gmelin

urial

O. o. vignei Blyth. Widespread in the Hindu Kush Mountains between Herat, Hazarajat, Wakhan (NE), Lataband (E) and Kandahar, between 500 and 4 500 m elevation. Four different forms are recognized (KH).

Appendix 5

CHECKLIST OF THE BIRDS OF AFGHANISTAN

For the taxonomical background and sequence of the species listed in Appendixes 5 and 6 the publication of Voous, K.M., 1973, 1977, 'List of recent Holarchic bird species' is followed. Subspecific taxonomy is mainly based on Vaurie, C., 1959, 1965 'The birds of the Palaearctic Region' and on Hué, F. and Etchecopar, R.D., 1970 'Les Oiseaux du Proche et du Moyen Orient'. For abbreviations of collectors and for name locations see Appendixes 2 and 3.

Much of the information is from collections made by K. Paludan but also from collections made by R. Meirertshagen, by W. Koelz and by R. Maconachie, referred to in H. Whistler, 1944-45. This publication also gives much information on the older collections. Recent collections were made by J. Niethammer et al., but most of the recent information is based on observation and not on collections and so the chance of error is much greater than in the older information. More important recent publications are those of F. Reeb, J. Vieillard, V. Vasic, E. Smith and especially J. Niethammer et al.

In Appendix 5 a split in information before and after 1930 is made because the older information is mainly from the turn of the century or before, and several of the major collections were made after 1930. Hardly any observations are made during the winter months in the mountains and so for many factual resident birds no winter data are available. In the last century more emphasis was given to the collection of gamebirds, such as partridge, sandgrouse, woodcock and snipe. Recent information on these birds is often scarce, but many populations have been decimated, especially pheasants and bustards.

KEY

- 1 one record only
- 2 2-4 records, uncommon
- 3 several records, rather common
- 4 many records, common
- 5 breeding
- 6 not recorded since 1930
- r resident
- w winter visitor
- s summer visitor, breeding not recorded
- b summer visitor, breeding recorded
- m passage migrant
- E eastern Afghanistan
- N northern Afghanistan
- W western Afghanistan
- S southern Afghanistan
- wE winter visitor to eastern Afghanistan
- C Central Afghanistan, Hindu Kush

NON-PASSERIFORMES	•	73	**		W	т	т		C	^	M	n	
	<u>J.</u>	F.											
Tachybaptus ruficollis	3	3	4	5	5	5	5	3	3	3	3	3	b,m,w
Podiceps cristatus	3	3	3	5	5	2	•	1	•	2	2	3	b,m,w
Podiceps grisegens		2	•	5	5	5	•	•	1	•	•	•	Ъ
Podiceps auritus		1	•	•	•	•	•	•	•	•	•	•	E
Podiceps nigricollis	2	2	3	3	5	5	5	3	2	2	2	2	b,m,w
Phalacrocorax carbo		2	3	5	3	•	•		1	1	•	•	b,m
Phalacrocorax pygmaeus				•	1	•	•	•		•		•	SW
Phalacrocorax niger	6				•	•	•	•	•	•	•	6	NW
Pelecanus onocrotalus	•		2		2	2	2	2	•	•	•	•	m
Pelecanus crispus	•	•	2	2	2	•	•	•	•	•	•	•	m
Botaurus stellaris	•	6	2	2	2	2	•	•	•	1	• .	6	b,m,w
Ixobrychus minutus	•	•	•	2	2	2	5	1	•	•	•	•	b,m,w
Nycticorax nycticorax	•	1	•	2	5	1	•	•	1	•	•	•	Ъ
Ardeola ralloides	•	•	•	•	1	•	•	•	•	•	•	•	E
Egretta garzetta	•	•	2	2	3	•	•	•	•	•	•	•	m
Egretta alba	2	2	3	3	2	2	•	•	•	•	1	2	m,w
Ardea cinerea	3	3	3	4	4	4	3	3	3	3	3	3	b,m,w
Ardea purpurea	•	•	2	1				•	•	•	•	•	mSW,E
Ciconia nigra		1		2	2	2	2	•	1	•	•	•	b,m
Ciconia ciconia	•	•	1	2	2	•	•	2	•	•	•	•	m
Plegadis falcinellus	•		2	2	1	•	•	•	•	•	•	•.	m
Platalea leucorodia	•	•	3	4	4	4	3	2	2	2	•	•	m
Phoenicopterus ruber	•	•	•	5	5	5	4	4	2	2	•	•	b,m,w
Phoenicopterus minor	•	•	•	•	•	6	•	•	•	•	•	•	SW
Cygnus olor	•	•	•	•	•	•	•	•	•	•	•	•	wSW.
Cygnus cygnus	•	•	•	•		•	•		•	•	•	6	SW
Anser albifrons	•	•	•	1	•		•	•	•	•	•		E
Anser anser	2		1						•	•	1	•	m,w
Anser indicus			•		•	•	5	2		•	•	•	bNE, m
Tadorna ferruginea	1	1	3	2	2	•	5	3	2		2	2	b,m,w
Tadorna tadorna	1	6	2	2	3	5	1	1	1	_	2	2	b,m,w
Nettapus coromandelianus	•	•	•	•	•	•	1	2	•	1	•	•	mE
Anas penelope	1	2	3	4	3	1	•	•	2	2	•	1	m,w
Anas falcata	•	•	•	•	•	•	•	•	•	•	•	6	NW
Anas strepera	2	2	2	3	1	•	5	•	1	2	2	2	bNE,m,w
Anas formosa	;	:	:	:	:	•	•	:	1	•	•	•	E
Anas crecca	4	4	4	5	5	3	3	4	-			4	bE,m,w
Anas platyrhynchos	3	4	4	5	3	2	•			-		4	bE,m,w
Anas acuta	3	3	4	4	3		1				3	3	BE,m,w
Anas querquedula	•	•	3	4	3	3	3			1	•	•	m
Anas clypeata	3	3	4	4	5	2	2	3	3	3	3	3	bE,m,w
Marmaronetta angustirostris Netta rufina	1	2	6 2		1	6	:	•	:	•	•	:	m.
Aythia ferina	1 3	3	4		1 3	3	5 5		1	•		2	bNE,m,w
	1	6	3			3)	5	5	3.	4	4	bE,m,w
Aythia nyroca Aythia fuligula	3	3	3	4	1	•	•	•	•	•	•	:	bW,m,w
		3		4	•	•	•	•	•	2		2	m,w
Aythia marila	1	•	•	•	•	•	•	•	•	•	•	1	E
Melanitta fusca	1	•	•	•	•	•	•	•	•	•	•	•.	E
Bucephala clangula Mergus albellus	6	1	1	÷	•	•	•	•	•	•	•	•	mN,SW
	6	1	6	:	•	•	•	•	•	•	•	6	mE
Mergus serrator Mergus merganser	6 1	•	1	1	•	•	•	•	•	•	•	:	wE,N
Oxyura leucocephala	1.	1	2	1	•		5	•	•	•	•	2	bNE,m,w
Pernis apivorus	T	_	2		2 1	2	•	1	•	6	•	•	m,w
Elanus caeruleus	•	•	•	-	T	•	•	•	•	•	•	•	C
Milvus migrans	3	3	3	1 4	3	3	3	3	3	•	•	•	E
mrPromo	,	,	,	4	,	J	J	J		3	3	3	r,b,m,w

	J.	F.	K.	A.	K.	J.	J.	A.	8.	0.	N.	D.	
Haliaeetus leucoryphus	6							•			1	•	mSW
Haliaeetus albicilla			1									•	mSW
Gypaetus barbatus		1	3	3	3	3	3	3	3	2	1	•	r
Neophron percnopterus			3	3	3	5	3	3	3	•	1	•	b.m
Gyps bengalensis						•		1		•	•	•	c ·
Gyps indicus			1			•	•	•		•	•	•	Ċ
Gyps fulvus	3	3	3	3	3	3	3	3	3	3	3	3	r,w
Gyps himalayensis				2	1	1	1	•				•	S
Aegypius monachus	1	1	3	3	3			1	3	3	1		m
Circaetus gallicus	3	3	3	3	3	2	2	6	•	•	3	3	m
Circus aeruginosus	3	3	5	5	5	5	2	5	3	2	3	3	bN,m,w
Circus cyaneus	6	6	1	1				•			6	6	m,w
Circus macrourus	3	3	3	3				1	2		3	3	m,w
Circus pygargus			1	3	2			2			•		m
Accipiter gentilis						٠.	. •					•	capt.
Accipiter nisus	1	3	3	3	5		•	1	3	2		1	bE,m,w
Accipiter badius		6		3	1		2			•	•		bNW,m
Butaster teesa								2					E
Buteo buteo		2	1	2	2				1	3			m
Buteo rufinus	3	3	3	3	2	3	5	2	2	3	3	3	bE,m,w
Aquila clanga		2	1	3	2	1	•	1		1	1	3	m
Aquila rapax			ī		2	2	2	2	•	ī	•	•	mE,C,S
Aquila heliaca	3	3	1	Ĭ.	1	1	1	•	·		3	3	m,w
Aquila chryaetos	3	3	3	2	2	2	2	3	2	2	3	3	s,m,w
Hieraaetus pennatus	i	ĺ	2	3	-	_	2	2	ī	ī	ī	ĭ	m,w
Hieraaetus fasciatus	_	_			•	•	-	2	_	_		6	m m
Pandion haliaetus	2	2	3	· 3	•	i		•	2	2	2	2	m
Falco naumanni	2	_	3	5	3	2	•	3	2		_		bNW,m
Falco tinnunculus	3	3	3	3	4	3	5	5	3	3	3	3	b,m
Falco vespertinus	-		1	2	7					_	,	_	m m
Falco columbarius	2	i	2	2	ì	:	•	•	:	i	:	2	m
Falco subbuteo	_	•	1	2	5	3	3	3	3	3	:	-	bN,m
Falco jugger	•	i	1	2	,	2	i	2			•	•	bSE
Falco cherrug	2	2	2	2	2	ī	2	2	2	2	2	2	8,m,w
Falco pelegrinoides	1	1	•	1	2	ĩ	2	2	•	-	•	•	r
Tetraogallus himalayensis	-		:	5	2	2	3	2	2	2		•	rE,C
Alectoris chukar	•	2	2	5	5	5	2	3	2	_	6	6	r
	6	1	6	2	5	2	5	1	î	•	U	U	r
Ammoperdix griseogularis Francolinus francolinus	U	r	1	1	,	2		6	•	6	•	•	rS
	•	•	-		•	•	i	-			•	•	rS
Francolinus pondicerianus Coturnix coturnix	•	•	2	3	3	2	5	3	2	ż	2	•	b,m
	•	•	1	ì	,	2	,	,	2	2	-	•	E
Pucrasia macrolopha Lophophorus impejanus	•	•	1		•	•	•	i	i	•	•	•	rE
Phasianus colchicus	•	6	6	6	5	•	•	•		i	•	6	rnw,n
	2		O	1	,	•	•	•	•	i	•	6	W
Rallus aquaticus	2	•	6	6	i	•	•	•	•	-	•	U	m
Porzana porzana	•	6	U			•	•	•	6	•	•	•	m
Porzana parva Porzana pusilla		6	•	2	2	i	•	•	2	i	•	•	bE,m
_	•		•		6	_	•	•			•	•	NW
Crex crex	•	•	•	3	3	5	2	2	2	i	•	•	be,sw,m
Gallinula chloropus	•	•	i	-	,	J	_		1		•	•	rSW,m
Porphyrio porphyrio	4	4	4	4	5	5	4	4	4	4	4	4	-
Fulica atra	4	4	4	3	J	ر	4	2	2	4	4	4	r,m,w
Grus grus	•	•	3		•	•	•	2	4	•	•	•	m ·
Grus leucogeranus	•	•	2	2	2	•	•	•	•	•	i	• .	m ` m
Anthropoides virgo	•	•	6		۷	•	•	•	•	•	6	6	m m,ws,n
Tetrax tetrax	•	i	2	6	•	•	•	•	•	•	6	U	-
Chlamydotus undulata	i	1	1	1	•	•	•	•	•	•	U	•	m,w mN
Otis tarda	r	Ţ	1	T	•	•	•	•	•	•	•	•	TTT/

-

	ว	. 1	. N	(. A	. N	(J	J. /	l. :	5.	0.	N.	D.	
Hydrophasianus chirurgus							2		2			_		bE
Rostratula benghalensis					6						•			E,S
Haematopus ostralegus		•		1	3	1			2			•	•	m m
Himantopus himantopus				4	4	4	4			4	2		•	bE,m
Recurvirostra avosetta			. :	3	3	4	2			2	2	6	:	bE,m
Burhinus oedicnemus			. (6 :	2 :	2		2		1	1		•	r r
Cursorius cursor			L :	3 :	2 :			2	•	1			·	b,m
Glareola pratincola				. :	L	1	2							m m
Glareola glareola		, ,		. 1		1		•						m
Glareola lactea	•			5 2		• .		1	•	•				PE
Charadrius dubius		. 1	. 3				3	3	l :	3	3			bE,m,w
Charadrius hiaticula	2						•			2	1	1	1	m
Charadrius alexandrinus	1	. 1	_			٠.				l	1	• .		bE,m,w
Charadrius mongolus	•	•		-				5 1		•	•		•	bNE, m
Charadrius leschenaultii Pluvialis dominica	•	6	•			2 5	5	. 1	L.	•	•	•	•	BE, m
	•	•				•	•	•	• •	•	6	•	•	m
Pluvialis squatarola	•	:						•		•	•	•	•	E
Hoplopterus indicus Chettusia gregaria	•	6	_					5.	•		•	•		bS,m
Chettusia gregaria Chettusia leucura	•	:	:			•			•		1	•		E
Vanellus vanellus	•					6	,	. 2	2		•	•	6	m,w
Calidris alba	3	3	2	_	-		•		•		3	3	3	m,w
Calidris minuta	· 1	:	:	1							•	•	•	m
Calidris minuta Calidris temminckii	T	3	-				2				3	•		m
Calidris ferruginea	•	•	1	•	2		•				•	•		m
Calidris alpina	2		•	•	3		-				2	•	•	m
Philomachus pugnax		•	2	3				1				2	6	m,w
Limnocryptes minimus	2	6	3 6	4	4	•	2	_	-			2	•	m.
Gallinago gallinago	3	3	3	1	•	•	•					2	1	m,w
Gallinago media	_		_		2	•	•	2				3	3	bNE,m,w
Gallinago solitaria	i	•	•	•	•	•	•	•	•		?	•	•	?NE
Scolopax rusticola	1	6	6	1	•	•	•	•	2		2	•	1	w
Limosa limosa		6	2	2	2	•	•	•	•	ı	L	6	1	m.
Limosa lapponica	•	Ü	2		1	•	1	•	2	•	•	•	6	m,w
Numenius phaeopus	•	•	•	i	1	•	•	•		•		•	•	m
Numenius tenuirostris	•	•	•			•	•	•	•	•		•	•	mE
Numenius arquata	•	•	i	3	3	•	•	i	i	•		•	•	?E
Numenius madagascariensis	•	•	•	•	1	•	•			•		•	•	m,w
Tringa erythropus		•	i	2	3	i	i	•	2	i		•	i	E
Tringa totanus	ì	2	2	3	3	2	5	3	2	1	1		1 .	M Larry
Tringa stagnatilis			1	3	3	-	3	2	2	•		L	τ.	,
Tringa nebularia	1			3		2	2			•		•	•	m.
Tringa ochropus	3	3	3	4	3	2	3	3	3	3	3		2	m,w
Tringa glareola	•		2	3	3	3	5	5	3	2			1	m,w
Xenus cinereus				•	2	1	2	2	2	~				m
Actitis hypoleucos	1		3	5	3	3	3	3	3	3	2		i	ри С в — Ш
Arenaria interpres				1	3			2	2		-	•	_	bN,C,E,m
Phalaropus lobatus				2	2	2		2	2	2	•		•	m.
Larus ichtyaetus		1	2	2				1	ī	-	•		•	DD
Larus ridibundus	•	6	4	4 .	4	3	2	2	2	2	i		2	
Larus genei		1		3.	5	5		2	2	-		•	-	m,w bE,m
Larus canus		•	1							•	•		•	mE
Larus fuscus	•	•	2		1		•	•	•		•			mE
Larus argentatus	1	1	2	2	2	•	•	•	•	•				m ·
Gelochelidon nilotica		•	2	3	5	3	3	2	1		•			bE,s,m
Sterna caspia Sterna aurantii	•	,. •	1	3	•	1	1	•	•	•		•		m
Sterna hirundo	•	•	•	•	•	•	•	1	•					mE
Sterna albifrons	•	•	2	2	3	3	5	2	•.	•				bNW,E,s
and atotitons	•	•	•	2	3	3	2	2	•	•	•			8

	<u> </u>	. 1	· 1	(. <i>E</i>	. M	• J	• J	• A	. S	. 0.	N.	. D.	
Chlidonias hybrida					2 /		3 2				-		
Chlidonias nigra		,			. 1	_	_	_	•	•	•	•	m an
Chlidonias leucoptera		, ,			. 2		•		•	•	•	•	?E mE
Pterocles coronatus		,			. 2				•	•	•	•	S S
Pterocles senegallus			. 1	L.				•		•	•	6	S S
Pterocles orientalis	1	. 2		2 2			2			6	•		-
Pterocles alchata						1			•	6	•	6	bE,m,w
Columba livia	3	4	. 4	, 4	5				•	•	•	•	rNW,S rE
Columba rupestris			2	· .					•	•	•	•	bne ·
Columba leuconota									i	•	•	•	rE
Columba oenas					?			•	•	:	6	•	wN
Columba eversmanni			1			2		•	•	:	-		rN, NE, m
Columba palumbus	•		6	2			5	5	•	i	•	•	bw,c,e
Streptopelia decaocta			3			5		3	3	3	•	•	bw,c,e
Streptopelia tranquebarica					5					•	•	•	E
Streptopelia turtur					5	3	5	3	2	:	•	•	
Streptopelia orientalis					5	5	2	2	2	2	•	•	bNW,N,m
Streptopelia senegalensis	3	3	3	3	5	5	5	5	5	3	3	. 3	bE,C,m
Streptopelia chinensis					ì	•	•		,	,	J	J	r,b E
Psittacula krameri	2	2	2	2	2	2	2	2	2	2	2	ż	E
Psittacula himalayana			_	2	1		1	-	-	_	2	_	sE
Psittacula eupatria			•				•	•	•	i	•	•	SL E
Cuculus canorus		•	•	i	5	5	2	2	i	1	•	•	_
Otus brucei	•	•		6	_	,	2		1	•	•	•	b NB
Otus scops	•	•	•	6	•	•	•	•	•	•	•	•	NE
Bubo bubo	•	6	•	Ū	2	•	i	3	•	•	•	•	S
Glaucidium brodiei	·		•	i	~	•	_	5	•	1	•	•	r
Athene noctua	3	3	5	5	5	3	5	3	3	1	3	3	E
Strix aluco	•	1	•	•	6	6		_	_	_	J	3	r rE
Asio otus	2	2	2	•		v	•	•	•	•	i	2	
Asio flammeus		ī	2	6	i	•	•	•	•	2	T	1	w,m
Caprimulgus mahrattensis			_	6	6	6	6	6	6	~	•	1	w,m
Caprimulgus asiaticus			•	6		•	Ū	Ü	U	•	•	•	8
Caprimulgus europaeus	•	•	:	•	2	2	i	2	i	i	•	•	8
Caprimulgus aegypticus			i	i			-	_		-	•	•	8 ~U CU
Apus apus		6	3	4	5	4	4	4	4	2	•	•	sW,SW
Apus melba	•	2	3	3	5	3	3	3	3	2	•	•	s,m
Apus affinis	i	_	1	5	1	,	1	1	1	2	•	•	s,m
Halcyon smyrnensis	1	2	•	2	•	•	_	-	1	•	i	i	s,m,wE rE
Alcedo atthis	1	2	2	3	3	3	2	3	3	2	1	1	_
Ceryle rudis	1	3	1	1	1		-	1	1		2	-	rS,s sE
Ceryle lugubris			-	2	•	•	•	•	-	•	2	•	E
Merops orientalis		•	•	-	2	•	•	•	•	•	•	•	E
Merops superciliosus	•	•	•	2	5	3	3	3	3	6	•	•	
Merops apiaster	•	•		4	3	5	3	4	2	O	•	•	s,m
Coracias garrulus	•	•	i	3	3	3	3	3	2	•	•	•	s,m
Coracias benghalensis	•	•	•	1	6			,		•	•	•	s,m
Upupa epops	2	2	3	4	5	3	3	3	3	2	2	•	5
Jynx torquilla	1	ī	i	4	2				2		_	2 1	s,m,r
Picumnus innominatus	_	-	-			•	i	-	_		_	1	m,w
Picus squamatus	•	i		i	5	5	2	i	•	6		1	E r
Dendrocopus himalayensis		-	2	3	5	5		-	•		•	-	sE
Dendrocopus auriceps			2	2	5	5		•	•		•	•	rE
Dendrocopus leucopterus	i		-	-	2		2	i	•	i	•		rN
• -,	-	•	•	•	_	•	-	-	•	-	•	• .	T 12

_

PASSERIFORMES	J	. F	. и	. A	. и	. J	. J.	. A.	. S.	. 0	. N.	D.	
Ammomanes cincturus	2	2	2									6	8SW
Ammomanes deserti	6	2	2	3	5	3	3	3	3	3	i	1	
Alaemon alaudipes	1	2	2	3	2	2	1	1	2	2	2	1	r rS
Melanocorypha calandra	1	_			5			3				1	
Melanocorypha bimaculata	1	3	2	2	5	5	2	3	3	2	i	1	rN,m
Calandrella cinerea	2	٠	4	2	4	1		3	3	3	2	2	bN,m
Calandrella acutirostris	1	i	1	2	5	5	2	3	3	2	1	1	s,m
Calandrella rufescens	2	1	i	1	5	,	1	1	2	2	2	2	6 b ==
Calandrella raytal	_	•		_	,	•	_	_	1	_		2	b,m,w E
Galerida cristata	3	4	4	5	5	5	3	3	3	3	3	3	r
Lullula arborea	1	•	•				,	,		,		J	SW
Alauda gulgula	î	i	i	5	5	3	5	3	2	i	2	. •	r r
Alauda arvensis	i	3	i	,	5		2	2		3	2	•	-
Eremophila alpestris	3	3	1	5	5	5	5	3	3	3	1	2	s,m
Eremophila bilopha	,	_	_		_	?	-	-			1		rNE
Riparia palundicola	•	•	•	•	5	2	•	2	2	•	•	•	1772
Riparia riparia	4	2	i	4	3	2	2	2	2	2	2	•	rNE
Ptyonoprogne fuligula	4		6	4					2	2	2	2	s,m,wE
Ptyonoprogne rupestris	•	•		•	•	1	•	•	•	•	:	•	C,S
Hirundo rustica	6	1	3	5	5	3	3	3	3	3	1	•	sN,E
Hirundo smithii	р		4	5	5	5	4	4	3	2	1	6	s,m
Hirundo fluvicola	•	•	•	3	5	3	3	3	3	2	1	•	8
	•	•	•	•	1	:	:	:	•	•	•	1	E
Hirundo daurica	•	•	3	5	3	5	3	3	3	1	•	•	εN,Ε
Delichon urbica	•	2	2	3	3	3	5	3	2	•	•	•	sN,E
Anthus novaeseelandiae	•	•	•	6	6	•	•	•	•	•	•	•	S
Anthus sylvanus	•	•	•	:	•	•	1	•	•	•	•	•	E
Anthus campestris	2	2	2	5	5	•	1	1	2	2	2	2	s,m,wS
Anthus similis	•	•	•	•	2	6	•	•	•	•	•		εE
Anthus hodgsoni	•	•	•	•	•	•	1	•	1	•		•	NE
Anthus trivialis	1	6	•	4	2	2	•	1	3	2	•	1	m,w
Anthus pratensis	•	•	•	•	•	•	•	•	•	•	•	•	m
Anthus cervinus	•	•	1	1	1	•	•	•	1	•	•	•	m
Anthus roseatus	:	:	6	•	•	•	•	•	•	•	•	•	E
Anthus spinoletta	1	2	2	1	1	•	•	•	•	1	1	2	m,w
Motacilla flava	•	:	2	3	3	•	2	3		2	•	•	s,m
Motacilla citreola	:	1	3	5	5	5	5		2	•	•	•	b,m
Motacilla cinerea	1	•	2	5	5	3	2			2		1	sC,m
Motacilla alba	3	3	3	3	5	5	2	3	3	3	3	3	s,m,w
Pericrocotus ethologus	•	1	•	1	2	•	•	•	•	•	•	•	sĚ
Pycnonotus leucogenys	1	2	1	3	5	2	2	1		2	1	2	rS,E
Hypsipetes madagascariensis	•	•	6	2	•	1	•	•	2	•		•	sE.
Bombycilla garrulus	•	•	•	•	•	•	•	•	•	•			m
Hypocolius ampelinus	•	•	•	1	2	•	1	•		1	. •		m
Cinclus cinclus	•	•		2	2	2		1	2	3	1		r
Cinclus pallasii	2	2	5	5	2			3	2	2		2	r.
Troglodytes troglodytes	2	2	2		1	1	5	1	3	3	2	2	rN,E
Prunella strophiata	•	•	•	•		2	•					•	bE
Prunella fulvescens	1	1	1	1		5	5			2		•	r,bN
Prunella atrogularis	3	3	3		1			•	1	2		3	W
Prunella himalayana	•	•	l	•			2	2			1		DNE,C
Prunella collaris	2	2	2			2	1	•	•	•	2 2	2	bE,m
Cercotrichas galactotes	•	•		2	3	•	2		2				s,m ·
Erythacus rubecula	•	• .	1	•	•		•		•		•		SE
Lucinia megarhynchos	•	•				2	2				•		bN
Lucinia svecica	•	2	3	3	1	5							b,m
Lucinia pectoralis	•				•	1	•	•	•			_	E

	J.	F.	ĸ.	٨.	ĸ.	J.	J.	A.	s.	٥.	N.	D.	
Lucinia brunnea	-			2	2	2	·	1					æE
Tarsiger cyanurus					•						•	•	rE
Tarsiger hyperythrus							?		•		-	•	?C
Irania gutturalis				1	2		1	•	•	•			8N
Phoenicurus erythronotus	2	3	3	2		•	•	•	1	2	2	3	m,
Phoenicurus caeruleocephalus		2	2	2	5	5	5	1	•	2	2	3	b,1
Phoenicurus ochruros	6	2	3	5	5	5	3	3	3	3	3	6	b,1
Phoenicurus phoenicurus				2	2							•	m.
Phoenicurus frontalis	•	•	•		1						•	•	g ·
Phoenicurus erythrogaster	2	2	2	6	-	i	5	3	2	2	2	2	b,:
Rhyacornis fuliginosus	1	2	2	5	5	2	2			1	1	1	rN.
Saxicola macrorhyncha	-	•	•	6		•	•	:	•		_	_	S
Saxicola rubetra	•			i	i	i				•	• .	•	C
Saxicola torquata	6	3	3	3	5	5	2	3	3	2	• .	•	
Saxicola caprata	2	2	2	5	5	2	2	3	2	2	2	2	b,π
Oenanthe isabellina	2	1	3	5	5	3	3	3	2			2	b, m
Oenanthe oenanthe		_		2					3	3	2	1	r,b
	•	•	1		3	2 2	2	3	1	•	•	•	s,m
Oenanthe pleschanka	;	1	2	2	3		3	3	3	3	2	•	s,m
Oenanthe deserti	6	1	2	2	1	5	5	2	3	3	2	1	b,m
Oenanthe finschii	2	2	2	5	5	2	5	3	2	2	2	2	rC,
Oenanthe xamnoprymma	2	2	2	3	5	2	2	3	3	3	2	2	b,m
Oenanthe picata	2	2	3	3	3	5	3	3	3	2	2	2	b,m
Oenanthe monacha	•	•	•	•	?	•	•	•	•		•	•	?W
Oenanthe alboniger	2	2	2	2	2			•	2	2	3	2	n,w
Chaimarrornis leucocephalus	2	2	2	3	1	1	5	5	1	2	2	2	rE,
Saxicoloides fulicata	•	•	•	•	•		•	•			•	•	
Monticola saxatilis		•	6	5	5	3	3	3					b,m
Monticola cyclorhyncha				2	5	5		1					bЕ
Monticola solitarius	6	6	2	5	5	5	3	2	1	•		6	Ъ
Myiophoneus caeruleus	2	3	2	5	5	3	5	3	3	2	2	2	r
Turdus unicolor		1						•					E
Turdus merula	2	2	2	2			5			2	3	2	ьNW
Turdus rubrocanus		•				1		•					E
Turdus ruficollis	2	3	3	2					2	2	2	2	m,w
Turdus pilaris		1				•	•	•				•	E
Turdus iliacus	1				•					•	ì	1	c,s
Turdus viscivorus	1	1		5	3	2	3	1			ī	1	b,w
Enicurus scouleri	•	•	1	2	2 .	•	2	•				1	rC,l
Enicurus maculatus	•		1	2	2							_	rE
Cettia cetti	6	6		1	1	2	2	1	•		•	•	8
Cisticola juncides	•				-	-	-	-	• .		•		E
Prinia gracilis	•	i	2		2	•	•	•	1	6	2	•	rS
Prinia inornata	•	-		•	-		•	•	ī		-	•	E
Prinia criniger	i	i	i	•	•	i	i	•	-	•	i	•	E,N
Scotocerca inquieta	•	î	•	2	5	•	•	•	•	•	ī	•	bС
Locustella certhiola	•	•	•	2	,	•	•	•	•	•	-	•	C
Locustella certhicia Locustella naevia	•	•	•		;	•	•	•	•	•	•	•	-
•		•	•	1 6	1	•	•	1	1	•	•	•	ш
Acrocephalus melanopogon	•	1	2		5	•	•	1	1	•	1	•	m, bì
Acrocephalus concinens	•	•	•	•	1	•	:	•	•	•	•	• .	N
Acrocephalus agricola	•	•	•	3	2	1	1	2	3	3	•	•	m
Acrocephalus dumetorum	•	•	•	1	5	•	1	3	3	3	•	•	b,m
Acrocephalus scirpaceus	•	•	•	•	5	•	1	•	•	•	•	•	Ь
Acrocephalus stentoreus	•	•	1	3	5	3	3	3	1	•	•	•	b,m
Acrocephalus arundinaceus	•	•	•	6	•	•	•	•	•	•	•	•	MM
Hippolais pallida	•	•	•	•	5	6	•	•	•	•	•	•	ЬN
				2	-	^			•				_
Hippolais caligata Hippolais languida	•	•	•	3	5 2	2 1	2	3 2	3	•	•	•	b,m

-

Mammals:

Macaca mulatta
Canis lupus
Vulpus cana
Cuon alpinus
All Felidae species (cats)
Equus hemionus
Cervus elaphus bactrianus
Capra falconeri
Ovis ammon

Birds:

Ciconia nigra
Pelecanus crispus
Platelea leucorodia
All Falconiformes species (birds of prey)
Otis tarda
Numericus tenuirostris
All Strigiformes species (owls)

Reptiles:

All Testudinidae species (turtles) Uromastyx species All Boidae species

Plants:

All Orchidaceae species.

	J.	F.	, M	. A.	. M	. J.	. J.	. A.	S.	0.	N.	D.	
Sylvia mystacea	-		1	6	1	6	•	•	•	•	•	•	sW
Sylvia nana			2	2		•		•	1	1			m
Sylvia hortensis	•	•		2	6		2	2	1	•		•	m
Sylvia nisoria	•	•	•	•	2	•	•	1	1	•	•	•	m
Sylvia curruca	•	•	2	3	3	5	2	3	4	3	2	•	b,m
Sylvia communis	•	•	•	•	2	5	5	•	2	•	•	•	bC,N,m
Phylloscopus occipitalis	•	•	•	2	2	5	2	•	•	•			ЪE
Phylloscopus tytleri	•	•,		1	2	2	1	•	•	•	•		ЪE
Phylloscopus nitidus	•		•	1	1	•	1	1	•	2	•		m
Phylloscopus trochiloides	•		•	2	2	1	2	3	3	•	•		sNE,m
Phylloscopus magnirostris	•	•	•	1	•	•	•	•	1	•	•	•	mE
Phylloscopus proregulus	•	•	•	•	•	•	•	•	•		•	•	E
Phylloscopus subviridus	•	•	1	5	5	1	1	•	•	•	•	•	bЕ
Phylloscopus inornatus	•	•	•	2	2	•	2	3	3	2	•	•	bN,m
Phylloscopus schwarzi	•	•	•	•	1	•	•	•	•	•	•	•	N
Phylloscopus griseolus	•	•	•	5	5	2	3	2	•	•	•	•	be,ne
Phylloscopus neglectus	•	1	•	2	1		2	•	•		•	•	m
Phylloscopus sindianus	•		•	2	1		•	•	•	•			N,E
Phylloscopus collibyta	•		3	4	1		2	2	1	4	4	1	m
Regulus regulus	2	2	2		1	5			•		2	3	bE,w
Muscicapa sibirica	•		•	•	1	1	•			•			E
Muscicapa ruficauda	•	•	•	1	5	5	1	•	•	•	•		ЪE
Muscicapa striata	•	•	2	6	3	5	2	2	2	•	1		bE,m
Fidecula superciliaris	•				5	5			•				ЪE
Fidecula parva	1		2	3	•	1		1	2	3	1		m
Terpsiphone paradisi	•		•	2	5	2	2	•	1	•		•	Ъ
Panurus biarmicus	•	•	•	•	•				•		•		sS,N
Turdoides caudatus	2	2	2	2	2	2	2	2	2	2	2	2	rS
Garrulax albogularis	•	•			•				•				E
Garrulax variegatus	2	1	•	2	2	2	•		•			2	вE
Garrulax lineatus	•		2	2	2	2	2	1	•		•		sN,E
Aegithalos leucogenys	•	•	2	5	2	5		•			•		rE
Parus rubidiventris	2	•	2	2	3	5	2	1	2	1	2	2	rE,N
Parus melalophus	•	2	2	2	5	5	•		•	•	•	•	rE
Parus cyanus	1	•	1	•	1	•	1	2	•	•	1	•	rN
Parus major	2	2	2	2	5	5	2	•	•	•	•	2	r
Parus bokharensis	•	1	•	•	5	•	1	•	2	•	1	6	TNE
Sitta leucopsis	•	•	•	•	2	5	•	•	•	•	•	•	rE
Sitta europaea	•	•	1	2	5	5	1	•	•		•		rE
Sitta tephronota	2	2	2	2	2	5	2	2	2	2		2	r
Trichodroma muraria	2	2	. 2	1	•		2	2	2	2	2	2	r
Certhia himalayana	•	•	2	1	5	5	2	•	2	1	1	2	rE,NW
Remiz pendulinus	•	•	2	1	1	•						•	IR.
Nectarinia asiatica	•	•	•	2	•	•		1	•	•	•	•	sE
Zosterops palpebrosa	•	•	•	•	•	•	•		•	•	•	1	E
Oriolus oriolus	•	•	•	2	5	3	3	3	1	•	•	•	вE
Lanius isabellinus	6	2	3	3	3	3	3		3	1	2	2	b,m,w
Lanius vittatus	•	:	•	2	2	5	2	2	•	•	•	•	ь
Lanius schach	•	2	2	5	5	3	3		3	2	1		b,m
Lanius minor	•	•	•	2	5	2	3	3		•		• .	ь
Lanius excubitor	2	2	2	1.	6	•	1	1		2	2	2	m,w
Lanius senator	•	1	1	•	•	•	•	•	•	•		•	s
Dicrurus macrocercus	•	1	6	3	5	2		2	1	•	•	•	PE
Dicrurus leucophaeus	•	:	:	1	1	1	1	•	•	•			sE .
Garrulus lanceolatus	:	1	2	2	2	•	•		•	•	•	•	rE
Pica pica	3	3 .	3	5	5	3	5	3	3	3	3	3	r ·
Nucifraga caryocatactes	•	•	•	1	2	2	•	•	•	•	•	•	rE
Pyrrhocorax graculus	•	2	2	2	2	2		2	1	• .		2	rC,E,NE
Pyrrhocorax pyrrhocorax	2	2	2	5	3	3	3	3	3	3		2	r

•

	<u>J</u> ,	. F.	M.	Α.	. M	. J	. J	. s.	0.	N.	D.		
Corvus monedula	2			1						2	3	3	bn,w
Corvus splendens							_		•	•	•	•	E E
Corvus frugilegus	3	3	3	3			_			2	3	3	w
Corvus corone	3	3	3	5	5	2	2		2	3	3	3	
Corvus macrorhynchos		2	2	3	5	2			•			•	b,w(not E) rE
Corvus ruficollis						•	i	i	:	•	•	•	
Corvus corax	3	3	5	3	3			3	3	3	3	3	N,E ·
Sturnus pagodarum					5	3		2	2			,	bE
Sturnus vulgaris	3	3	4	3	5	•	ī	ĩ	•	4	4	4	
Sturnus roseus				4	3	2	4	4	4	2		-	bN,m,w b,m
Acridotheres tristis	3	3	3	3	5	5	5	3	3	3	3	3	r.m
Acridotheres ginginianus						•		•	•				S
Passer ammodendri						•	•	•	•	•	•	•	NW
Passer domesticus		2	2	4	5	5	4	3	3	3			b,m
Passer hispaniolensis	1	•	2	3	2	2	2	2	2	2	•		bN,m
Passer rutilans				?	•					. •			E
Passer moabiticus	•		6		1					6	•	•	S,NW
Passer montanus	4	4	4	4	5	5	5	5	4	4	4	4	r
Petronia brachydactyla					2			•					W
Petronia xanthocollis				2	1			•					bS
Petronia petronia	2	2	2	5	5	2	2	3	3	2	2	2	r,m
Montifringilla theresae				2	2	3	2	2	2	2			r,m
Montifringilla nivalis	2	2	2	2	5	5	3	2	2	2	2	2	r
Lonchura punctulata									1				E
Fringilla coelebs	3	3	3							2	2	3	w
Fringilla montifringilla	2	3	3	2						2		3	w
Serinus pusillus	•		2		2	5	2	2			1		r
Carduelis chloris		2	2									2	w
Carduelis spinoides													?E
Carduelis carduelis	6		1	2	5	5	3	2	2	2	1	2	r.w
Carduelis spinus	1	1									1		C,E
Carduelis cannabina	2	2	1		1	•	2	2		2	1	1	rC,N
Carduelis flavirostris	•	1		5	2	5	2	2	1	1		1	rN,NE
Loxia curvirostra:	1				•		•						wE
Leucosticte nemoricola	•	1		1	1	3	5	2	1	•	•		rC,NE,E
Leucosticte brandti	•	•		•	1	2	3	2	1				rne
Rhodopechys sanguinea	•	•	•	•	2	2	2	2	1	•			r
Rhodospiza obsoleta	6	1	2	3	3	3	5	3	2	1	2	1	r
Bucanetes mongolicus	•	•	•	6	1	5	1	2	1	1	1		r
Bucanetes githagineus	•	1	1	2	5	2	5	1	•		•		r
Carpodacus erythrinus	•	•	•	3	5	5	5	2	2	• •	•	• ,	s,m
Carpodacus synoicus	•	•	•	2	•	1	2	•	2	1	•	•	r
Carpodacus thura	•	•	•	•	•	•	•	•	•	•	•	•	bЕ
Carpodacus rhodochlamys	•	•	•	•	2	2	2	•	2	1	1	•	rE,N
Carpodacus rubicilla	•	•	•	•	•	•	3	•	•	•	•	•	rne
Mycerobas icterioides	•	•	•	:	•	6	•	•	•	•	•	•	E
Mycerobas carnipes	•	٠	•	6	•	1	1	2	_	1	• •	•	r .
Coccothraustes coccothraustes	1	•	•	6	•	•	1	•	•	1	•	•	?b
Emberiza leucocephalos	2	3	1	6	•	•	•	•	• ;	•	1	2	m,w
Emberiza citrinella	1	2	•	•	•	•	•	•	•	•	•	•	E,N
Emberiza stewarti	•	6	•	3	5	2	2	2	2	•		• .	bC,E,m
Emberiza cia	2	2	3	5.	3	3	3	2	2	2	2	2	bE, NE, m
Emberiza striolata	•	•	•	1	1	5	•	•	•	•	•	•	ЪE
Emberiza cineracea	1	•	•		•	•	•	•	•	•	•	•	E
Emberiza hortulana Emberiza buchanini	•	•	•	6 3		•	•	•	•	•	•	•	NW ·
Emberiza buchanini Emberiza fucata	•	•	•	3	5	3	2	2	2	•	•	•	b,m
Emberiza rustica	•	•	i	•	•	•	6	•	•	•	•	•	?
mmnetiva instite	•	•	1	•	•	•	•	•	•	•	•	L	E

.

	J.	F.	M.	A.	M,	J.	J.	A.	S.	0.	N.	D.	
Emberiza pusilla													wSW.E
Emberiza schoeniclus	1	1	1								1		w
Emberiza bruniceps	•	•	•	3	3	5	3	4	2	1			b.m
Emberiza melanocephala Miliaria calandra	•	•	•	6	•	•	•	•					S
miliaria calandra	2	2	•	•	5	•	•	•	•	?			bN.?r

Appendix 6

BIRDS OF AFGHANISTAN

ORDER PODICIPEDIFORMES

Family Podicipedidae

(NA).

- Tachybaptus ruficollis (Pallas) little grebe T. r. capensis Salvadori. Passage migrant and winter visitor. Breeding recorded from Kole Hashmat Khan IV-IX/65,66 (JN), V/67 (AP) and Kuhak (Seistan) VII/04 (JC). Recorded from Kole Hashmat Khan III, VI-X/79 (JS), III/78 (TE), VII/74 (FR), IV/69 (JV), br IV/65-IV/66 (JN), Qargah X/79 (JS), summer 65,66 (JN), Sarobi III/78 (TE), Darunta IV/69 (JV), VI/75 (FR), Ab-i-Estada V/79 (JS), Bande Amir IX/78 (Wassey), Hamun-i-Puzak III/78 (RP), Lab-i-Baring (SW) XII/18
- 2. Podiceps cristatus Linnaeus great crested grebe P. c. cristatus. Passage migrant and winter visitor, breeding in Seistan (KP, HE) Recorded from Kole Hashmat Khan IV-V/79 (JS) II/77 (TE), Qargah XII/64, II-III/65 (JN), X-XI/79 (JS), Sabs Au (E), juv. 1965 (JN), Sardeh IV/78 (TE), Darunta IV/69 (JV), XII-I/73-75 (FR), XII/71-VI, VIII/72 (ES), Hamun-i-Puzak III/78 (RP), II/04 (JC).
- 3. Podiceps grisegena (Boddaert) red-necked grebe P. g. grisegena. Breeding in Kole Hashmat Khan: summer/65 (JN) and elsewhere recorded from Kabul Valley II/77 (TE), Qargah II, IX/66 (JN).
- 4. Podiceps auritus (Linnaeus) Slavonian grebe P. a. auritus. Passage migrant (HE). Unsubstantiated record from Kabul Valley II/77 (TE).
- 5. Podiceps nigricollis Brehm black-necked grebe P. n. nigricollis. Passage migrant and winter visitor to lower elevations of Afghanistan, breeding at Kole Hashmat Khan: summer 65 (JN), V/67 (AP), IV/69 (JV). Recorded from Kole Hashmat Khan III-VIII/79 (JS), Qargah winter 65-66 (JN), Sarobi III/78 (TE), Jalalabad II, III, VI/66 (JN), X-XII/73-75 (FR), Kunar (E), Lashkargah (SW) X-XII/73-75 (FR) Ab-i-Estada IV/79 (JS), Bande Amir VI/76 (Wassey).

ORDER PELECANIFORMES

ramily Phalacrocoracidae

1. Phalacrocorax carbo (Linnaeus)

common cormorant

P. c. sinensis (Shaw). Summer visitor and passage migrant. Recorded from Kole Hashmat Khan V/76, III/78 (TE), IV/76 (RP), Qargah III-V/65 (JN), IV/74 (FR), X/79 (JS), Darunta (E) III-IV/72 (ES), Bande Amir IX/49, Kokcha River (N) VII/48 (KP), Arghandab (SE) III/1881 (CS), Seistan II/04 (NA), Hamun-i-Saberi III/49 (KP), Hamun-i-Puzak III/78 (RP), Lower Farah Rud III/49 (KP), Lashkargah juv. IV/74 (FR), Badghis (NW) III/1885 (CY).

- 2. Phalacrocorax pygmaeus (Pallas) pygmy cormorant

 Passage migrant, probably breeding in Seistan (KP). Recorded from Danaghori (N) V/37 (RM).
- 3. Phalacrocorax niger (Vieillot) little cormorant
 Straggler (KP). Recorded from Bala Murghab (N) XII/1884, I/1885 (JA).

Family Pelecanidae

- Pelecanus onocrotalus Linnaeus great white pelican Passage migrant. Recorded from Hamun-i-Saberi III/49 (KP), Hamun-i-Puzak III/78 (RP), Kole Hashmat Khan V-VIII/79 (JS), Kabul bazar III/72 (No), Qargah l juv. VII/74 (FR).
- Pelecanus crispus Bruch

 Passage migrant. Recorded from Hamun-i-Puzak III/78 (RP), Qargah
 IV,V/65 (JN), III/73 (FR), Kabul bazar V/65, III/66 (JN), IV/71 (No)
 Darunta IV/69 (JV).

ORDER CICONIIFORMES

Family Ardeidae

1. Botaurus stellaris (Linnaeus)

bittern

B. s. stellaris. Breeding, passage migrant and winter visitor.

Reported from Kole Hashmat Khan IV-V/79 (JS), br V/67 (AP), Kabul
bazar III/73 (FR), II/80 (JS), IV/37 (RM), N of Baghlan summer/78 (Ti),
Kunar (E) X/73 (FR), Kandahar III/1881 (CS), Bala Murghab (N) XII/1884,
Badghis (NW) II/1885 (JA), Farah Rud (W) XII/04 (JC), Bagrami
III/34 (Ma).

2. <u>Ixobrychus minutus</u> (Linnaeus)

little bittern

I. m. minutus. Passage migrant and winter visitor. Breeding known from Lashkargah VII/73 (FR), probably Kole Hashmat Khan VI/72 (ES). Recorded from Kole Hashmat Khan V-IX/79 (JS), Kunar (E) V/73, V,VII/74 (FR) Kandahar IV,VI/1879 (St.J) Kwaja Ali (SW) IV/03 (JC), Faizabad (SW) IV/49 (KP), Bagrami V/34 (Ma), Panjao (C) VI/49, Chasma-i-Sher (N) VIII/49 (KP).

- 3. Nycticorax nycticorax (Linnaeus) night heron

 N. n. nycticorax. Breeding bird and winter visitor (KP). Recorded from Kabul IX/79 (JS), bazar IV/71 (No), Farah V/48, Shin Dand (S) VI/48, Lashkargah (KP), IV/73, juv. II/75 (FR), Jalalabad V/34 (Ma).
- 4. Ardeola ralloides (Scopoli) squacco heron
 Probably in Afghan Seistan and Turkestan (HE). Recorded from Qargah
 V/74 (FR).
- 5. Egretta garzetta (Linnaeus) little egret

 E. g. garzetta. Passage migrant. Recorded from Kole Hashmat Khan

 V/79, Ab-i-Estada V/79 (JS), V/79, Dashte Nawar V/76 (TE),

 Hamun-i-Puzak III/78 (RP), Sarobi III/78, Gozara (NW) IV/72, Qargah

 IV/72 (ES).
- 6. Egretta alba (Linnaeus)

 E. a. alba. Passage migrant and winter visitor. Recorded from Kole Hashmat Khan III-VI/79 (JS), III/78 (TE), VI/33 (Ma), Qargah XI-XII/79, Ab-i-Estada IV/69 (JV), IV/79 (JS), V/76 (TE). III/77 (Savey), Mukur IV/69 (JV), Sardy Lake I/79 (JS), Dashte Nawar (E) IV/69, Hamun-i-Puzak III/78 (RP), Farah IV/49 (KP), Naghlu IV/69, Darunta IV/69 (JV), Jalalabad, Sarobi, Mazar-i-Sharif, Ghazni XII-VI/73-75 (FR), Kandahar II/1881 (CS).
- 7. Ardea cinerea Linnaeus grey heron

 A. C. cinerea. Common passage migrant and winter visitor. Recorded from all wetlands throughout the country. The only heron breeding colony is in Charbujak (Seistan) XI/71 (ES). Project records from Kole Hashmat Khan III-IX/79 (JS), Ab-i-Estada I/79 (JS).
- 8. Ardea purpurea Linnaeus purple heron

 A. p. purpurea. Passage migrant. Recorded from Seistan III/49, Farah

 TV/49 (KP), Qargah III/66 (JN).

Family Ciconiidae

Passage migrant, probably breeding in Bamian (ES), in E Hazarajat (KP) and the high areas of Kandahar and Zabul (ES). Recorded from Panjao VI-VII/49, Doab VIII/49 (KP), Ajar Valley V-VI/77 (TE), Jam (C) V/77 (RP), Qargah IK/64 (JN), Jalalabad IV/69 (JV) Kandahar-Kalat V/74 (FR), Bande Amir VII/74, Surkhab (W) II/72, Qalat, Lake Dahla, Maruf (SE) VI/72 (ES), Bamian IV/37 (RM).

Ciconia ciconia (Linnaeus) white stork
 C. c. asiatica Severtzov. Passage migrant. Recorded from Salang Pass VIII/65 (JN), Kole Hashmat Khan V, VII/79 (JS), Dashte Nawar V/77, north of Charikar several captured, spring 64/65 (JN), Kabul (bazar) IV/71, III/72 (No), Darunta IV/69 (JV), Samangan (N) V/75 (FR).

Family Threskiornidae

1. Plegadis falcinellus Linnaeus glossy ibis
P. f. falcinellus. Passage migrant. Recorded from Kole Hashmat Khan
III/66 (JN), Kabul V/37 (RM), Darunta IV/69, Ab-i-Estada IV/69 (JV),
Pushut (E) III/1840 (St.J).

2. Platalea leucorodia Linnaeus

spoonbil1

P. 1. leucorodia. Passage migrant and winter visitor. Recorded from Kole Hashmat Khan IV/IX/79 (JS), IX,X/64, IV/65 (JN), Kabul Valley IV/77, Qargah IV,VII,X/74, Ab-i-Estada V/76 (TE), III/77 (Savey), IV,V/79 (JS), VI/77 (Crook), IV/69, Darunta up to 130 III-VI/72 (ES), IV/69, Najhlu (E) IV/69 (JV), Qala-i-Nau (NW) III/1885 (JA).

ORDER PHOENICOPTERIFORMES

Family Phoenicopteridae

1. Phoenicopterus ruber Linnaeus

flamingo

P. r. roseus Pallas. Scarce passage migrant and winter visitor, breeding in Ab-i-Estada and Dashte Nawar. Recent records from Ab-i-Estada: VI/69 5 100 (JV), IV/79 3.000+, V/79 6 000+ (JS) Dashte Nawar V/77 3 000 (TE). Other records from Kole Hashmat Khan IX-X/64, Qargah IX-X/64, Sarobi IX/61 (JN) Kuhak (SW) IV/05 (JC).

2. Phoenicopterus minor Geoffroy lesser flamingo Rare straggler (KP). Recorded from Kuhak (SW) VI/04 (JC).

ORDER ANSERIFORMES

Family Anatidae

- 1. Cygnus olor (Gmelin) mute swan
 Breeding in Seistan and winter visitor (KP).
- Cygnus cygnus (Linnaeus) whooper swan
 Winter visitor in Seistan (KP). Recorded from Hamun-i-Saberi XII/04 (JC).
- 3. Anser albifrons (Scopoli) white-fronted goose

 A. a. albifrons. Rare passage migrant (KP). Recorded from Darunta
- Anser anser (Linnaeus)

 A. a. rubrirostris Swinhoe. Breeding in Seistan, winter visitor and passage migrant. Recorded from Hamun-i-Saberi III/49 (KP), XII/04, III/05 (JC), Hamun-i-Puzak III/78 (RP), Kabul bazar III/65 (No), Charburjak (SW) 150 XI/71, Juwien (SW) 1/72, Lokhi (SW) 1/72 (ES).
- 5. Anser indicus (Latham)

 Breeding in Wahan and Pamir, passage migrant. Recorded from Zor Kol

 (Big Pamir) VIII/72 (JN), 72 (RP), Kol-e-Tshaqmaqtin (Small Pamir)

 VII/71 (No) VIII/72 (JN).
- Common passage migrant and winter visitor to all lakes and wetlands of Afghanistan. Breeding recorded at Hari Rud River 1884 (JA), Kole Hashmat Khan IV/67 (AP) Kol-e-Tsaqmaqtin (Pamir) VII/71 (No), Dashte Nawar and probably Ab-i-Estada (JN). Project records from Kole Hashmat Khan III/79 (JS), III/76 (RP), Taimani Wat (E) III/80 (JS), Ab-i-Estada IV/76, Hammin-i-Puzak III/78 (RP).

7. Tadorna tadorna (Linnaeus)

(JS).

shelduck

Passage migrant and winter visitor and breeding in Ab-i-Estada 64, 65 (JN) Seistan IV/04 (JC). Recorded from Ab-i-Estada, I, IV, V/79 (IV: 1 000+) (JS), VI/77 (Crook), V/76 (TE), III/77 1 000+ (Savey), Dashte Nawar V/76, 77 (TE), Qargah V-XII/64, 65 (JN) Oxus river (N) II/1884 (CY), Lab-i-Baring XII/18 (NA), Kabul XI/34 (Ma).

- 8. Nettapus coromandelianus (Gmelin) cotton teal N. c. coromandelianus. Recorded from Qargah VII/73 (FR), VIII/73 (ES), VIII/64, Jalalabad VIII/66 (JN), Sarobi X/71 (ES).
- 9. Anas penelope Linnaeus wigeon Common passage migrant and winter visitor to all Afghan wetlands. Recorded from January till June and September till December. Project records from Kole Hashmat Khan V-VI, IX-X/79, Qargah X-XII/79, II/80
- 10. Anas falcata Georgi falcated teal Straggler to SW Afghanistan (HE). Recorded from Bala Murghab (N) XII/1884 (JA).
- 11. Anas strepera Linnaeus gadwall Passage migrant and winter visitor. Breeding in Wakhan and probably

western Afghanistan. Recorded from Kole Hashmat Khan V, IX-X/79 (JS), XII/64-I-III/65 (JN), Qargah X-XII/79, Sardy Lake I/79, Ab-i-Estada IV/79 (JS), Dashte Nawar V/76 (TE), Wakhan, Small Pamir VII/71 (No), Kandahar I-III/1881 (CS), Bala Murghab (N) II/1885 (JA), Seistan III/05 (JC).

- 12. Anas formosa Georgi Baikal teal The only record for Afghanistan is from Sarobi (E), 3 couples IX/71 (ES).
- 13. Anas crecca Linnaeus A. c. crecca. Very common passage migrant and winter visitor to Afghan wetlands. Breeding in Kole Hashmat Khan IV-VI/69 (AP). Project reports from: Hamun-i-Puzak III/78 (RP), Ab-i-Estada I, IV/79, Sardy Lake I/79 (JS), Kole Hashmat Khan III-VI/79, 400+ VIII-IX/79, Qargah X/79-II/80 (JS).
- 14. Anas platyrhynchos Linnaeus mallard A. p. platyrhynchos. Common passage migrant and winter visitor to Afghan wetlands, especially in Seistan and around Kabul. Breeding bird of Kole Hashmat Khan IV/67 (AP). Project reports from Kole Hashmat Khan IV-VI/79, VIII-IX/79, Qargah 300+ X/79-II/80, Ab-i-Estada I, IV, V/79, Ajar Valley III/79, Sardy Lake (E) I/79 (JS), Hamun-i-Puzak III/78 (RP).
- 15. Anas acuta Linnaeus A. a. acuta. Common passage migrant and winter visitor to Afghan wetlands. Breeding in Kole Hashmat Khan VII/67, IX/69 (AP). Project reports from Hamun-i-Puzak III/78 (RP), Ab-i-Estada I, IV, V/79, Kole Hashmat Khan III-VII, IX-X/79, Qargah X-XII/79 (JS).

16. Anas querquedula innaeus

garganey

Passage migrant. Recorded from Kole Hashmat Khan IV-V, VIII/79 (JS). VII-IX/64-65, 11I-VI/64-65 (JN), IV/69 (JV), Sarobi III/78 (TE), Daronta (E) IV/69 (JV), Ab-i-Estada IV/69 (JV), IV, V/79 (JS), Dashte Nawar V, VI/78 (TE), Panjah (Wakhan) IV/1874 (Bi), Darah-e-Shaur (Big Pamir) VIII/72 (JN), Bamian IX-X/49 (KP), Naghlu (E) IV/69 (JV), Kamdesh (E) IV/69 (JV), Mazar-i-Sharif, Baghlan, Daronta, Qargah III-VIII/73-75 (FR), Kandahar III/1881 (CS).

17. Anas clypeata Linnaeus

shoveler

Common passage migrant and winter visitor. Breeding in Kole Hashmat Khan V, VII/67 (AP) and probably Seistan (JC). Recorded throughout the country. Project reports from Kole Hashmat Khan III-X/79, Ab-i-Estada up to 600+ I, IV, V/79 (JS), Sardy Lake I/79 (JS), Zor Kol (Big Pamir) VIII/72, Hamun-i-Puzak III/78 (RP).

- Passage migrant and breeding in Seistan (Kuhak) IV-V/05 (JC) and probably Turkestan (KP). Recorded from Kole Hashmat Khan V/65 (JN), Kandahar III/1881 (CS), De-Kamran De Doda (JA), Kham-i-ab (Oxus R) VI/1884 (CY).
- 19. Netta rufina (Pallas)

red-crested pochard

Passage migrant and winter visitor, breeding in the Pamirs and in Kole Hashmat Khan (JV). Recorded from Kole Hashmat Khan IV/69 (JV), Qargah XII/65-I, IX/66 (JN), XII/73 (FR), Sarkhad (Wakhan) VII/71 (JN), Sarobi II/77 (TE), Ab-i-Estada V/79 (JS), IV/69, Daronta IV/69, Naghlu IV/69 (JV), Kandahar III/1881 (CS), Lab-i-Baring (SW) XII/18 (NA).

20. Aythia ferina (Linnaeus)

pochard

Common passage migrant and winter visitor. Breeding in Kole Hashmat Khan VII-VIII/65 (JN), V/67 (AP). Recorded in considerable numbers from all wetlands. Project records Kole Hashmat Khan up to 2 500+ III-X/79, Qargah X-XII/79 (JS), Bande Amir IX/78 (Wassey), Hamun-i-Puzak III/78 (RP), Ab-i-Estada IV-V/79 (JS).

21. Aythia nyroca (Glildenstädt)

white-eyed pochard

Passage migrant and winter visitor, breeding in Seistan and the Hari Rud Valley (HE). Used to be common in the Kandahar area II/1881 (CS), 1878-1881 (St.J). Elsewhere recorded from Kole Hashmat Khan X/65, III/66 (JN), IV/69 (JV), Qargah I, III, IX/66 (JN), IV/74 (FR), Kila Wali (Murghab, N) III/1885, Maruchak (N), III/1885 (LY), Ab-i-Estada V/76 (RP).

22. Aythia fuligula (Linnaeus)

tufted duck

Passage migrant and winter visitor. Recorded from Kole Hashmat Khan IV/79 (JS), V/76 (TE), III-IV/76 (RP), X-IV/64-66 (JN), IV/69 (JV), Qargah X-XI/79 (JS), X-IV/64-66 (JN), X-IV/73-75 (FR), Sardeh IV/78, Sarobi III/78, Kabul Valley II/78, Dashte Nawar V/76 (TE), Daronta IV/69, (JV), III/73, II/75 (FR), Hamun-i-Puzak III/78 (RP), Kandahar II/1881, Kokeran III/1881 (CS), Maimanah II/1885 (CY).

23. Aythia marila (Linnaeus)

greater scaub

The only record is one pair from Qargah 20/XII/73 (FR).

24. Melanitta fusca (Linnaeus) velvet scooter
M. f. fusca. One record from Qargah lake 26/I/73 (FR).

- 25. Bucephala clangula (Linnaeus) golden eye

 B. c. clangula. Rare passage migrant. Recorded from Qargah II/65 (JN),
 Farah Rud (Seistan) III/49 (KP), Seistan I/05 (JC), Bala Murghab
 I/1885 (JA), Maruchak (N) III/1885 (CY), Chahar Shamba (N) II/1885 (CY).
- Rare passage migrant in eastern Afghanistan. Recorded from Kabul river west of Jalalabad II/48 (KP), Argandab river near Kokeran (SE) I/1881, (CY).
- 27. Mergus serrator (Linnaeus) red breasted merganser
 Rare winter visitor. Recorded from Daronta IV/69 (JV), Sarobi 5/III/73
 (FR) and Bala Murghab (N) I/1885 (JA).
- 28. Mergus merganser Linnaeus goosander

 M. m. comatus Salvadori. Breeding in Wakhan, rare passage migrant and winter visitor (HE). Recorded from Sarkhad (Wakhan) VII/71, Kol-e-Tshaqmaqtin (Small Pamir) VII/71 (No), Qargah II-III/65, I/66 (JN), XII/79 (JS), IV, XII/74 Daronta I. II/75 (FR).
- 29. Oxyura leucocephala (Scopoli) white headed duck

 Scarce passage migrant and winter visitor, breeds in Seistan (HE).

 Recorded from Kole Hashmat Khan IV/69 (JV), III, VI, VIII/79 (JS),

 VI/77 (Crook), III/78 (TE), Ab-i-Estada V/77 (Savey), Hamuns (Seistan)

 common during winter 1904/05 (JC), Jameh river near Khelat-i-Ghilzai

 X/1879 (St.J), Maruchak III/1885 (CY).

ORDER ACCIPITRIFORMES

Family Accipitridae

- 1. Pernis apivorus (Linnaeus) honey buzzard
 Only once recorded, from the Shibar Pass (C) 29 migrating V/77
 (Thiollay).
- 2. Elanus caeruleus (Desfontaines) black winged kite

 E. c. caeruleus. Few records from Afghanistan: Kabul (Whistler, 1945),

 Pech valley, Wama (E) IV/48 (KP).
- 3. Milvus migrans (Boddaert) black kite

 M. m. migrans. Breeding up to 3 000 m elevation; M. m. lineatus

 (Gray), common passage migrant, M. m. govinda Sykes straggler (KP).

 Recorded throughout the country. Project records Kole Hashmat Khan:

 III-V/79, IX-X/79 (JS).
- 4. Haliaectus leucoryphus (Pallas)

 Passage migrant in Seistan (HE). Recorded from Hamun-i-Saberi I/05
 (JC), Charburjak (SW) XI/71 (ES), Ab-i-Estada V/77 (Thiollay).
- 5. Haliaectus albicilla (Linnaeus) white-tailed eagle Passage migrant and winter visitor. Recorded from Hamun-i-Saberi III/49 (KP).

Appendix 2

GAZETTEER OF LOCATIONS IN AFGHANISTAN

The list includes many of the locations mentioned in the lists of mammals, birds, reptiles and amphibians. For many of the locations only approximate longitudes and latitudes could be given and some could not even be traced on the available maps.

Abakhan Valley NE Afghanistan Ab Gatsch = Abgaj 37000' 72037' Ab-i-Estada 32032' 67057' Adreskan 33°35' 62°10' Aftao + 35062' Ajar Valley = Hadjar V 35025'. 76030' Akcha 36°57' 66°06' Akrobat Valley 34057' 67040' Aksu Small Pamir Alingar R Nuristan Amu Darya R NE boundary Andkhui 36°56' 65°05' Anjuman Pass 35050' 70007' Aq Chah near Mazar-i-Sharif Archy 37005' 69028' Argandab 32030' 67008' Arghestan 31°33' 66°33' Ashraf Central Afghanistan Atin Jalao Kokcha R Baba Wali 31039' 65040' Badakhshan NE Province Badghis region 35°50' 62°00' Bagame Valley Nuristan Baghlan 36011' 68044' Baghrami 34030' 69015' Bagrabad = Baqrabad Bala Murghab = Murghab 35°35' 63°20' Balkh 36047' 66055'

Bamian = Bamiyan 34050' 67053' Bande Amir 34°50' 67°10' Bande Turkestan area north of the Hindu Kush Baqrabad 31°45' 61°00' Barfak 68°08' 35°25' Barikot 35°18' 71°36' Basawal = Bharawal BashGal = Bashgul Valley Bash Gul Valley tributary Kunar River Bassoolah SE Afghanistan Bharawal 34013' 70053' Bel Chiragh 35050' 65011' Big Pamir region 37°10' 73°00' Bisut 70030' 34034' Bokan 63032' 35043' Bolan Cave 31055' 66058' Burchao Pass 35º36' 64º37' Bust = Bost = Qala Bist 31028' 64021' Byan Khel SE Afghanistan Chagai Sarai 34048' 71005' Chah-i-Ab 37025' 69050' Chah-i-Angir 30°10' 65°55' Chahar Shamba 35°45' 63°58' Chaharasia central Afghanistan Chailtan = Chohiltan central Afghanistan Chakar Darra 34039' 69000' Chakhcharan 34028' 65001'

6. Gypaetus barbatus (Linnaeus)

bearded vulture, lammergeier =

G. b. aureus (Hablizl). Resident and common, but in small numbers in all mountain ranges, also in Nuristan and the Pamirs.

- Neophron percnopterus (Linnaeus)

 Regyptian vulture

 N. p. percnopterus. Common summer visitor and breeding throughout Afghanistan, except for the highest parts. The Afghan population partly migrates to NW India during the winter (KP). Many records throughout Afghanistan between March and September, most numerous in the northwest. Winter visitor of Seistan XI/71 (ES).
- 8. Gyps bengalensis (Gmelin) white backed vulture
 Breeding in southern Afghanistan (HE). It followed the army in 1878/
 VIII/72 (VV). Recorded from Panjao (C)
- 9. Gyps indicus (Scopoli) long-billed vulture
 Only once recorded from Pul-i-Khumri (C) III/72 (ES).
- 10. Gyps fulvus (Hablizl)

 G. f. fulvus. Common throughout the country, sometimes in large groups, breeding and winter visitor.
- 11. Gyps himalayensis Hume

 Sometimes considered conspecific with G. fulvus (in Voous, 1972).

 Summer visitor to eastern Afghanistan. Recorded from Jalalabad IV/69 (JV), Sarobi IV-VI/72 (ES), Salang Pass VII/70 (SM) (western-most
- 12. Aegypius monachus (Linnaeus) black vulture
 Breeding in southern Afghanistan (HE), passage migrant. Recorded
 throughout the country, especially in the northern and eastern parts,
 between September and May. Summer records from Hamun-i-Saberi (SW)
 VI/49 (KP), Murghab (N) VIII/72 (VV).
- 13. Circaetus gallicus (Gmelin)

 C. g. gallicus. Passage migrant. Recorded from Kabul VII/72, Ghazni VI/72, Charkh (E) VII/72 (ES), Sarobi IV/69, Chagha Sarai IV/69, Mukur IV/69 (JV), Kandahar VIII/1879 (St.J), VI/72, Zabul VI/72 (ES), Salang S, Baghlan, Pol-i-Chamri (N) III-V/73-75 (FR), Ghorband IV/37, Shibar Pass IV/37, Bamian IV/37 (RM).
- Circus aeruginosus (Linnaeus)

 C. a. aeruginosus. Throughout the year recorded from all Afghan wetlands. Locally breeding between III-VI/73-75 in the north; Kunduz (FR), Haibak (N) VIII/49 (KP). Project records from Kole Hashmat Khan IV. V/79 (JS).
- 15. Circus cyaneus (Linnaeus)

 C· C· Cyaneus . Passage migrant and winter visitor. Recorded from Qargah IV/66 (JN), Ab-i-Estada III/77 (Savey), Murghab XII/1884 (JA), III/1885, Sulfikar (Badghis) XI/1885, Chahar Shamba (N) II/1885 (CY), Kandahar I/1879 (St.J).

16. Circus macrourus (Gmelin)

Passage migrant and winter visitor. Recorded from many places. between September and April, especially April Taloqan-Faizabad (NE) 30/VIII/72, VV). (earliest record

- 17. Circus pygargus (Linnaeus) Passage migrant. Recorded from Ab-i-Estada V/79 (JS), IV/69 (JV), Kunduz IV/78 (TE), Mukur IV/69, Herat IV/69 (JV), Ghazni, Kabul IV/73-75 (FR), Murghab IV/1885 (CY), Barfak V/37 (RM), Logar III/34 (Ma), Farsi (C), Qala-i-Nau (N), Daulatabad (N) VIII/72 (VV).
- Accipiter gentilis (Linnaeus) FR recorded captured specimens from Charikar IV/73, II/74 and in
- Accipter nisus (Linnaeus) A. n. nisosimilis (Tickel). Passage migrant and winter visitor throughout the country, breeding in Nuristan (JV). Project records from Paghman, summer 79, Qargah XII/79-[[/80 (JS), West Nuristan VIII-IX/77 (RP), A. n. nisus also passage migrant (HE).
- 20. Accipter badius (Gmelin) A. b. cenchroides (Severtzov). Passage migrant, breeding in NW Afghanistan (KP). Recorded from Kandahar IV/1881 (St.J), (CS), Chahar Shamba (N) II/1885 (CY), Ghorband (C) IV/37, Haibak (N) V/37 (RM), Haibak, Maimanah VII, VIII/49 (KP), II/1885 (CY), Khyber, Barikot Kamu (E), Kabul (JV).
- 21. Butaster teesa (Franklin) Two records only from Kamu Valley (E) VIII/63 (JN) and Kole Hashmat white-eyed buzzard eagle 22.
- Buteo buteo (Linnaeus) B. b. vulpinus (Gloger). Passage migrant. Recorded from Jalalabad-Sarobi IV/69, Kandahar IV/69 (JV), X/73-75 (FR), Ab-i-Estada IV/69 (JV), V/76 (TE), III/77 (Savey), Pol-i-Chamri (N), Baghlan, Kunduz, Mazar-i-Sharif X/73-75, Ghazni II, X/73-75, Samangan V, X/73-75 (FR),
- Buteo rufinus (Cretzschmar) B. r. rufinus. Passage migrant, breeding in Muristan (KP), Safed Koh (WR) and probably Badakhshan. Many records throughout the country outside the high mountains, most common in open areas. Project records Ab-i-Estada I, V/79, Kabul I-III/80 (JS).
- 24. Aquila clanga Pallas Passage migrant, usually solitary. Recorded from Kabul V/73, XII/74, IV/75 (FR), Qargah VIII/71, Sarobi XI/71 (ES), Jalalabad IV/69, Mukur IV/69 (JV), Ab-i-Estada III/77 (Savey), Bamian X/49 (KP), Kandahar II/75 (FR), II/1879 (St.J), Qalat (SE) XII/71, Arghestan (SE) VI/72, Juwien (SW) XII/71 (ES), Danaghori (N) V/37 (RM).
- 25. Aquila rapax (Temminck) A. r. nipalensis Hodgson. Passage migrant in eastern, central and southern Afghanistan. Recorded from Kole Hashmat Khan V/76 (TE), Qargah VI/75 (FR), Ab-i-Estada III/77 (Savey), Bande Amir X/49 (KP), VII/71, 72 (ES), Ghazni-Kandahar VI/72 (ES), Panjao (C), Jam, Farsi (C) VIII/72 (VV), Danaghori (N) V/37 (RM).

26. Aquila heliaca Savigny

imperial eagle

A. h. heliaca. Passage migrant and winter visitor. Recorded from Kabul II/72, Ghazni XI/71, Farah, Nimroz winter 71-72 (ES), Girishk (S) II/49 (KP), Kandahar winter 1882 (CS), Mazar-i-Sharif, Pol-i-Chamri (N), Baghlan (N) and Lashkargah (S) XI-II/73-75 (FR).

27. Aquila chrysaetos (Linnaeus)

golden eagle

A. c. daphanea Severtzov. Common passage migrant and winter visitor but found throughout the year in all mountainous areas although probably not breeding. Project records from Kole Hashmat Khan VIII/79, Ajar Valley III/79, Ab-i-Estada V/79 (JS), Herat Province V/77, W Nuristan VIII-IX/77 (RP).

28. Hierasetus pennatus (Gmelin)

booted eagle

H. p. pennatus. Winter visitor and passage migrant, breeding in Central Afghanistan and Nuristan (HE). Recorded from Nuristan: Pech Valley III-IV, VIII/48 (KP), Sarobi, Kamdesh IV/69 (JV), Hindu Kush VIII/68 (JV), Ghazni VII/71, Mukur VI/72 and elsewhere (18 records between) VIII/71-IV/72 (ES), Ashraf (C) VIII/38 (Ma), Bamian, Ab-i-Estada, Bande Amir, Herat V/77 (Thiollay).

29. Hieraaetus fasciatus Vieillot

Bonelli's eagle

H. f. fasciatus. Passage migrant. Recorded from Bala Murghab XII/1884 (Clarke) and Bamian VIII/72, Sia Koh (C) VIII/72, Kokcha R.

Family Pandionidae

Pandion haliaetus (Linnaeus)

osprey

P. h. haliaetus. Passage migrant. Recorded from Kabul, Ghazni, Kundur $\overline{X}I-\overline{V}/7\overline{3}-75$ (FR), Kargah IX/65, IV/66 (JN), Kabul River basin IX/71-V/72, one in I/72, Enjil (NW) IV/72 (ES), Pashat (E) II/1840 (Griffith), Pech R (E) III/48 (KP), Sarobi (E) VI/77 (Crook), III/78 (TE), Sardeh IV/78 (TE), Daronta IV/69 (JV), Jalalabad XI-V/73-75 (FR), Kandahar IV/69

ORDER FALCONIFORMES

Family Falconidae

Falco naumanni Fleischer

lesser kestrel

F. n. naumanni. Passage migrant, breeding in colonies in northwestern Afghanistan: Gozara IV/72, Khusk IV/72 (ES). Recorded from III-V(-VI) and from VIII(-IX) throughout Afghanistan. Project records from Hamun-i-Puzak III/78 and Herat V/77 (RP), Bande Amir IX/78

Falco tinnunculus Linnaeus

kestrel

F. t. tinnunculus. Passage migrant, F. t. stegmanni (Portenko) breeding and passage migrant. Widespread throughout Afghanistan and many records. Project records from Ab-i-Estada I, IV/79, Kabul I/79 (JS), Herat V/77, Hamun-i-Puzak III/78 (RP).

3. Falco vespertinus Linnaeus

red-footed falcon

F. v. vespertinus. Passage migrant. Recorded from Herat, Farah III-IV/72 (ES) and Bamian many IV/37 (RM).

Falco columbarius Linnaeus

- F. c. insignis (Clark). Passage migrant. F. c. pallidus (Sushkin). Passage migrant of southern Afghanistan (HE). Recorded from Kabul II/80 (JS), IV/78 (TE), XII/74 (FR), Kunduz IV/78 (TE), Logar III, XII/33 (Ma), Ajar Valley V/77 (TE), Bande Amir VII/78 (Wassey), Ab-i-Estada III/77 (Savey), Kandahar I/1879 (St.J), X/1880, II/1881
- Falco subbuteo Gray

hobby

 \underline{F} . \underline{s} . $\underline{subbuteo}$. Passage migrant (HE) and breeding in northern Afghanistan: Haibak V/37 (RM). Recorded from Kabul VIII-IX/79 (JS), IV, VII, IX, X/65-66 (JN), VIII/33 (Ma) V-X/73-75 (FR), VI/77, Karizmir VI/77 (Crook), Unai Pass IX/33 (Ma), Bamian IV/37, Doshi V/37 (RM), Faizabad (NE) VIII-IX/65 (JN), Kokcha (NE) VIII/48, Nuristan III/48 (KP), Kunduz, Samangan (S) V-X/73-75 (FR).

Falco jugger Gray

lagger falcon

Breeding in southeastern Afghanistan. Recorded from Khairabad, Chazni, Jalalabad, Badakhshan (HE). Ghazni V. VII/49 (KP), II/75, Kabul VI/75 (FR), Kole Hashmat Khan IX/66 (JN), Sardeh (E) IV/78 (TE), E of Jalalabad IV/69 (JV), Kunduz III/75 (FR), Jalalabad VI/37 (RM), VIII/72 (VV), Ghazni-Herat V/77, Shibar Pass V/77 (Thiollay).

Falco cherrug Gray

saker falcon

F. c. cherrug. Passage migrant (Seistan). F. c. milvipes Jerdon. Probably breeding in northwestern Afghanistan (HE). Recorded from Sar-i-Bulak IX/33 (Ma), Sia Koh (C), Farsi (C) VIII/72, Kokcha VIII/72 (VV), Dashte Nawar V/76 (TE), VII/71, Bande Amir VII/72 Sarobi V/72 and 19 records elsewhere between VIII/71-IV/72 (ES), Murghab XII/1884 (CY), Kunduz X/73, Mazar-i-Sharif II/74 (FR).

Falco pelegrinoides Temminck 8.

barbary falcon

F. p. babylonicus Sclater. Resident and breeding, uncommon (HE). Recorded from Kole Hashmat Khan I, IV, V/66 (JN), Kabul Gorge VI/77 (Crook), V/49, Herat, Obeh, Kwaja Chisht (W) VII/49 (KP), Kandahar II, VII/1879 (St.J), Farsi, Jam, Sia Koh (C) VIII/72, Bande Amir (C) VIII/72 (VV), Qala-i-Nau (C) V/77 (RP), Small Pamir VII/71 (CN), throughout Afghanistan V/77 (Thiollay).

ORDER GALLIFORMES

Family Phasianidae

Tetraogallus himalayensis Gray

- Himalayan snowcock T. h. himalayensis. Resident of eastern Afghanistan, Nuristan and
- T. h. incognitus Zarudnyi. Resident of northern Afghanistan from the Hindu Kush west till Bandi-Turkestan (HE). Recorded from Kabul bazar I-III/65-66 (No) I-III, XII/79, I/80 (JS), W. Nuristan VIII-IX/77 (RP), Wama (E) IV-V/48, Bamian VI, X/49 (KP), IV/37 (RM), Salang Pass (C) VII/70 (SM), V/75 (FR), Dhap Darah (C) X/1884 (CY), Paghman (C) VI/33 (Ma), Safed Koh (E) summer 66 (CW), Small Pamir VIII/72 (GN), VIII/71 Big Pamir VII/70 (No), VIII/72, Wakhan VII/71 (GN). Sanglich (E) br VII/37, Munjan Pass (E) VI/37, Safed Sang (NW) IX/39, Burchao Pass

2. Alectoris chukar (Gray)

chukar

The chukar occur in high alpine valleys up to 4 500 m. A. graeca (Meisner) is the European partridge. Four subspecies are distinguished for Afghanistan: A. c. chukar, resident in Nuristan. Reported from Pashki and Stiewe II-IV/48 (KP), western Nuristan VIII-IX/77 (RP); A. c. koroviakovi (Zarudnyi) resident of western and southern Afghanistan; A. c. falki Hartert, resident of the central Hindu Kush and A. c. pallescens (Hume), resident of Badakhshan and the Pamirs. The last three subspecies are reported from Darra-i-Shikari (C) VI/49, Obeh (W) VII/49 (falki), Kokcha Valley (N) VII/48 (KP), Ajar Valley (C) common, III/79 (JS), Kabul Gorge VI/77, Faizabad (N) VI/77 (Crook), Hari Rud XI/1884, Karabagh XI/1884, Bala Murghab XII/1884 (JA), Logar VIII/33 (Ma), Kabul, Ghorband (C), Dar-e-Shikari (C), Bamian IV/37 (RM),

3. Ammoperdix griseogularis (Brandt)

Resident and widely distributed in Afghanistan. Two subspecies see-see partridge A. g. griseogularis, resident of the south and A. g. peraticus Koelz, resident of the northwest. Reported from Qala Birt (S), Tarnak Valley (S) V/49, Obeh, Murghab, Maimanah (N) VI-IX/49, Jalalabad-Sarobi II/48 (KP), Kabul bazar XI-IV/65, 66 (No), Dohab (C) VII/74 (FR), IV/37 (RM), Baghlan VI/74 (FR), Haibak (N) V/37, Barfak (N) V/37, Bamian IV/37 (RM), Lala Maidan (N) br VIII/37 (WK), Asraf (C) VII/33 (Ma), Qala Sarkari (N), Chash maegawan (N) IX/39, Burchao X/39 (WK), Karez-dasht (W) XI/1884 (JA), Kandahar V/1879 (St.J), I/1881 (CS), X/37, Khairabad (NE) VII/37 (WK), Gazechah III/1869 (FM).

Francolinus francolinus (Linnaeus)

black francolin F. f. bogdanovi Zarudnyi. Resident of southern Afghanistan. Recorded from Farah Rud Estuary and Faizabad (SW) III/49 (KP), Seistan X/1884 (JA), Kuhak (SW) VIII/1904 (JC), Argandab, Kandahar 1880 (CS), Lashkargah 73/75 (FR), Girishk X/39 (WK).

Francolinus pondicerianus (Gmelin) 5.

Indian grey partridge

- F. p. mecranensis Zarudnyi and Härms. Southern Afghanistan (HE), reported from Arghestan (SE) VI/72 (ES). Outside Afghanistan it occurs in Persian and Pakistan Baluchistan (Vaurie).
- 6. Coturnix coturnix (Linnaeus)

C. c. coturnix. Summer visitor and passage migrant. Widely common partridge distributed as breeding bird (KP). Recorded from all parts, including Wakhan and Nuristan.

Pucrasia macrolopha (Lesson)

kokloss

P. m. castanea Gould. Nuristan. Griffith collected it in Chagha Sarai III/1840, KP from Gusalek and Wama III-IV/48.

8. Lophophorus impejanus (Latham)

Resident of Nuristan and Safed Koh. The sole collected specimen from Afghanistan was collected in Kurder (E) III/48 (KP). But also recorded from Chaga Sarai 1840 (WG), Mimagi (E) VIII-IX/77 (RP), Kamdesh (E) numerous (Robertson, 1896), Safed Koh numerous 1906 (CW).

9. Phasianus colchicus Linnaeus

P. c. principalis Sclater. Resident in northwestern Afghanistan, Murghab and Hari Rud Basins, P. c. bianchii Buturlin. Resident of Afghan Turkestan, f.e. the Amu Darya Valley. Last century still very

common as reported by Hutton, 1847 for Herat and Yate, 1884 and Aitchison, 1884 for the Hari Rud Valley, Scully, 1884-87 for Bala Murghab but at present very restricted and in high need of protection

ORDER GRUIFORMES

Family Rallidae

Rallus aquaticus Linnaeus 1.

water rail

- R. a. korejewi Zarudnyi. A winter visitor but possibly also breeding (HE). Recorded from Bamian X/49 (KP), Kunduz I/73 (FR), Lundi Rudbar X/1884, Murghab I/1885 (JA), XII/1884 (CY), Kabul IV/37 (RM).
- 2. Porzana porzana (Linnaeus) spotted crake Passage migrant (KP). Recorded from Murghab IV/1885, Chabar Shamba (N) IV/1885 (CY), Danaghori (N) V/37 (RM), Kandahar III/1881 (St.J).
- Porzana parva (Scopoli) little crake Passage migrant. Recorded from Kandahar II/1881 (CS) and Kuhak (Seistan) IX/1904 (JC).
- 4. Porzana pusilla (Pallas) Baillon's crake P. p. pusilla. Possible breeding in north and west Afghanistan, only recorded from Kole Hashmat Khan V/67 (AP), but mainly passage migrant. Recorded from Pashki (E) V/48, Shibar Kotal (C) VI/49, Bamian IX-X/49 (KP), Kole Hashmat Khan V-VI/79 (JS), Kabul IV/37 (RM), Chaharasia IV/33, Rahkol IX/33 (Ma), Kila Wali (N) IV/1885 (CY), Gaz-i-Cha (W) X/1884 (JA), Kandahar II/1881 (CS).
- 5. Crex crex (Linnaeus) corn crake Possibly summer visitor and passage migrant (KP). Recorded from Maruchak (Murghab) V/1885 (CY).
- Gallinula chloropus (Linnaeus) moorhen G. c. indicus Blyth. Breeding bird and possibly winter visitor. Recorded from Kole Hashmat Khan, VI/65 (JN), br. 67 (AP), IV-X/79 (JS), IV/69 (JV), V/77 (Crook), Baqrabad (Seistan) III/49, Chasma-i-Sher VIII/49 (KP), Darunta IV/69 (JV), Kandahar IV/1879 (St.J), Seistan
- 7. Porphyrio porphyrio (Linnaeus) purple coot P. p. seistanicus Zarudnyi. Resident and breeding in Seistan, possible passage migrant in Kabul area. Recorded from Hamun-i-Puzak III/78 (RP) and Jalalabad ? IX/77 (In), Seistan br 1880 (CS), Kabul XI/34 (Fia), Kandahar III/1881 (CS). 8.
- Fulica atra Linnaeus coot F. a. atra. Resident, passage migrant and winter visitor throughout Afghanistan. Breeding in Seistan and Kole Hashmat Khan V/67 (AP). Project records Kole Hashmat Khan 5 000 IV/76 (RP) and III/79 more than 10 000 (JS), p between III-X/79, Sardy Lake 1 000+ I/79, Ab-i-Estada 2 000 IV/79, 3 000 V/79 (JS), Ajar Valley (C) IX/78 (Wassey), Hamun-i-Puzak III/78 (RP), Qargah (Kabul) X/XII/79 (JS).

Family Gruidae

Grus grus (Linnaeus)

common crane

g. lilfordi Sharpe. Passage migrant. Recorded from Qargah 200, 300 III/66, Salang 185 IX/65, Small Pamir 90 VIII/72 (JN), 160 IX/72 (RP), Ab-i-Estada 175 III/77 (Savey), Hamun-i-Puzak 350 III/78 (RP), Hamuni-Saberi 9 III/49, Faizabad (SW) 150, 5, 8, 6 III-IV/49 (KP), Kabul flights of 300 in /73, III/74 and III-IV/75, Kunduz 100 III/75 (FR), Balkh 360 III/72, Mazar-i-Sharif 830 III/72 (ES).

- 2. Grus leucogeranus Pallas Siberian white crane Regular passage migrant, in spring in the bazar in Kabul (HE). Recorded from Ab-i-Estada 76 spring /70 (Koning), 3 winter /70 (No), 2 spring /74 (RP), 56 III/77 (Savey).
- Anthropoides virgo (Linnaeus) demoiselle crane Passage migrant. Recorded from Kabul bazar III-IV/66, Charikar bazar III-IV/71, III/72 (No), Qargah 1 III/76 (JN), III/74 (FR), Ab-i-Estada 2 V/76 (RP), IV/79 (Tischoff), Dashte Nawar V/76 (TE), Kandahar 9, 51 XI/75 (FR), Daronta 2 IV/69 (JV).

Family Otididae

- Tetrax tetrax (Linnaeus) Passage migrant and winter visitor. Kandahar XI/1880 (CS), Murghab little bustard III/1885, Chaman-i-Bed (Badghis) XII/1884 (CY).
- Chlamydotis undulata (Jacquin) C. u. macqueenii (Gray). Passage migrant, winter visitor and probably breeding. Recorded from Hari Rud (Kafir Qala) where in IV/1903 Zarudnyi found it abundant, VII/49, Girishk (S) II/49, Farah-Rud/ Faizabad III/49 (KP), Logar III/34 (Ma), Seistan plentiful in spring, III/05 (JC), Kandahar common 1880 (St.J), Abdul Rahman XI/1881 (CS).
- Otis tarda Linnaeus O. t. tarda. Passage migrant and perhaps breeding in Afghan Turkestan. Recorded from the Kabul bazar II/65 (No), Bala Murghab I/1885 (JA), Danaghori (N) I/35 (Ma), leaving in IV/37 (RM).

ORDER CHARADRIIFORMES

Family Jacanidae

Hydrophasianus chirurgus (Scopoli) Endemic in the Seistan swamps (HE). pheasant-tailed jacana VI-VIII/66 with juveniles (JN). Recorded from the Jalalabad Basin

Family Rostratulidae

Rostratula benghalensis (Linnaeus) l. R. b. benghalensis. Probably breeding in Wakhan, but not recorded since 1847. Jalalabad IV/1840 (WG), Kandahar 1847 (Hutton).

Family Haematopodidae

Haematopus ostralegus Linnaeus

oyster catcher

H. o. longipes Buturlin. Rare passage migrant. Recorded from Farah Rud-Seistan IV/49, Hari Rud (Herat) VII/49 (KP), Ab-i-Estada IV/69 (JV), VIII/67 (Puget), IV/76 (RP), III/77 (Savey), Qargah VIII/72 (ES), V/66, Kabul bazar IV/66 (JN). Gozara (NW) IV/72 (ES).

Family Recurvirostridae

Himantopus himantopus (Linnaeus)

black-winged stilt

H. h. himantopus. Common passage migrant, breeding in Ab-i-Estada (JN) but also elsewhere (Voous). Recorded from the wetlands throughout the country. Project records from Kole Hashmat Khan, IV-V, VII-X/79, Ab-i-Estada IV-V/79 (JS).

Recurvirostra avosetta Linnaeus 2.

R. a. avosetta Linnaeus. Passage migrant, breeding in Ab-i-Estada (JN) and Dashte Nawar (Klockenhoff and Madel). Recorded from the wetlands throughout the country. Project records from Kole Hashmat Khan V-VI, VIII/79, Qargah X/79, Ab-i-Estada V/79 (JS).

Family Bhurinidae

Burhinus oedicnemus (Linnaeus)

stone curlew

B. o. harterti Vaurie. Breeding and probably resident. from Ab-i-Estada v/49 (KP), V/76 (TE), IV, V/79 (JS), Kabul market IV/72 (No), Herat VII/49, Bamian X/49, Hamun-i-Saberi (Seistan) III/49 (KP), Gulran (W) III/1885 (JA), Helmand (Lashkargah) large numbers IX/72 (VV).

Family Glareolidae

Cursorius cursor (Latham)

cream-coloured courser

Breeding bird, passage migrant and winter visitor (HE). Recorded from Herat VII/49 (KP), V/77 (RP), lower Farah Rud (Seistan) III/49 (KP), Hamun-i-Puzak III/78 (RP), Mukur (E) VI/49, Balkh (N) VII/49, Dilaram (S) II/49 (KP), Dashte Nawar br VI/65 (JN), Kandahar-Mukur IV/76 (RP), Ab-i-Estada III/77 (Savey), Lijjikarez (S) III/1896 (FM), Nadali (S) III/03, Khwaja Ahmed (S) IV/03 (JC), Girishk (S) IV/69 (JV), Kandahar III/74, VII/73, Lashkargah III/73, IV/73, Herat IX/72 (FR), Karez, Khorasan VI/1885, Gulran III/1885 (JA).

2. Glareola pratincola (Linnaeus) pratincole Recorded from Ab-i-Estada VI/72, Maruf (SW) VI/72 (ES), Kandahar IV/1879 (St.J), Lashkargah (S) V/74, Mazar-i-Sharif IV/74 (FR).

- Glareola glareola (Linnaeus) 3. collared pratincole G. g. glareola. Paludan only mentions it as once recorded. Also recorded from Ab-i-Estada V/76 (TE), IV/76 (RP).
- 4. Glareola lactea Temminck little pratincole Breeding in the lower Kabul valley VI/68 (HE). Recorded along the Kabul R. IV/77 (TE), Qargah VII/73 (FR), Jalalabad IV/69 (JV), III/1840 (in Whistler).

Family Charadriidae

- 1. Charadrius dubius Scopoli

 C. d. curonicus Gmelin. Summer visitor and breeding in Jalalabad (JN), common passage migrant and winter visitor. Recorded throughout the country. Project reports: Kole Hashmat Khan IV-VI, VII-X/79, Ab-i-Estada IV, V/79 (JS), Hamun-i-Puzak III/78 (RP).
- 2. Charadrius hiaticula Linnaeus ringed plover
 Passage migrant. Recorded from Qargah IX-X/65 (JN), I, IV, V/73,
 Ab-i-Estada IX/72 (FR), I/79 (JS), Dashte Nawar V/77 (TE), Lashkargah
 (S) IV/73, Kunduz V/75, VIII/74, XII/73, Douchy (C) VII/74 (FR),
 Charburjak (SW) XI/71 (ES).
- 3. Charardrius alexandrinus Linnaeus Kentish plover

 C. a. alexandrinus. Breeding in Ab-i-Estada 67 (JN), VI/77 (Crook),
 passage migrant and winter visitor. Recorded from Qargah and Kole
 Hashmat Khan III-V, VIII-X/64-66 (JN), Ab-i-Estada V/49 (KP), III/77
 (Savey), V/76 (TE), I, IV, V/79 (JS), IV/69 (JV), Dashte Nawar V/76, 77
 (TE), Hamun-i-Puzak III/78 (RP), Hamun-i-Saberi III/49 (KP), Kandahar
 II/1881 (CS), Lashkargah, Kandahar, Kabul, Gardez II-V/73-75 (FR).
- Charadrius mongolus Pallas

 C. m. atrifrons Wagler. Passage migrant and breeding in the Small Pamir: Kol-e-Tshaqmaqtin br VII/71 (No), Terghen Qorum (not br) VIII/72 (JN). Other records from Ab-i-Estada, Dashte Nawar V/76, 77 (TE), Lijjikarez III/1896 (FM).
- Passage migrant, breeding in Ab-i-Estada and Dashte Nawar VI/65 (JN), V/69 (JV). Recorded from Ab-i-Estada V/79 (JS), Dashte Nawar V/77 (TE), Bamian-Bande Amir VI/49 (KP), Shorawak (S) II/1896 (Finn), Qargah VIII/71 (ES), Ghazni-Mukur IV/69 (JV).
- 6. Pluvialis dominica (MUller)

 P. d. fulva (Gmelin). Occasional passage migrant (HE). Recorded from Qalat-Ghilzai (S) X/1879 (St.J).
- 7. Pluvialis squatarola (Linnaeus) grey plover Recorded from Ab-i-Estada IV/69 (JV), III/76 (TE).
- 8. Hoplopterus indicus (Boddaert) red-wattled lapwing
 H. i. aigneri Laubman. Breeding bird of southern and western
 Afghanistan (HE). Recorded from Herat (Hari Rud) VII/49, Lashkargah
 V/49, Farah IV/49 (KP), Hamun-i-Puzak III/78 (RP), Duganan III/1896
 (FM), Jalalabad IV/69 (JV), Lashkargah (S) br V/74, IV/73, VII/73 (FR),
 Kandahar II, III (CS).
- 9. Chettusia gregaria (Pallas) sociable lapwing
 Occasional passage migrant (HE). Recorded from Kabul bazar III/71, 72
 (No), Kalayal X/34 (Ma), Ab-i-Estada IV/69 (JV).
- 10. Chattusia leucura (Lichtenstein) white-tailed plover

 Passage migrant and winter visitor, especially of southern Afghanistan,
 perhaps breeding in the west (HE). Recorded from Kole Hashmat Khan
 IX/79 (JS), III/78 (TE), VIII, IX/65 (JN), Qargah VIII/65 (JN), Ab-iEstada III/77 (Savey), Seistan III-IV/49 (KP), Hamun-i-Puzak III/78 (RP),

Kandahar XII/1880, III/1881 (CS), Helmand (Nadali), Shahgul (S) III/03 (NA), Nasratabad (SW), V/05 (CS), Lashkargah (S) II-IV/73-75 (FR), Kham-i-Ab (N) VI/1885 (RY).

11. Vanellus vanellus (Linnaeus)

lapwing

Common passage migrant and winter visitor between X and III-IV. Project records from Qargah XI/79-II/80 (JS), Hamun-i-Puzak III/78 (RP).

Family Scolopacidae

1. <u>Calidris</u> <u>alba</u> (Pallas)

sanderling

Passage migrant. Recorded from Ab-i-Estada V/49 (KP), V/79 (JS), IV/69 (JV), Kole Hashmat Khan V/65 (JN), spring/66 (AP).

2. <u>Calidris minuta</u> (Leisler)

little stint

Very common passage migrant in all Afghan wetlands. Project records from Kole Hashmat Khan IV-VI, VII-X/79, Ab-i-Estada I, IV, V/79 (JS), Hamun-i-Puzak III/78 (RP).

3. <u>Calidris temminckii</u> (Leisler)

Temminck's stint

Passage migrant. Recorded from Kole Hashmat Khan VIII-IX/79, Ab-i-Estada V/79 (JS), V/76 (TE), III/77 (Savey), Qargah V/65, IX/65 (JN), Sarobi IX/72 (ES), Danaghori V/37 (RM).

- 4. Calidris ferruginea (Pontoppidan) curlew sandpiper

 Passage migrant, recorded from Ab-i-Estada V, IX/64-66 (JN), IV/69
 (JV), V/79 (JS), V/78 (TE), Qargah VIII-X/65 (JN), V/72 (ES), Kole
 Hashmat Khan V-VI, VIII-IX/79 (JS), Kabul River Basin VII-VIII/71-72
- 5. <u>Calidris alpina</u> (Linnaeus)

dunlin

C. a. alpina. Scarce passage migrant and winter visitor. Recorded from Qargah X/71 (ES), IX/72, XI/73, I/74, V/75 (FR), Kole Hashmat Khan V/65 (JN), V, VIII/79 (JS), Ab-i-Estada I, V/79 (JS), V/76 (TE), III/77 (Savey), IV/69 (JV), Sardy Lake (E) I/79 (JS), Dashte Nawar V/77 (TE), Hamun-i-Puzak III/78 (RP), Farah IV/69 (JV), Lab-i-Baring XII/18 (NA), Nimroz (SW) XI/71, Herat IV/72 (ES).

6. Philomachus pugnax (Linnaeus)

ruff

Passage migrant. Recorded from Kole Hashmat Khan VII-XI/64-66 (JN), IV/69 (JV), III/78 (TE), V, VIII-IX/79 (JS), Qargah IV/74-75 (FR), Ab-i-Estada 2 000-4 000+ IV, V/79 (JS), 1 300 IV/69 (JV), 2 000+ III/77 (Savey), V/76, Dashte Nawar V/76 (TE), Sardeh Lake III/78 (TE), Hamun-i-Puzak III/78 (RP), Shaharistan (SW) III/03 (NA), Chahar Shamba (N) IV/1885 (CY), Panjah IV/1874 (Bi), Kandahar III/1881 (JC), Danaghori (N) V/37 (RM).

7. Lymnocryptes minimus (Brunnich)

jack snipe

Passage migrant and winter visitor (KP). Recorded from all over northern Afghanistan between X/1884-III/1885 (JA), Kandahar, Seistan 04 (CS, JC, St.J), Kabul IV/37 (RM), I/74 (FR), Kunduz I/73, X-XII/73, 74 (FR).

8. Gallinago gallinago (Linnaeus)

common snipe

G. g. gallinago. Breeding in NE, common passage migrant and winter visitor between X and IV. Project records from Kole Hashmat Khan III, V, VIII, IX/79 (JS).

Chaman = Kole Hashmat Khan Chaman (Pakistan) 30055' 66027' Charburjak 30018' 62004' Charikar 35002' 69013' Chasma-i-Cher 36005' 68035' Chasma Sabz 34050' 61035' Chasma Sabz Pass 34053' 61037' Chiga Sarai = Chaga Sarai Chilik Turkestan Choiltan = Chahiltan Dahnai Ghar 34024' 69002' Danaghori = Dahana 35°54' 68°29' Darband-i-Kul Rekhta 35019' 63027' Darqad region 37025' 69030' Darra-i-Chahidan Bamian Darrah-e-Shaur 37020' 73042' Darra-e-Shikari central Afghanistan Darre Nau E. Afghanistan Darunta = Daronta 34030' 70020' Darwashan = Darweshan 31°02' 64°08' Gusalek = Gursalek 35°00' 70°52' Darwaz N. Badakhshan Dasht-i-Barang 32005' 61035' Dashte Nawar 33044' 67045' Dashte Margo region S. Afghanistan Daulatabad NW 36028' 64052' Daulatabad E 32048' 62039' De Doda 31010' 62005' Delaram = **Dilaram** 32010' 63025' De Kamran 31008' 62005' Deh Tang (Ghorband) 34010' 69029' Doab = Dohab 35015' 68000' Djelallabad = Jalalabad Doshy = Douchy 35938' 68043' Dragon Ajar Valley Dubrai S. Afghanistan Enjil NW Afghanistan Eshkashem = Ishkashim = Ishkamish Faizabad NE 37005' 70040'

Faizabad SW 31028' 61032' Farah 32023' 62007' Farah Rud R SW Afghanistan Farakhulm 34029' 68009' Farkhar NE Afghanistan Farsi 34023' 63006' Firindjal 35000' 68042' Galichah 34058' 61014' Gardez 33037' 69009' Gatai 31005' 66022' Gaz-i-Chah 30025' 63015' Ghazni 33037' 68028' Ghilzai = Gilzai 34°30' 67°35' Ghorband Pass 34010' 69029' Girishk 31050' 64035' Gulam-i-Maidan 35000' 61015' Gulbahar 35010' 69016' Gumanara SW Afghanistan Gungazai NE Afghanistan Hadda Mts SE of Jalalabad Hadjijak Pass 34040' 38015' Haibak 36015' 68003' Hamadan desert SW of Herat Hamun-i-Puzak 31030' 61043' Hamun-i-Saberi 31030' 61015' Hariab R E of Herat Hari Rud R central and NW Afghanistan Harut R central and S. Afghanistan Hazara 32048' 65054' Hazarajat region 33045' 66000' Helmand River S and SW Afghanistan Herat 34020' 62010' Hindu Kush central mountains Imam Sahib 37015' 68045' Ishkashim = Ishkamish 36043' 71034' Ishkatul 36024' 71018' Ishkarzir 36002''61005'

- 9. Gallinago media (Latham) great snipe
 One record from Tolibai (Big Pamir) X/72 (RP) (but perhaps
 G. gallinago).
- 10. Gallinago solitaria Hodgson solitary snipe

 G. s. solitaria. Scarce winter visitor. Recorded from Bamian
 IX, X/49 (KP), Qargah I/80, Kole Hashmat Khan X/79 (JS), Deh Tang,
 Ghorband (C) X/1884 (CY), Chohiltan (C) XII/31 (Ma), Paghman IV/37
 (RM).
- 11. Scolopax rusticola Linnaeus Eurasian woodcock

 Passage migrant (HE). Recorded from Arghandab (SE) between X-III/1880,
 1881 (CS, St.J) Chahiltan XII/31 (Ma), Chahar Shamba (N) II/1885 (CY),
 Kunduz X/73, I/75 (FR).
- 12. Limosa limosa (Linnaeus)

 L. 1. limosa. Passage migrant and winter visitor. Recorded from Kole Hashmat Khan V/66 (JN), V, VII/79 (JS), Qargah V/65 (JN), Ab-i-Estada IV/79 (Tischoff), Ab-i-Estada IV/69 (JV), III/77 (Savey), Hamun-i-Puzak III/78 (RP), Lab-i-Baring (SW) XII/18 (NA), Nasratabad (SW) V/05 (JC), Kandahar II/1881 (CS).
- 13. Limosa lapponica (Linnaeus)

 Occasional passage migrant. Recorded from Ab-i-Estada IX/66 (JN),

 IX/72 (FR), V/76 (TE), Qargah IX/65 (JN).
- 14. Numenius phaeopus (Linnaeus) whilbrel
 Rare passage migrant. Recorded from Dashte Nawar V/77 (TE), Ab-iEstada IV/69 (JV).
- 15. Numenius tenuirostris Vieillot slender-billed whimbrel
 One possible record from Ab-i-Estada V/79 (JS).
- Numenius arquata (Linnaeus)

 N. a. arquata. Passage migrant and winter visitor. Recorded from III/79 (JV), Kole Hashmat Khan VIII-IX/79 (JS), Ab-i-Estada (TE), Hamun-i-Puzak III/78 (RP), Hamun-i-Saberi III/49 (KP), Hamuns common 1905 (JC), Daronta IV/69 (JV).
- 17. Numenius madagascariensis (Linnaeus) Far-eastern curlew Rare passage migrant. One record from Qargah V/75 (FR).
- 18. Tringa erythropus (Pallas)

 Passage migrant. Recorded from Kole Hashmat Khan XII/65, III/66 (JN), VI, VII, IX/79 (JS), Qargah V, IX, X/65 (JN), V/74 (FR), Ab-i-Estada Charburjak (SW) XI/71 (ES).
- 19. Tringa totanus (Linnaeus)

 T. t. eurhinus Oberh. Breeding in Kol-e-Tshaqmaqtin (Small Pamir)

 VII/71 (No) and Dashte Nawar 70 (Klockenhoff and Nadel) and probably
 Ab-i-Estada (JN). Common passage migrant and winter visitor,
 Khan VI-IX/79, Ab-i-Estada IV/79 (JS), Hamm-i-Puzak III/78,
 W. Nuristan VIII-IX/77 (RP), Qargah XI/79 (JS).

- 20. Tringa stagnatilis (Bechstein)

 Rare passage migrant (KP). Recorded from Kole Hashmat Khan IV-V,
 Hamun-i-Puzak III/78 (RP), Danaghori (N), Haibak (N) V/37 (RM),
 Bamian VII/74, Qargah IV. VII/73 (FR).
- 21. Tringa nebularia (Gunnerus)

 Passage migrant and winter visitor. Recorded from Kole Hashmat
 Khan (and Qargah) IV-V, VII-X/64-66 (JN), V/76 (TE), V-IX/79 (JS),
 i-Estada IV/69 (JV), IV-V/79 (JS), Bande Amir VI/77 (Crook), HamunDarunta IV/69 (JV), Kunduz X/73, Kabul I-V/73-75 (FR), Kandahar
 common 1880/81 (CS, St.J), Farakhulm IX/33 (Ma).
- 22. Tringa ochropus Linnaeus green sandpiper

 Common passage migrant and winter visitor. Many records from all the wetlands. Project reports from Kole Hashmat Khan VIII-IX/79 (JS).
- 23. Tringa glareola Linnaeus wood sandpiper

 Common passage migrant. Breeding in Qala Panjan, Ab Gatsh, Karish,
 Neshtkhawar (all Wakhan) VII-VIII/71 (No). Project records from Kole
 Hashmat Khan autumn /79, Ab-i-Estada IV, V/79 (JS).
- Zenus cinereus (Güldenstädt)

 Occasional passage migrant (HE). Recorded from Kole Hashmat Khan (and Qargah) VII-IX/64-66 (JN), VII-IX/79 (JS), Ab-i-Estada V/65, VII-VIII/71, 72 (ES).
- 25. Actitis hypolancos (Linnaeus)

 Breeding in the northern, central and eastern parts of Afghanistan, common passage migrant. Reported from all parts of Afghanistan.

 Project records from Kole Hashmat Khan V-IX/79, Ab-i-Estada IV, V/79 (JS), Hamun-i-Puzak III/78 (RP).
- Arenaria interpres (Linnaeus)

 A. i. interpres. Passage migrant, recorded from Qargah VIII/66 (JN), IX/66 (JN), V/76 (TE), IV/69 (JV), IX/72 (FR), Dashte Nawar V/76 (TE).
- Phalaropus lobatus (Linnaeus)

 Passage migrant. Recorded from Bamian X/49 (KP), Kole Hashmat Khan (Qargah) V, VIII-X/64-66 (JN), V-VI, VIII/79 (JS), IX/72, VI/74 (PR), IX/72 (RP), Adreskan (W) IV/69 (JV).

Family Laridae

- Larus ichthyaetus Pallas great black-headed gull Passage migrant. Recorded from Hamun-i-Saberi, Lower Farah Rud III-III/49 (KP), Seistan common III, IV/04 (NA), (JC), Hamun-i-Puzak Panjah (NE) IV/1874 (Bi).
- Larus ridibundus Linnaeus
 Common passage migrant and winter visitor. Project records from Kabul 111/80, Kole Hashmat Khan III-IX/79, Qargah X-XI/79, Ab-i-Estada IV/79, 1 000+ V/79 (JS), Hamun-i-Puzak III/78 (RP).

3. Larus genei Breme

slender-billed gull

Passage migrant, breeding in Ab-i-Estada, few thousands summer/66 (JN). Recorded from Kole Hashmat Khan and Qargah VIII-IX, III-V/64-66 (JN), VI/74 (FR), Ab-i-Estada V/49 (KP), IV/69 (JV), V/76 (TE), VI/77 (Crook), IV/79, 1000+ V/79 (JS), Dashte Nawar V/77, Sardeh Lake IV/78 (TE), Hamun-i-Saberi II/49 (KP), Kuhak (SW) IV/05 (JC).

4. Larus canus Linnaeus

common gull

- L. c. heinei Homeyer. Passage migrant, one record from Qargah III/73 (FR). It occurs in the south Caspian districts of Iran to Turkmenya
- Larus fuscus Linnaeus L. f. fuscus. Passage migrant. Recorded from Qargah (Kabul) V/66 (JN), Kabul bazar III/71 (No), Ab-i-Estada III/77 (Savey). lesser black-backed gull
- 6. Larus argentatus Pontoppidan

herring gull

L. a. cachinnans Pallas and/or L. a. heuglini Bree. Passage migrant. Recorded from Qargah (Kabul) II/66 (JN), III/74 (FR), Dashte Nawar V/77 (TE), Ab-i-Estada V/49 (KP), IV/69 (JV), Hamun-i-Saberi III/49, Farah V/49 (KP), Jalalabad I/75 (FR).

Rissa tridactyla (Linnaeus)

kittiwake

Recorded from Qargah (IX/65) but probably wrongly identified. It is recorded from the USSR coast, west Mediterranean and inland Turkestan

Family Sternidae

Gelochelidon nilotica (Gmelin) 1.

gull-billed tern

G. n. nilotica. Summer visitor, breeding in Ab-i-Estada 64-66 (JN) and probably also elsewhere, common passage migrant. Recorded throughout the country but not in the east and the Pamirs.

2. Sterna caspia Pallas

Caspian tern

- S. c. tschegrava (Lepechin). Passage migrant, possibly breeding. Recorded from Kole Hashmat Khan V/79 (JS), Hamun-i-Saberi III/49 (KP), Hamun-i-Puzak III/78 (RP), Kuhak (SW) IV/05 (JC), Qargah IV, VI/74,75 (FR), Daronta IV/69 (JV), IV/72 (ES), Sarobi VI/75 (FR), Ab-i-Estada IV/69 (JV).
- 3. Sterna aurantia Gray

river tern

One observation from Qargah VIII/66 (JN).

4. Sterna hirundo Linnaeus

common tern

- S. h. hirundo. Common summer visitor and breeding recorded from Obeh (W) VII/49 (KP) and Kol-e-Tshaqmaqtin (Small Pamir) VII/71 (No). Project records from Kole Hashmat Khan V, VI/79, Ab-i-Estada V/79 (JS),
- 5. Sterna albifrons Pallas

little tern

S. a. albifrons. Summer visitor, breeding in Seistan and probably northwest (NE). Recorded from Kole Hashmat Khan V-VII, VIII/79 (JS), VI/79 (Crook), V/76 (TE), Qargah X/65 V/66 (JN), Ab-i-Estada V/79 (JS), V/76 (TE), IV/69 (JV), Baghlan VI/77 (Crook), Lashkargah V/49, Herat VII/49 (KP), Kham-i-Ab (NE) VI/1886 (CY), Khwaja Ahmed, Nasratabad (SW) V/05 (JC), Girishk (S) IV/69 (JV), Kunduz IV-IX/73-75

Chiidonias hybrida (Pallas) 6.

- whiskered tern C. h. hybrida. Probably passage migrant. Recorded from Kabul V/37 (RM), Kole Hashmat Khan V/76 (TE), V-VI, VIII-IX/64-66 (JN), V-IX/79 (JS), Ab-i-Estada V/79 (JS), IV/69 (JV), V/76, Dashte Nawar V/76 (TE), Daronta IV/69, Dilaram IV/69 (JV), Sarobi, Kabul, Jalalabad
- Chlidonias nigra (Linnaeus) 7. black tern C. n. nigra. Unsubstantiated records from Ab-i-Estada VI/77 (Crook)
- 8. Chlidonias leucoptera (Temminck) white-winged black tern Recorded from Ab-i-Estada VI/77 (Crook), V/79 (JS), Qargah V/74 (FR).

ORDER PTEROCLIDIFORMES

Family Pteroclididae

- Pterocles coronatus Lichtenstein cornetted sandgrouse P. c. atratus Hastert. Breeding bird of the steppes of southern Afghanistan. Recorded from Girishk (S) V, VI/49, Shin Dand (W) VII/49 (KP), Bassoolah (SE) (Gr) and Shorawak (SE) 1896 (Ma), Lashkargah V/74 (FR). In the Helmand desert it was common but nowhere numerous, 1880 (St.J).
- 2. Pterocles senegallus (Linnaeus) spotted sandgrouse Recorded from southern Afghanistan (KP), Juwain-Farah III/78 (RP), Lijjikarez III/1896 (Ma) and Seistan XII/18 (NA). In XI/71 at Nimroz 450 were assembled at water pockets in the dry Helmand River bed (ES).
- Pterocles orientalis (Linnaeus) black-bellied sandgrouse P. o. arenarius (Pallas). Uncommon passage migrant and winter visitor of the steppes throughout Afghanistan. Breeding in Dashte Nawar VI/65 (JN), northwest III/1885 (JA), Kandahar IV/1879 (St.J). Project records from Ab-i-Estada I, IV, V/79 (JS). Not recorded from Nuristan and the Pamirs.
- Pterocles alchata (Linnaeus) 4. pin-tailed sandgrouse P. a. caudacutus (Gmelin). Resident of northwestern and southern Afghanistan. Recorded from Farah (SW) VI/49 (KP), Kham-i-Ab (N) VI/1885, Andkhui (NW) X/1885 (CY), VII/49 (KP), Kandahar XII/1880

ORDER COLUMBIFORMES

Family Columbidae

Columba livia Gmelin

rock dove

C. 1. neglecta Hume. Resident of eastern Afghanistan, recorded from the Pech Valley (E) III-IV/48 (KP). C. 1. gaddi Zarudnyi and Loudon. Resident of the whole of Afghanistan, except the eastern part. Recorded throughout the country, especially the river gorges (VV), common in the south and formerly especially attended in pigeon houses. Project records from Ab-i-Estada IV/79 (JS).

2. Columba rupestris Pallas

- eastern rock dove
- C. r. turkestanica Buturlin. Breeding bird of northeastern Afghanistan above 1 00° m elevation (RP). Recorded from Gusalek (E) III/48, Kokcha Valley (N) Warduj Valley (N) VII/49 (KP), Wakham Valley (NE) VII-VIII/71 (No).
- 3. Columba leuconota Vigors

snow pigeon

- C. 1. leuconota. Resident and breeding bird of eastern Afghanistan above 2 000 m elevation (RP). Recorded from the Parun Valley (E) V-VI/48, Weran (NE) VI/48 (KP), Qala-i-Pandja (NE) IX/65 (JN). The only record outside the area is from the Shibar Pass (C) VI/72 (ES).
- 4. Columba oenas Linnaeus

stock dove

- C. o. yarkandensis Buturlin. Rare winter visitor to northern Afghanistan (HE). Recorded from Karabagh (NW) XI/1884 (JA) and probably Herat V/77 (RP).
- 5. Columba eversmanni Bonaparte eastern stock dove Summer visitor of the Hari Rud, Paropamusus and the northern and northeastern Hindu Kush, migrating through southern Afghanistan (HE). Recorded from Danaghori (N), Kunduz (N), Haibak (N) br V/37 (RM), Tirpul, Toman Aghi (NW) IV/1885 (JA), Maimanah VI/1885 (CY), Qala Nau br, Murghab, Maimanah, Mazar-i-Sharif VII-VIII/49, Herat, Obeh (W) br VII/49 (KP), Kuhak (SW) IV/05 (JC), Ajar V/77, Dashte Nawar V/77 (TE), and Ab-i-Estada III/77 (Savey).
- Columba palumbus Linnaeus 6.

wood pigeon

- C. p. casiotis (Bonaparte). Breeding bird of northwestern, central and eastern Afghanistan. Recorded from Obeh (W) VII/49 (KP), Murghab br VIII/72, Daulatabad VIII/72 (VV), Kamdesh (E) IV/69 (JV), Pech-Parum Valley (E) IV, V/48, Kachari (NE), Kwaja Chist (N) br VII/48 (KP), Hariab (E) br IV/1879 (WR), Kandahar III/1879 (St.J), Ashrai
- Streptopelia decaocta (Frivaldszky) 7. Indian ring pigeon S. d. decaocta. Common summer visitor to towns, villages and oases, not recorded from the forested east and Badakhshan but common in the lower Kabul Valley.
- Streptopelia tranquebarica (Hermann) 8. red turtle dove S. t. tranquebarica. Recorded from Jalalabad br V/37 (RM), common in Pakistan (Vaurie).
- 9. Streptopelia turtur (Linnaeus)

turtle dove

- S. t. arenicola (Hartert). Common summer visitor to northwestern and northern Afghanistan, arrives in May and immediately starts breeding /37 (RM). Big migrating flocks were seen in Qala-i-Panja IX/72 (JN). Migrates also through southwestern Afghanistan (HE).
- 10. Streptopelia orientalis (Latham)

rufous turtle dove

S. o. meena (Sykes). Summer visitor to eastern and central Afghanistan and passage migrant. Recorded from Pech-Parun Valley (E) VIII-IX/77 (RP), V-VIII/48 Bamian IX-X/49, Gardez V/49, Farah VI/49 (RP), Jalalabad IX/77 (In), Abakhan Valley (NE) IX/71 (RP), Tolibai (NE) VIII/72, Darwaz VII/72 (JN), Atin Jalao (NE) VIII/72 (VV), Haibak (N), Doshi (N), Ghorband (C) V/37 (RM), Safed Koh arriving in large numbers V/1879 (WR).

- 11. Streptopelia senegalensis (Linnaeus) palm dove

 S. s. cambayensis (Gmelin). Resident or summer visitor to southern and eastern Afghanistan. S. s. ermanni (Bonaparte). Resident or summer visitor to northern Afghanistan (HE). Common and breeding in the Afghan towns and villages. In Kabul from I-XII/79 I-IV/80 (JS).
- 12. Streptopelia chinensis (Scopoli) spotted dove

 S. c. suratensis Gmelin. Eastern Afghanistan, presumably breeding bird of Kandahar and Kabul regions (HE). Recorded from Jalalabad V/37 (RM) and Kandahar summer 1880 (Hutton).

ORDER PSITTACIFORMES

Family Psittacidae

- 1. Psittacula krameri (Scopoli) rose ringed parakeet
 P. k. borealis (Neumann). Introduced in Jalalabad and Kabul (HE).
 Many records also from Kunduz (FR).
- 2. Psittacula himalayana (Lesson) slaty-headed parakeet

 P. h. himalayana. Summer visitor of Nuristan and Safed Koh (HE).

 Recorded from Pech Valley (E) V, VII/48 (KP), Kunar Valley IV/69 (JV),

 Pashat IV/1840 (WG).
- 3. Psittacula eupatria (Linnaeus)

 P. e. nipalensis (Hodgson). Recorded from Jalalabad X/69, probably not introduced (AP). Nearest records from Peshawar in Pakistan (Ali et al).

ORDER CUCULIFORMES

Family Cuculidae

1. Cuculus canorus Linnaeus

cuckoo

C. c. subtelephonus Zarudnyi. Summer visitor and breeding bird. Recorded throughout Afghanistan. First records in April and May, last in September. In the northern part the great reed warbler is recorded as common host V/37 (RM).

ORDER STRIGIFORMES

Family Strigidae

- 1. Otus brucei (Hume) striated scops owl One single record from Panjah (Wakham, NE) IV/1874 (Bi).
- 2. Otus scops (Linnaeus)

 O. s. pulchellus (Pallas). Perhaps breeding in Kandahar (KP).

 Recorded from Kandahar IV/1879 (St.J), IV/1881 (CS).
- 3. <u>Bubo bubo (Linnaeus)</u>
 <u>B. b. nikolskii Zarudnyi.</u> Resident of the whole of Afghanistan except for Badakhshan (HE). Recorded from Girishk (S) V/49, Obeh (W) VII/49, Haibak (N) VIII/49 (KP), Kabul bird market V/79 (JS), Ajar Valley

- V/77 (TE), Danaghori (N) V/37 (RM), Kandahar II/1879 (St.J).

 B. b. auspicabilis Dementiev (or B. b. hemachalana Hume). Resident of Badakshan (HE). Recorded from Tolibai (Big Pamir) VIII/72 (JN), Kol-e-Tshaqmaqtin (Small Pamir) VIII/72 (RP).
- 4. Glaucidium brodiei (Burton) collared pygmy owlet

 G. b. brodiei. Recorded from the forests of Payopari Char (E) X/69

 (AP), Kamdesh (E) IV/69 (JV).
- Athene noctua (Scopoli)

 A. n. bactriana Blyth. Widespread but uncommon resident. Recorded throughout the country except Nuristan. Breeding recorded in Kunduz V/37 (RM), Pamir (Langar) VII/71 (No).
- 6. Strix aluco Linnaeus tawny owl
 S. a. biddulphi (Scully). Resident of northeastern and eastern
 Afghanistan, south to Safed Koh (HE). Recorded from Hariab Valley
 (E) V-VI/1879 (WR), and Gusalek (E) II/48 (KP).
- 7. Asio otus (Linnaeus)

 A. o. otus. Passage migrant and winter visitor and probably breeding (KP), Darre Naw (E) VIII, IX/77 (RP), Kunduz winter/73/74, I/75, Sarobi XI/74, Lashkargah II/75 (FR), Kandahar ruins common 1880 (St.J), Maimanah III/1885 (CY), Murghab II/1885, Maruchak II/1885 (JA).
- 8. Asio flammeus (Pontoppidan)

 A. f. flammeus. Passage migrant and winter visitor. Recorded from (KP), Kunduz XII/73, Mazar-i-Sharif II/74 (FR), Murghab XII/1884 (JA), Baghrami (C) III, X/33, Logar X/33 (Ma).

ORDER CAPRIMULGIFORMES

Family Caprimulgidae

- 1. Caprimulgus mahrattensis Sykes Sykes' nightjar
 Southern Afghanistan, probably summer visitor (HE). Recorded from
 Lijjikarez (SE) 1896 (FM), Helmand br IV-IX/05 (JC).
- 2. Caprimulgus asiaticus Latham Indian nightjar
 C. a. gurgaoni Koelz. Straggler, recorded from NW Afghanistan:
 Tirpul IV/1885, Khusan IV/1885 (JA).
- 3. Caprimulgus europaeus Latham

 European nightjar

 C. e. unwini Hume. Summer visitor and passage migrant. Recorded from Lashkargah V/49 (KP), Kandahar V/1879 (St.J), Panjao (C) VI/49, Bamian IX/49, Balkh (N) VII/49 (KP), Bande Amir (C) VIII/70 (SM), Qalai Pandja (NE) VIII/71 (No), Kabul X/73 (FR), VI/34 (Ma), Khusan (NW) V/1885 (JA), Kila Wali V/1885 (CY), Logar VI/33 (Ma).
- Caprimulgus aegypticus Lichtenstein

 C. a. arenicolor Severtzov. Summer visitor of western and southwestern Afghanistan (HE). Recorded from Faizabad (SW) III-IK/49 (KP), and USSR part of Amu Darya (N) (Vaurie).

ORDER APODIFORMES

Family Apodidae

1. Apus apus (Linnaeus)

swift

A. a. pekinensis (Swinhoe). Common summer visitor and passage migrant throughout the country. First records February in Kandahar, March in Kabul. Project records from Kabul X/79, Kole Hashmat Khan V-IX/79, Ab-i-Estada IV, V/79 (JS).

2. Apus melba (Linnaeus)

Alpine swift

A. m. tuneti Tschusi. Common summer visitor and passage migrant. Recorded throughout the country except Nuristan. First spring records Darunta, hundreds II/72 (ES), last record October, Kabul. Project records Kabul X/79, Kole Hashmat Khan IX/79 (JS), Darre Nau (E) IX/77, Herat br V/77 (RP).

3. Apus affinis (Gray)

little swift

A. a. galilejensis (Antinori). Summer visitor and passage migrant. Probably winter visitor of the Lower Kabul Valley. Breeding recorded from Kandahar 1879 (St.J), Chagai Sarai (E) IV/69 (JV). Elsewhere recorded from Jalalabad IX/77 (In), I/75 (FR), IV/69 (JV), Sarobi III/78 (TE), Ab-i-Estada IV/69, Kalat, Kandahar IV/69 (JV), Murghab (NW) VII/49, Farah, Faizabad (S) IV, V/49 (KP).

ORDER CORACIIFORMES

Family Alcedinidae

1. Halcyon smyrnensis (Linnaeus)

white-breasted kingfisher

H. s. smyrnensis. Resident in lower Kabul and Kunar Valley (NE). Recorded from Kunar III/1840 (WG), Darunta II/48 (KP), II/77 (TE), II/79 (JS), Jalalabad IX/77 (In), XII/73, XI/74, I/75 (FR), IV/69 (JV), Rost (E), Chagha Sarai IV/66 (AP).

2. Alcedo atthis (Linnaeus)

common kingfisher

A. a. atthis. Summer visitor, resident of southern and eastern Afghanistan. Breeding in Sarobi, Kunduz, Baghlan /73-75 (FR), common in Kandahar and Seistan (JC, CS, St.J). Elsewhere recorded from Kabul VIII-XII/79 (JS), VI/49 (KP), Kole Hashmat Khan VIII-IX/79, Ajar Valley (C) III/79 (JS), Bamian IX-X/49 (KP), Bande Amir VIII/70 (SM), IX/33, Logar VI/33 (Ma), Chagha Sarai (E), Kamdesh (E), Darunta (E) IV/69 (JV), Kunduz V/37 (RM), Shin Dand (W), Farah, Herat, Faizabad (SW) IV-VII/49 (KP), VI/77 (Crook), Mazar-i-Sharif VIII/49 (KP), Pech Valley VIII/77 (RP).

3. Ceryle rudis Linnaeus

pied kingfisher

- C. r. leucomelanura Reichenbach. Breeding bird and probably resident of eastern Afghanistan. Recorded from Pech River (E) II-III/48, Darunta (E) II/48 (RP), IV/69 (JV), XI/74, I/75 (FR), II/77 (TE), II/79 (JS), Sarobi XI/73 (FR), Jalalabad II/48 (KP), IX/77 (In), Laghman V/37 (WK), Hindu Kush VIII/72 (VV).
- 4. Ceryle lugubris (Temminck) great pied kingfisher

 C. 1. guttulata Stejneger. Recorded along the Kamdesh River (E) in

 (Pakistan) (Vaurie). It occurs in Kashmir

Family Meropidae

- Merops orientalis Latham little green bee-eater
 M. o. beludschicus Neumann. Recorded from Jalalabad V/ (HE), V/37,
 Torkhan (E) V/37 (RM).
- 2. Merops superciliosus Linnaeus blue cheeked bee-eater

 M. s. persicus Pallas. Common summer visitor to the plains and passage migrant. Breeding recorded from Sarobi 71, 72 (ES), Danaghori (N), Kunduz and Jalalabad V/37, large breeding colonies (RM), Seistan V/05 (JC). Many records between end of April and September (Seistan October). Project record from Herat surroundings V/77 (RP).
- 3. Merops apiaster Linnaeus European bee-eater
 Summer visitor and numerous passage migrant. It breeds in considerable
 numbers at moderate elevations. Common between middle of April and
 August. Project records from Kole Hashmat Khan, Qargah VIII/79,
 Ab-i-Estada V/79 (JS), Herat, Qala-i-Nau V/77 (RP).

Family Coraciidae

1. Coracias garrulus Linnaeus

roller

- C. g. semenowi Loudon and Tschusi. Very common summer visitor and passage migrant throughout the country between March and September.
- Coracias benghalensis (Linnaeus)
 Linnaeus
 Indian roller
 benghalensis
 Recorded as breeding bird from Afghan Siestan
 Further records from Hariab Valley V/1879 (WR), Jalalabad
 IV/69 (JV).

Family Upupidae

1. Upupa epops Linnaeus

hoopoe

<u>U. e. epops.</u> Common summer visitor and passage migrant, a few are winter visitors (KP). In Kabul present throughout the year (JS), Ab-i-Estada IV/79 (JS).

ORDER PICIFORMES

Family Picidae

1. Jynx torquilla Linnaeus

wryneck

- J. t. torquilla. Passage migrant. Recorded from Kabul XI, XII/74 (FR), VIII-XI/79 (JS), Kole Hashmat Khan IV/65 (JN), IV/79 (JS), Pashki (E) V/48, Kunar Valley (E) III/48 (KP), Ab-i-Estada IV/79 (JS), Bande Amir VIII/70 (SM), Qala-i-Pandja (NE) IX/65 (JN), Kunduz IV/74, Mazar-i-Sharif IV/74, Jalalabad I/75, Kandahar II/74 (FR), VIII/1879 (St.J), IV/1881 (CS), Girishk, Adreshkan (W) IV, V/69 (JV), Logar IV/34 (Ma), Bamian very common IV/37 (RM).
- 2. Picumus innominatus Burton speckled piculet
 P. i. innominatus: Recorded from Nuristan: Begame Valley XII/63,
 Gusalek VII/65 (JN). Outside Afghanistan it occurs from NW Punjab
 (Pakistan) eastwards (Vaurie).

3. Picus squamatus Vigors

P. s. squamatus.

X/1884 (JA).

- scaly-bellied green woodpecker Resident of eastern Afghanistan from Safed Koh and Kabul to Nuristan (HE). Recorded from Gusalek II/48 and Wama (E) IV/48 (KP), Pech River (E) br V, VI/66 (LT), Kabul VIII/72 (VV), Gardez VI/35 (Blanchard), Logar XII/33 (Ma). P. s. flavirostris (Menzbier). Resident of southern and western Afghanistan. Recorded
- Dendrocopus himalayensis (Jardine and Selby) 4. Himalayan pied woodpecker D. h. albescens (Baker). Summer visitor of eastern Afghanistan. Recorded from the Pech-Parun Valley (E) III-VI/48 (KP), Pech River (E) br V, VI/66 (LT), Barikot (E), Kamdesh (E) IV/69 (JV), Hariab Valley (E) abundant br V-VI/1879 (WR).

from Herat and Obeh (W) VII/49 (KP), Padda Sultan, De Kamran (W)

- 5. Dendrocopus auriceps (Vigors) brown-footed pied woodpecker D. a. auriceps. Resident in Nuristan. Recorded from Pashat, Chagha Sarai III/1847 (WG), Pech Parun Valley IV-V/48 (KP), Pech River (E) br V, VI/66 (LT), Kamdesh, Kamu (E) IV/69 (JV), Pashar, Bharawal III/1840 (WG).
- 6. Dendrocopus leucopterus (Salvadori) white-winged spotted woodpecker Resident in northern Afghanistan from the Paropamisus and northern Hindu Kush till Badakhshan. Recorded from Faizabad (NE), Jurm and Supian (NE) VII/48, Haibak (N) VIII/49 (KP), N. Salang VII/70 (SM), Darqad X/78 (Wassey), Salang I/73 (FR), Doshi (N), Haibak (N), Baghlan V/37 (RM).

ORDER PASSERIFORMES

Family Alaudidae

1. Ammomanes cincturus (Gould)

bar-tailed desert lark

1

A. c. zarudnyi Hartert. Breeding bird of southern Afghanistan. Recorded from Nimroz (W) I/72, Helmand, Kandahar II-III/72 (ES), Lab-i-Baring XII/15 (NA).

2. Ammomanes deserti (Lichtenstein)

desert lark

Three subspecies are distinguished A. d. iranica Zarudnyi. Resident of the southwest and west. Recorded from Shin Dand VII/49, Seistan II-III/49 (KP), Farah VIII-IX/37, Kandahar X/37 (WK). A. d. orientalis Zarudnyi and Loudon. Resident of the north, recorded from Murghab (N) and Maimanah VII/49 (KP), Faizabad VI/77 (P&C), Doab, Barfak, Haibak (N) IV, V/37 (RM) Tala (N), XII/37, Chigzar (N) IX/37, Kamard (N) VIII/37, Turkestan IX/37 (WK). A. d. phoenicuroides (Blyth). Resident of the east and south, recorded from Kole Hashmat Khan IX, X/65 (JN), Kabul VIII/72 (VV), IV/37 (RM), Turukpul (E of Kabul) br V/37, Nazhil (E) V/37 (WK), Lataband III/78 (TE), Jalalabad II/48 (KP), IV/69, Chaga Sarai IV/69 (JV), Kandahar III-V/1879 (St.J), I/1881, Malkarez IV/1881, Dubrai IV/1881 (CS).

Jalalabad 34°27' 70°25'

Jalalabad Valley lower Kabul
R Valley

Jam 34023' 64022' Jawzjan Northern Province Juwein = Juwien 31042' 61036' Jurm 36°50' 70°46' Kabul 34°30' 69°10' Kabul Gorge E. of Kabul Kabul R E. Afghanistan Kachari Kokcha Valley Kafiristan = Nuristan Kajakay = Kayakal 32°20' 65°05' Kalat = Qalat 32008' 66053' Kalat-i-Ghilzai = Kalat Kalat us Seraj 34°40' 70°18' Kambao 35005' 61026' Kamiab = Kham-i-Ab Kamdesh 35°25' 71°22' Kamu Valley 35°23' 71°23' Kandahar 31037' 65047' Kang 31°07' 61°52' Karabagh = Qarabagh C 340501 1690081 Karabagh = Qarabagh NW 34057' 61046' Karawal Khana 35043' 63014' Karez Dasht 33017' 61030' Karizmir 34°30' 69°10' Kargah = Qargah 34030' 69002' Kathagan Pass - 37069' Kathakhak 34°50' 68°00' Khairabad NE Afghanistan Kham-i-Ab 37032' 65042' Khanabad 36042' 69008' Khash R central and S Afghanistan

Khawak Chapchi Pass 35040' 69051'

Khelat-i-Ghilzai Badghis

Khost 330221 700011

Khossik Pass SE Afghanistan

Khusan = Kohsan 34°38' 61°11'
Khusah = Kohsan 34°38' 61°11'
Khushk 32°51' 71°05'
Khwaja Ahmed S. Afghanistan
Khwaja Ali 30°25' 63°15'
Khwaja Chist = Chisht-i-Sharif
34°19' 63°45'

Kila Wali = Qala Vali 35047' 63044' Kin 32°15' 61°35' Koaja Palounda Badghis Koh-i-Baba central mountain spur Koh-i-Duzdan = Kouh Dazd 32013' 63029' Kotal-i-Dalez Wakhan Kohsan 34038' 61011' Kokcha 37000' 69030' Kokeran S. Afghanistan Kole Hashmat Khan 34030' 69010' Kol-e-Tshaqmaqtin 37015' 74012' Kuhak SW Afghanistan Kunar R 34045' 70055' Kunduz 36047' 68051' Kushk 34053' 62030' Kustaki 35°29' 70°56' Lab-i-Baring SW Afghanistan Laghman Province 34035' 70008' Lal 34030' 66015' Lalander 34°25' 69°10' Lashkargah = Lashkari Bazar 31°30' 64°20' Lataband Pass 34030' 69034' Lijjikarez Samuli SE Afghanistan Lokhi SW Afghanistan Logar Valley

Lundi 30°12' 62°45'

Madan Sorb Cave 35°00' 68°42'

Magnaul 36°05' 70°58'

Maidan 32°28' 66°45'

Maidan Tal 34°27' 68°50'

Maimanah 35°55' 64°46'

3. Alaemon alaudipes (Desfontaines)

- Hoopoe lark, bifaciated lark
- A. a. doriae (Salvadori). Resident of southern Afghanistan. Recorded from Tarnak Valley (S), Pirzada (W of Kandahar), IV/49, Dilaram II-VI/49, Hamun-i-Saberi II-IV/47 (KP), Farah-Juain IV/76, Hamun-i-Puzak III/78 (RP), Kandahar IX/1879 (St.J), 73-75 (FR), Girishk X/37 (WK), IV/69 (JV), Kuhak (SW) IX/04 (JC), Zagin (W) XI/1881 (JA), Kang (SW), Salyan (SW) XI/37 (WK).
- 4. Melanocorypha calandra (Linnaeus)
- calandra lark
- M. c. psammochroa Hartert. Resident of northern Afghanistan and passage migrant. Recorded from Khanabad br V/37 (RM), Maimanah, Murghab, Mazar-i-Sharif, Pul-i-Khumri(N) VIII/72 (VV), Mazar-i-Sharif winter 73-75, large flocks (FR), Andkhui (NW) XI/37, Taligan (NE) VIII/37 (WK). Often in bird markets (Kabul, Charikar).
- 5. Melanocorypha bimaculata (Ménétries) eastern calandra lark

 M. b. torquata Blyth. Breeding bird in northern and northeastern
 Afghanistan, common passage migrant (HE). Breeding recorded from
 Tashkotal (W) V/37 (RM). Many records throughout the country. Often
 kept as pet and sold in bird markets, especially during the migratory
 season.
- 6. <u>Calandrella cinerea</u> (Gmelin)

short-toed lark

- C. c. longipennis (Eversmann). Summer visitor and breeding bird of the north. Probably br in Danaghori V/37 (RM). Very common passage migrant, sometimes in enormous numbers (Kandahar, 1879, 1880 (St.J), Kole Hashmat Khan VIII-X/79 (JS), Ab-i-Estada IV/79 (JS), 5000+ III/77 (Savey)). Recorded throughout the country, except the high central, eastern and northeastern mountains. Often kept in cages, especially during the migratory seasons. The birds are sometimes intermediates between C. c. longipennis and C. c. artemisiana Banikovski.
- 7. Calandrella acutirostris Hume

 C. a. acutirostris. Breeding bird of the Alpine steppes of western, central and northeastern Afghanistan, east to Gardez. Here rather common in small numbers between April and August, some remained in the winter 71/72 (ES). Breeding recorded in the Shibar Pass and Unai Pass V/37 (RM). Also recorded from Sust-e-Bala (Pamir) VIII/72 (JN), Panjah (NE) IV/1874 (Bi) and Hariab Valley V/1879 (WR).
- 8. Calandrella rufescens (Vieillot)

 C. r. persica (Sharpe). Breeding bird of Seistan (HE). C. r. heinei (Homeyer). Passage migrant and winter visitor (HE). Recorded from Mukur (E), br (E) V/49 (KP), X/37 (WK), Ab-i-Estada V/76 (TE), II/77 (Savey), V/79 (JS), Dashte Nawar V/76 (TE), Sardeh Lake IV/78 (TE), Nimroz (SW) common winter visitor 71/72 (ES) Douchy (C), Sarobi, Jalalabad I/73-75, Kandahar VII/73-75, Ghazni XI/73-75 (FR) Qala-i-Nau (N) VIII/72 (VV), Kang (SW), Farah (SW) (heinei) XI/37 (WK).
- 9. Calandrella raytall (Blyth) Indian sand lark
 C. r. ?adamsi (Hume). One record from Jalalabad IX/77 (In).
- 10. Galerida cristata (Linnaeus) crested lark

 G. c. magna Hume. Resident. Widely distributed throughout the country, but not in Nuristan and the Pamirs. It avoids the high mountains and in the winter it is found at lower altitudes, common in the south and west. Many records, project records Ab-i-Estada I, IV, V/79 (JS).

11. Lullula arborea (Linnaeus)

wood lark

L. a. pallida Zarudnyi. One single record from Nimroz (SW) I/72 (ES). Vaurie gives as easternmost record SW Iran.

12. Alauda gulgula Franklin

lesser skylark

A. g. inconspicua Severtzov. Resident of Afghanistan except for Badakhshan. Reported throughout the country, common between April and September. Breeding reported from Sanglich (NE) VII/37 (WK), Kabul, Bamian, Danaghori IV-V/37 (RM). A. g. lhamarum Meinertshagen. Resident of Badakhshan. Recorded from Tilli (NE) VII/48, Qala-i-Pandja (NE) IX/65 (JN), Wakhan common till VII/71 (No).

13. Alauda arvensis Linnaeus

skylark

A. a. dulcivox Hume. Summer visitor to northern Afghanistan, passage migrant. Recorded from Panjah (NE) IV/1874 (Bi), Gandachesma (Badakhshan) VII/37 (WK), Ardewan Kotal (W) VII/49 (KP), Ab-i-Estada I/79 (JS), V/76 (TE), Hamun-i-Puzak III/78 (RP), Herat br V/77 (RP), Samangan, Mazar-i-Sharif, Ghazni and Lashkargah migrating II, X/73-75, Salang V/75 (FR), Bande Amir VIII/72, Dalautabad (N) VIII/72 (VV), Kandahar II/1881 (CS), Aftao (W) II, XII/1884 (JA), Seistan X-XI/37, Andkhui (NW) X/39, Balkh XI-XII/37, Jalalabad XII/37 (WK).

14. Eremophila alpestris (Linnaeus)

shore lark

E. a. albigula (Bonaparte). Resident of the central, eastern and northeastern mountains and breeding above 2 500 m elevation. Recorded throughout this area, common in Wakhan and Pamirs (No), (JN).

Eremophila bilopha (Temminck)

Temminck's horned lark

The record from the Kabul Gorge VI/77 (P&C) probably refers to \underline{E} . alpestris. The easternmost record in Vaurie is from the Syrian deserts.

Family Hirundinidae

1. Riparia paludicola (Vieillot)

brown-throated sand

R. p. chinensis (Gray). Recorded breeding from Kabul V/37 (WK), and also considered common in VI (HE), hundreds E of Sarobi VIII, IX/71 (ES).

2. Ripara riparia (Linnaeus)

sand martin

R. r. riparia. Uncommon breeding bird of Afghanistan (HE). Recorded from Barfak (N) grasslands V/37 (RM). R. r. diluta (Sharpe and Wyatt). Passage migrant. Recorded throughout the country. Project records Ab-i-Estada IV-V/79 (JS).

3. Ptyonoprogne fuligula (Lichtenstein)

pale crag martin, rock martin

(Syn. <u>Hirundo obsoleta</u>) <u>P. f. pallida</u> (Hume). Perhaps breeding in the south <u>Lijjikarez</u> and Samuli III/1896 (FM). Crook recorded it from the Salang Pass VI/77.

4. Ptyonoprogne rupestris (Scopoli)

crag martin

Summer visitor and breeding bird to the mountains of north, central and east Afghanistan. Recorded throughout that area, between April (first 30 March, Kabul and 15/II/74 Sarobi (FR)) and beginning November.

5. <u>Hirundo rustica</u> Linnaeus

swallow

H. r. rustica. Common summer visitor and breeding bird, passage migrant between the end of February (Kabul: March) and September. Recorded throughout the country.

6. <u>Hirundo smithii</u> Leach

wire-tailed swallow

- H. s. filifera Stephens. Localized but common summer visitor (HE). Recorded throughout the country except in the extreme west and south and the higher mountains from April till October (November).
- 7. Hirundo daurica Linnaeus red-rumped swallow
 H. d. rufula Temminck. Summer visitor to northern and eastern
 Afghanistan (KP). Recorded throughout this area, between the end of
 March and October.
- 8. <u>Hirundo fluvicola</u> Blyth Indian cliff swallow Recorded from Kabul V/37 (WK) and Sarobi XII/71 (ES).
- 9. Delichon urbica (Linnaeus) house martin

 D. u. urbica. Summer visitor to the mountainous areas of central and northeastern Afghanistan. In the Pamirs nesting places till 4 000 m elevation. Also passage migrant. Many records between February and September.

Family Motacillidae

1. Anthus novaeseelandiae (Gmelin)

Richard's pipit

- A. n. waitei Whistler. Indian subspecies, probably a summer visitor to southern Afghanistan (KP). Recorded from Gatai (S) IV/1881 (CS) and Kandahar V/1881 (St.J).
- 2. Anthus sylvanus (Hodgson) upland pipit
 Probably resident of eastern Afghanistan. One record from AndarabBanu VII/33 (Ma).
- 3. Anthus campestris (Linnaeus)

tawny pipit

- A. C. campestris. Passage migrant in winter in south Afghanistan (Vaurie), recorded from Bamian IX-X/49 (KP), IV/37 Kabul IV/37 (RM). A. C. griseus Nicoll. Scarce breeding bird of northern Afghanistan and passage migrant. Recorded from Tashkotal (N) and Haibak (N) br IV, V/37 (RM). Various records throughout the country as passage
- 4. Anthus similis Jerson

long-billed rock pipit

- A. s. decaptus Meinertzhagen. Summer visitor to northeastern
 Afghanistan, passage migrant. Recorded from Faizabad (SW) III/49 (KP),
 Safed Koh br VI/1879 (WR), Kandahar VI/1879 (St.J). A. s. jerdoni
 Finsch. Jalalabad V/37 (RM).
- Anthus hodgsoni Richmond olive-backed tree pipit
 Passage migrant, three subspecies breed in USSR and winter in India.
 Recorded from Qala-i-Pandja, common IX/65 (No), common VII/72 (JN).
- 6. Anthus trivialis (Linnaeus) tree pipit

 A. t. trivialis and A. t. haringtoni Witherby. Common spring passage migrant. Recorded from Wama and Pashki (E) IV-V/48 (KP), Paghman IX/33 (Ma), IV/37, Kabul IV/37 (RM), IV/34 (Ma), IV, XII/74 (FR),

I, IV/80 (JS), Kabul valley IV/77, N of Kunduz IV/78 (TE). Bande Amir VIII/70 (SM), Bamian IV/37 (RM), IX-X/49, Panjao (C) VI/49 (KP), Kunar (E) II/1840 (WG), Khyber, Jalalabad, Chagha Sarai, Kamu (E), Kamdesh (E), Sarobi (E) IV/69 (JV), Panjah (NE) IV/1874 (Bi), Kandahar IV/1881 (CS).

7. Anthus pratensis (Linnaeus)

meadow pipit

A. p. pratensis. Passage migrant of Afghanistan Turkestan (N) (HE).

- 8. Anthus cervinus (Pallas)

 Passage migrant. Recorded from Jalalabad IX/77 (In), Ab-i-Estada
 V/79 (JS), Hamun-i-Puzak III/78 (RP), Panjah (NE) IV/1874 (Bi), Kabul
 IV/37 (RM).
- 9. Anthus roseatus Blyth
 Summer visitor of the east (Safed Koh) (Vaurie), one record from Maruchak (NW) III/1885 (CY).
- 10. Anthus spinoletta (Linnaeus) water pipit

 A. s. japonicus Temminck and Schlegel, one record Paghman IV/37 (RM),

 A. s. coutelii Audouin. Passage migrant, winter visitor. Recorded from Ab-i-Estada-Mukur III/77 (Sa), Hamun-i-Puzak III/78 (RP), Dashte Nawar III/77, Kabul Valley II/77 (TE), Kandahar common XII/1880-iii/1881 (CS), (St.J), Murghab I/1885 (JA) Chahar shambai (E) III/1885 (CY), Chaharasia (C) III/34 (Ma), common between X-III/73-75 in more humid areas (FR).
- Motacilla flava Linnaeus

 M. f. melanogrisea (Homeyer). Uncommon summer visitor and breeding bird of northern and western Afghanistan. Recorded from Farah Rud Estuary (SW) III/49, Herat VII/49, Chasma-i-Cher (W) III/49 (KP). Five subspecies are known as passage migrant, M. f. flava, M. f. beema (Sykes) only in April, M. f. thumbergi Billberg, large flocks, M. f. feldegg Michahelles breed in northern Afghanistan (Vaurie) and M. f. leucocephala (Przewalski). Recorded throughout the country. Project records Kole Hashmat Khan IV-V, VII-IX/79 Ab-i-Estada IV, V/79 (JS), Hamun-i-Puzak III/78, Western Nuristan VIII-IX/77 (RP).
- Motacilla citreola Pallas

 M. c. calcarata Hodgson. Breeding bird of central, northwestern and northeastern Afghanistan, passage migrant. Recorded from Wakhan Big and Small Pamir br VII-VIII/71 (No), 72 (JN), Bamian br IV/37, Ghorband Valley br IV/37, Doab (C), Shibar Pass IV/37 (RM), Kandahar few br, common or passage 1879/80 (St.J), Miyan Deh, Tilli (Badakhshan) VII/48, Darra-i-Shikari (C), Panjao (C), VI/49, Herat VII/49, Bamian IX/49 (KP). M. c. citreola and M. c. werae (Buturlin) are common passage migrants. Project records Kole Hashmat Khan IV-VI, VII/79, Ab-i-Estada IV/79 (JS) Western Muristan VII-IX/79 (RP).
- Motacilla cinerea Tunstall grey wagtail

 M. c. cinerea. Summer visitor, passage migrant and winter visitor.

 Breeding recorded from Paghman, very common 64-66 (JN), IV/37 Ghorband IV/37 (RM), Hariab Valley (E) 1879 (WR). Many records as passage migrant and few as winter visitor. Project records Paghman VII/79, Qargah XII/79 (JS).

14. Motacilla alba Linnaeus

white wagtail

M. a. personata Gould. Common summer visitor and breeding in north (RM) and Safed Koh (WR), passage migrant and winter visitor.

M. a. dukhunensis Sykes and possibly M. a. baicalensis Swinhoe are passage migrants and winter visitors from October on. Project records from Kole Hashmat Khan IV-V, IX/79, Ajar Valley III/79, Ab-i-Estada IV, V/79, Kabul winter /79, IV/80 (JS), Western Nuristan VIII-IX/77, Big Pamir 72 (KP).

Family Campephagidee

Pericrocotus ethologus (Bangs and Phillips) flame-coloured minivet

P. e. favillaceus Bangs and Phillips. Scarce but widely distributed summer visitor of eastern Afghanistan (KP), and the lower Kabul Valley (Vaurie). Recorded from Pech-Parun Valley (E) IV-V/48 (KP), Kamdesh (E) IV/69 (JV), Otipore (E) IV/1840 (WG), Nimla (E) II/77 (TE).

Family Pycnonotidae

1. Pycnonotus leucogenys (Gray)

white-cheeked bulbul

- P. 1. leucogenys. Resident of the east (HE). Recorded from Laghman $\overline{V}/37$ (WK), Jalalabad V/37 (RM), 73/75 (FR), IX/77 (In), IV/69, Chagha Sarai IV/69 (JV), Sarobi III/78 (TE), Wama (E) IV/48 (KP).
- P. 1. leucotis (Gould). Resident of the south. Recorded from Farah (S) II/76 (RP), II-VI/49 (KP), X/37 (WK) Kandahar common XII/1880, II/1881 (CS), II/49 (KP), X/37 (WK), Herat VII/49 (KP).
- 2. Hypsipetes madagascariensis (Statius Müller) black bulbul (Syn. Microscelis psaroides). H. m. psaroides (Vigors). Probably summer visitor to Kunar Valley (E) and Safed Koh (HE). Recorded from Chagai Sarai (E) III/1880 (WG), IV/69 (JV), Kamu, Kamdesh (E) VI/66, IX/69 (AP).

Family Bombycillidae

- 1. Bombycilla garrulus (Linnaeus) waxwing

 B. g. centralasiae Pokakov. Probably a passage migrant (HE).
- 2. Hypocolius ampelinus Bonaparte grey hypocolius

 Passage migrant. Recorded from Kandahar X/37 (WK), V/74 (FR),

 Faizabad (SW) IV/49, Lashkargah (S) V/49 (KP), V/74, VII/73 (FR).

Family Cinclidae

1. <u>Cinclus cinclus</u> (Linnaeus)

dipper

C. c. leucogaster Bonaparte. Western, central and eastern Afghanistan (HE). Recorded from Obeh (W) VII/49, Hazarajat VI/49, Bamian IX-XI/49 (KP), IV/37, Ghorband IV/37 (RM), Paghman V-VII/79 (JS), Salang (C) VII/70 (SM), VII/74, Doab VII/74 (FR), Hadjijak Pass VI/77 (P&C), Jam V/77 (RP), Big and Small Pamirs VII-VIII/71 (No), IX/75, VII/72 (JN), Sufak (C) IX/39, Mak (C), Gharchi (C), Burchao Pass (C)

2. Cinclus pallasii Temminck

brown dipper

C. p. tenuirostris Bonaparte. Resident in central and northeastern Afghanistan. Recorded from Munjar Pass VII/37, Paghman VI/37, Baltistan Valley VII/36, Kargasi Pass (NE), Sanglech Valley (NE) VIII/37 (WK), VII/48, Kokcha Valley (NE) VII/49 (KP), Big Pamir VII-VIII/71 (No), Kamdesh (E) br IV/69 (JV), Pech-Parun Valley II-VI/48 (KP), Salang (C) VII/70 (SM), 73-75 (FR), Panchir (C) V/73 (FR), Iskar (N), DehTang (C) Ghorband (C) X/1884 (CY), IV/37, Shibar (C) juv. V/37, Paghman IV/37 (RM), Khawak Chapchi VII/33 (Ma).

Family Troglodytidae

1. Troglodytes troglodytes (Linnaeus)

wren

T. t. neglectus Brooks. Resident of Nuristan, collected from Gusalek (E) II, III, V/48 (KP). T. t. magrathi (Whitehead). Resident of Safed Koh (HE). Recorded from Sirotai (E) VI/37 (WK).

T. t. tianschanicus Sharpe. Resident of northern and northeastern Afghanistan. Recorded from Ajar Valley III/79 (JS), Salang X-III/73-75 (FR), br V/70 (SM), Wakhan VIII/71 (No), Paghman IV/37 (RM), Chahiltan XII/33, Logar I/35 (Ma), Kabul, Kunduz X-III/73-75 (FR), Chahar Shambai (N) I, II/1885 (SY), Maimanah, Safed Sang, Sufak (N) Burchao (C), Balkh, Pul-i-Khumri (N) IX-XI/39 (WK).

Family Prunellidae

1. Prunella strophiata (Blyth)

rufous-breasted accentor

 $\frac{P.\ s.\ jerdoni}{Recorded\ from}$ (Brooks). Breeding bird of eastern Afghanistan. Recorded from Sirotai Pass VI/37 (WK), Stieive (E) VI/48 (KP), Safed Koh (E) (Vaurie).

2. <u>Prunella fulvescens</u> (Severtzov)

brown accentor

- P. f. fulvescens. Breeding bird and probably resident of the northern Hindu Kush and Badakhshan (HE). Recorded from Weran Valley (NE) VII/48 (KP), Big and Small Pamir common VIII/65, VII-VIII/72 (JN), VII-VIII/71 (No), Western Nuristan VII-IX/77 (RP), Paghman, Ghorband IV/37 (RM), Salang (C) IX/66 (JN), VII/70 (SM) I-VII/73-75 (FR), Bamian IX, X/49 (KP), Sanglech (NE) br, Munjan Pass VI, VII/37, Sabz, Safed Sang, Burchao (C) VIII-X/39 (WK).
- 3. Prunella atrogularis (Brandt) black-throated accentor

 P. a. atrogularis. Winter visitor of southern Afghanistan (HE).

 Recorded from Farah Rud Estuary (SW) II/49 (KP), Maimanah, Andkhui,
 Balkh XI/37, Girishk X/37, Pul-i-Khumri XI/39, Balkh X/39 (WK).

 P. a. huttoni (Horsfield and Moore). Winter visitor of northern and
 eastern Afghanistan (HE). Recorded from Kabul XI-XII/79, I-III/80

 (JS), Kole Hashmat Khan II/65 (JN), Gusalek (E) III/48 (KP), Big Pamir
 IX/65 (JN), Kandahar X/1879 (St.J), Balkh XI/37, Jalalabad XII/37,
 Pol-i-Khumri XI/39 (WK). (FR) gives records throughout Afghanistan
 between XII and III/73-75 and Salang V/75.
- 4. Prunella himalayana (Blyth) Himalayan accentor
 Breeding bird of the northern Hindu Kush and Badakhshan (HE).
 Recorded from Salang IX/66 (JN), VII/70 (SM), Sarobi III/78 (TE),
 Darwaz (NE) VII/72, Wakhan, Big Pamir VIII/72, Big, Small Pamir
 VII-VIII/71 (No), Murjan (NE) VII/37 (WK).

5. Prunella collaris (Scopoli)

Alpine accentor

P. c. rufilata (Severtzov). Breeding bird of northeastern Afghanistam and Safed Koh, passage migrant of northwestern Afghanistan (HE). Recorded from N Salang VII/70 (SM), VI/74, S Salang I/75, Kabul Gorge XI-III/73-75 (FR), II/77 (TE), Murghab XII/1884 (JA), Bulola (E) XII/37 (WK), Pech-Parun Valley (E), II, VI/40 (KP).

Family Turdidae

1. Cercotrichas galactotes (Temminck)

grey-backed warbler

C. g. familiaris (Menetries). Summer visitor and passage migrant. Recorded from Faizabad (SW), Farah IV/49, Obeh (W), Qala Nau (NW), Murghab (NW) VII/49, Bamian IX/49 (KP), Adreskan (W) V/69 (JV), Kunduz V/73, 75, Lashkargah V/74 (FR), Danaghori (N) V/37 (RM), Tirpul IV/1885, Kambao V/1885 (JA), Kilawali IV/1885, Darbund-i-Kil-Rekhta (N) V/1885. (CY), Kandahar 1880 (St.J).

2. Erithacus rubecula (Linnaeus)

robir

E.r. tataricus Grote or E.r. hyrcanus Blanford. Easternmost distribution is Iran (Vaurie). The only record for Afghanistan is from Maywand (SE) III/72 (ES).

3. Lucinia megarhynchos Brehm

nightingale

L. m. africana Fischer and Reichenow. Probably summer visitor of northwestern Afghanistan (HE). L. m. hafizi Severtzov. Summer visitor of northern Afghanistan and passage migrant (KP). Recorded from Faizabad (NE) VII/48, Panjao (C) VI/49 (KP), Bande Amir VII/70 (SM), Baharak (NE) IV/77 (P&C), Obeh (W) VII/49 (KP), V/77, Jam V/77 (RP), Danaghori (N) br V/37 (RM).

4. Lucinia svecica (Linnaeus)

bluethroat

L. s. abbotti (Richmond). Breeding bird of Badakhshan and central Afghanistan. Recorded from Kabul II/33 (Ma), Tilli (NE) VII/48, Panjao (C) br VI/49 (KP), Dashte Nawar br VI/65 (JN), VI/77 (TE). Four more subspecies L. s. svecica, L. s. pallidogularis Zarudnyi, L. s. kobdensis (Tugarinov) and L. s. saturatior (Sushkin) are recorded as common passage migrants (HE). Project records: Kabul IX-XI/79, III/80, Ab-i-Estada IV/79 (JS), western Nuristan VIII-IX/77 (RP).

- 5. Lucinia pectoralis (Gould)

 L. P. ballioni (Severtzov). Breeding in northeastern Afghanistan

 (HE). One record from Stiewe (E) VI/48 (KP).
- 6. <u>Lucinia brunnea Modgson</u> Indian blue robin

 L. b. brunnea. Recorded from Stiewe and Pashki (E) IV-VI/48 (KP)

 and Bande Amir VIII/70 (SM).
- 7. Tarsiger cyanurus (Pallas)

 T. c. rufilatus (Hodgson). Resident and breeding in eastern

 Afghanistan, up to 3 000 m. Recorded from Safed Koh (HE).

 Tarsiger hyperythrus (Blyth)

 rufous-breasted bush robin

Presumably recorded from Bande Amir VIII/70 (SM). Vaurie gives the distribution Himalayas from Garhwal (Pakistan) eastwards.

- 8. Irania gutturalis (Guerin-Meneville) white-throated robin

 Summer visitor, breeding bird of northern mountainous Afghanistam, south to Kabul and passage migrant (HE). Recorded from Kabul V/34 (Ma), Bamian IV/37 (RM), Obeh (W) VII/49 (KP), Qala-i-Nau (W) V/77, Jam V/77 (RP).
- 9. Phoenicurus erythronotus (Eversmann) Eversmann's redstart

 P. e. erythronotus. Common passage migrant throughout Afghanistan between the end of October and March/April. Breeding bird of the Pamirs and Wakhan. Project records Kabul XI/79, III/80, Qargah XII/79 Ajar Valley III/79 (JS). Summer records from breeding area: Qala-i-Panja (NE) IV/1874 (Bi), IX/65 (JN).
- 10. Phoenicurus caeruleocephalus Vigors blue-headed redstart
 Breeding bird of eastern Afghanistan and passage migrant. Recorded
 from the Pech-Parun Valley (E) br III-VI/48 (KP), Kamdesh (E) IV/69
 (JV), Byankhel (E) br V/1879 (WR), Kabul X/79 (JS), N Salang
 VIII/70 (SM), VII/74 br, III, V/75 (FR), Darunta II/77 (TE).
- 11. Phoenicurus ochruros (Gmelin)

 P. o. phoenicuroides (Horsfield and Moore). Common breeding bird of mountainous Afghanistan, passage migrant and winter visitor throughout Afghanistan. Project records from Kabul IX-XI/79 (JS).
- 12. Phoenicurus phoenicurus (Linnaeus) redstart

 P. p. phoenicurus and P. p. samamisicus (Hablizl). Passage migrants.

 Recorded from Kabul IV/80 (JS), Yawkalang (C) V/72 (ES), Ardreshkan
 V/69 (JV), Mazar-i-Sharif IV/74 (FR).
- 13. Phoenicurus frontalis (Vigors) blue fronted redstart
 Only recorded from Pashki (E) V/48 (KP). Main distribution from
 Kashmir (Pakistan) eastwards (Vaurie).
- Phoenicurus erythrogaster (Güldenstädt) Güldenstädt's redstart

 P. e. grandis (Gould). Passage migrant (KP), breed in NE Afghanistan and the Pamirs (Vaurie). Recorded from Panjah (NE) common IV/1874 (Stolizka), Pamir common IX/65, VIII/72 (JN), Salang (C) VIII/65 (JN), VI/77 (P&C), br VII/70 (SM), Bamian X/49 (KP), and as winter visitor between XI-III/73-74 from Salang, Douchy (C), Kunduz, Sarobi, Kabul, Samangan (N) and Mazar-i-Sharif (FR).
- Rhyacornis fuliginosus (Vigors) plumbeous redstart

 R. f. fuliginosus. Resident of northeastern and eastern mountainous
 Afghanistan. Recorded from Sarobi (E) vi/37 (WK), Pech-Parun (E) br
 II-VI/48 (KP), Chagha Sarai, Bandikot, Kamu br IV/69 (JV), Sarobi
 X-III/73-75, Pansjir IV/75, V/73, 75, Doab VII/74 (FR), Wakhan VII/71
 (No), Keshem (NE) VI/77 (P&C), N Salang VII/70 (SM).
- 16. <u>Saxicola macrorhyncha</u> (Stolizka) Stolizka's whinchat Very local resident of southern Afghanistan, recorded from Kandahar IV/1881 and Dubai IV/1881 (CS).
- 17. Saxicola rubetra (Linnaeus) whinchat

 Recorded from Kaymard, Sayghan (C) IV-V/75 (ES), S Salang VI/73 (FR).

 Vaurie gives as easternmost record Iran.

18. Saxicola torquata (Linnaeus)

stonechat

S. t. maura (Pallas). Breeding bird of Afghanistan except in the southwestern part, common passage migrant and winter visitor. Recorded throughout Afghanistan. Project records from Kabul IX/79, III-IV/80, Ab-i-Estada IV/79 (JS).

19. Saxicola caprata (Linnaeus)

pied bush chat

S. c. rossorum (Hartert). Common summer visitor, breeding on lower altitudes, also passage migrant and winter visitor (KP). Recorded throughout the country.

20. Oenanthe isabellina (Temminck)

isabelline wheatear

Common resident and in the north summer visitor, breeding in desert areas. Also a passage migrant. Recorded throughout the country. Project records Ajar Valley III/79, Ab-i-Estada I, IV, V/79 (JS).

21. Oenanthe oenanthe (Linnaeus)

common wheatear

<u>O. o. oenanthe.</u> Summer visitor and passage migrant. Recorded throughout the country and common in less dry habitats.

22. Oenanthe pleschanka (Lépéchin)

pied wheatear

O. p. pleschanka. Summer visitor and passage migrant. Breeding in western, central and eastern Afghanistan. Recorded throughout the country between February and October. Project records from Kabul III/80 (JS), Qala-i-Nau deserts V/77, Jam V/77 (RP).

23. Oenanthe deserti (Temminck)

desert chat

- 0. d. atrogularis (Blyth). Passage migrant and winter visitor.
- O. d. oreophila (Oberholser). Summer visitor, passage migrant and winter visitor, breeding at high altitudes in the central Hindu Kush and the Pamirs (JN). Recorded throughout the country. Project records from Hamun-i-Puzak III/78 (RP).
- 24. Oenanthe finschii (Heuglin)

Finsch's wheatear

- O. f. barnesi (Oates), Summer visitor to northern Afghanistan. Widespread resident of central and winter visitor of southern Afghanistan. Also passage migrant. Breeding recorded from Kabul VII/65 (JN), Atin Jalao (NE), most easterly known breeding place VIII/72 (VV), Haibak (N), Doshi (C), Doab (C), Kunduz (N) IV, V/37 (RM). The only Nuristan record is from VIII-IX/77 (RP).
- 25. Oensathe xanthoprymma (Hemprich and Ehrenberg) red-tailed chat

 O. x. chrysopygia (DeFilippi). Common breeding bird of the central and eastern mountains, south to Safed Koh; passage migrant. Winter visitor to the low altitudes such as Sarobi, Kandahar and Jalalabad (KR).
- 26. Oenanthe picata (Blyth)

eastern pied wheatear

Common summer visitor and passage migrant, winter visitor to the lower Kabul Valley. Three colour phases are distinguished which overlap each other in distribution area, and with many intermediates:

O. p. capistrata more concentrated in central and northern Afghanistan;

O. p. opistoleuca more concentrated in eastern and northeastern Afghanistan, predominant in Kabul, not found in the winter;

O. p. picata least common of the forms, more in the south, not found in the Salang (KP, JV, FR). Project records from Kole Hashmat Khan VIII-IX/79 (JS), Hamun-i-Puzak III/78, Qala-i-Nau, Herat V/77 (RP).

Oenanthe monacha (Temminck)

hooded wheatear

Probably southern Afghanistan (HE). An unsubstantiated record from Herat-Farsi V/77 (RP).

27. Oenanthe alboniger (Hume)

Hume's chat

Probably resident of southern Afghanistan (HE). Recorded from Kole Hashmat Khan IX/79 (JS), Kabul Valley IV/77 (TE), Herat-Qala-i-Nau V/77, Chakcharan-Jam V/77 (RP), Qala-i-Bost IX/72 (VV), Delaram, Farah, Adreshkan IV, V/69 (JV) Kandahar, Sarobi, Jalalabad XI-III/73-75 (FR), Zagu (W) XI/1885 (JA), Kandahar, Delaram X/37, Farah XI/37, Takhtipul XI/39 (WK).

28. Chaimarrornis leucocephalus (Vigors) white-capped redstart

Resident of central and eastern Afghanistan, south to Safed Koh (HE). Recorded from Pech-Parun Valley (E) II-VI/48 (KP), Chagai Sarai (E) III/1840 (WC), br IV/69, Barikot (E) IV/69, Sarobi IV/69 (JV), X-III/73-75 (FR), II/48 (KP), Kabul Gorge II/77 (TE), Salang (C), Pansjir br 73-75 (FR), Ajar (C) III/79 (JS), Bamian IX/49 (KP), Salang N VII/70, Khawak Pass VII/33 (Ma), Sarghaz (NE) br VII-VIII/71 (No).

- 29. <u>Saxicoloides fulicata</u> (Linnaeus) Indian robin

 <u>S. f. cambaiensis</u> (Latham). Only once recorded by Griffith, 1840.
- 30. Monticola saxatilis (Linnaeus)

 Summer visitor and passage migrant (KP). Common breeding bird of the Hindu Kush (JN). Recorded from Koh-i-Baba, Firozkoh, Guldara (C) VIII/72 (VV), Adreskhan (W) V/69 (JV), Shibar (C) br V/37 (RM), VII/33 (Ma), VI/49, Panjao (C) VI/49 (KP), Paghman br, Unai Pass (C) br, Ghorband br IV, V/37 (RM), Sar-i-Chasma (C) VI/34 (Ma), Pansjir, Bamian br 73-75, Samangan (N) VII/73 (FR), NW Afghanistan III, IV/1885 (JA), Bande Amir VIII/70 (SM), Salang VI/77 (P&C), Dashte Nawar (E) V/76, 77 (TE), Qala-i-Nau (W) V/77, Jam V/77 (RP), Darwaz VII/72 (JN).
- 31. Monticola cinclorhyncha (Vigors) blue-headed rock thrush
 Summer visitor of mountainous eastern Afghanistan, including Safed
 Koh. Recorded from the Pech-Parun Valley (E) IV-V/72 (KP), Hariab
 Valley (E) br V, VI/1879 (WR), Parsi (C) VIII/72 (VV).
- Monticola solitarius Linnaeus

 M. s. longirostris (Blyth). Summer visitor to western and central Afghanistan, and in the last century recorded as winter visitor. Recorded breeding from Kabul, Ghorband, Akrobat, Doab, Haibak, Paghman IV, V/37 (RM) and from Kabul V/49, Obeh (W) VII/49, Shibar (C) IX/49, Bala Murghab (NW) VII/49, Waibak (N) VIII/49 (KP).

 M. s. pandoo Sykes. Summer visitor to Nuristan. Recorded as scarce breeding bird of the Pech-Parun Valley br IV-VI/48 (KP). Other records throughout the country except the Pamirs.
- 33. Myiophoneus caeruleus (Scopoli) blue whistling thrush

 M. c. turcestanicus Zarudnyi. Resident of central, eastern and northeastern Afghanistan in rocky places and always near water. Winter
 visitor to the south. Recorded throughout the country.
- 34. Turdus unicolor Tickel Tickel's thrush

 Recorded from Chitral (Pakistan) (HE). One record from Sarobi (E)
 II/72 (ES).

Maiwand SE Afghanistan Mak central Afghanistan Malik Dorkhand Mts S. Afghanistan Mangalpur 34006' 69051' Marmakhel 34°20' 70°00' Maruchaq 35048' 63008' Maruf SW Afghanistan Matun = Khost Mazar-i-Sharif 36043' 67005' Meterlam Laghman Miana Bam 35010' 63033' Minla E. Afghanistan Mukur 32052' 67042' Murghab R NW Afghanistan Murghul 34035' 70048' Musa Qala 32023' 69049' Nadali Helmand basin Naghlu 34042' 69045' Nangahar province E. Afghanistan Nao = Nau 35054' 70044' Nasratabad SW border Neshtkaur 37000' 73018' Nijrao 34040' 69045' Nimroz Seistan Nuristan area N. of Jalalabad Obeh 34023' 63006' Oruzgan central Afghanistan Otipore E. Afghanistan Oxus R = Amu Darya R Padda Sultan 31005' 62005' Paghman 34033' 68055' Paktia province 33030' 70000' Pamir E. part of Badakhshan Panjsjir R 35025' 69040' Panja = Qala Panjah 36059' 72040' Panjah R 37000' 72040' Panjao 34021' 67000' Parendah R tributary Panjsjir R

Par-i-Kham Wakhan Paropamisus NW mountain spur Parun 35028' 70055' Parwan Pashat = Pushut 34042' 71003' Pashki 35025' 70056' Payopari Char E. Nuristan Pech R tributary Kunar R Peiwar Kotal 33058' 69051' Pirzada 31037' 65005' Pol-i-Shamri = Puli Khumri 35056' 68044' Puistagoli 35025' 70055' Puzak Gish Mt Doshakh Qachqar Cave Jalalabad Valley near Murgul Qala Bist = Qala Bost 31°28' 64°21' Qala-i-Jadid = Spin Baldak

Qala-i-Jadid = Spin Baldak
Qala Kang 31007' 61052'
Qala-i-Nau 34058' 63004'
Qala Safed Quanai 34030' 69000'
Qala-i-Shahi Nangahar
Qala Vali 35047' 63044'
Qal-i-Charak central Afghanistan
Qalat = Kalat 32007' 66054'
Qarabagh = Karabagh
Qoul Kharideh 75 km W of Kabul
Rah Kol 34050' 68000'
Registan S. Afghanistan
Robate Khusk R = Khush Rud R
32040' 62016'

Rost Safed Koh

Rudbar 30008' 62029'

Sabz Pass 34053' 61037'

Sabzak Pass 35030' 67035'

Safed Koh 34015' 70000'

Safed Sang 35033' 65017'

Sahib Zada S. of Kandahar

Salang Pass 36024' 69008'

35. Turdus merula Linnaeus

blackbird

T. m. intermedius (Richmond). Rare summer visitor to northwestern Afghanistan (KP) and winter visitor. Recorded from Obeh (W), Kwaja Chist (W) VII/49 (KP), Ajar Valley (C) III/79 (JS), X/78 (Wassey), Khusan (W), Murghab (NW) XI, XII/1885 (JA), Chahar Shamba (N) II/1885 (CY), Farakar br VII/37, Maimanah, Murghab, Balkh XI-XII/37, Kandahar VII/37 (WK) XII/1880 (CS), XI-IV/73-75, Kabul, Kunduz, Samangan, Mazar-i-Sharif XI-IV/73-75 (FR).

36. Turdus rubrocanus Hodgson

grey-headed thrush

T. r. rubrocanus. Breeding bird of Safed Koh (Vaurie). Recorded from Pech River (E) VI/66 (LT).

37. Turdus ruficollis Pallas

dark-throated thrush

T. r. ruficollis. Rare passage migrant and winter visitor.

T. r. atrogularis Jarocki. Passage migrant and winter visitor.

Common in some years. Recorded from Kabul XI/33 (Ma), IV/37 (RM), X/79, II-III/80 (JS), Kabul Valley II/77 (TE), Qargah III/66 (JN), Qala-i-Pandja (NE) IX/65 (JN), Wama (E) IV/48, Bamian IX/49 (KP), Ajar Valley III/79 (JS), Chahiltan XII/33, IX/34 (Ma), Chahar Shamba I, II, IV/1885 (CY), Kushk-Rud XI/1884, Murghab (NW), I/1885 (JA), between XI, IV/73-75 winter visitor in Salang, Kabul, Sarobi, Kunduz, Samangan, Ghazni, Lashkargah, Mazar-i-Sharif (FR).

38. Turdus pilaris Linnaeus

field fare

Winter visitor. One record from Kabul II/74 (ES).

39. Turdus iliacus Linnaeus

redwing

T. i. iliacus. Winter visitor. Recorded from Salang(N) XII/72, I/73 and Lashkargah (S) XI/73 (FR).

40. Turdus viscivorus Linnaeus

mistle thrush

T. v. bonapartei Cabanis. Scarce breeding bird, also winter visitor. Recorded from Kamdesh (E) IV/69 (JV), Pech-Parun Valley (E) br IV-VI/48, Obeh (W), Sauzak Valley (W) VII/49 (KP), Bande Amir VIII/70 (SM), Khinjan Pass (C) VII/33 (Ma), Safed Koh br V/1879 (WR), Salang V/75, VII/74, XI-II/73-75, Kandahar, Lashkargah, Douchy (C) Mazar-i-Sharif XI-II/73-75 (FR).

41. Enicurus scouleri Vigors

little forktail

Scarce resident of northeastern Afghanistan. Recorded from Pech-Parun Valley (Nuristan) II-VII/48, Kokcha Valley: Kachari (NE) VII/48 (KP), N. Salang (C) VII/70 (SM), XII/73, Panchir (C) V/75 (FR).

42. Enicurus maculatus Vigors

spotted forktail

E. m. maculatus. Scarce resident of Nuristan: Pech-Parun Valley (E) III-V/48 (KP).

Family Sylviidae

1. Cettia cetti (Temminck)

Cetti's warbler

C. c. albiventris Severtzov. Summer visitor, breeding north of the Hindu Kush (Vaurie). Recorded from Panjao (C) VI/49 and Obeh (C) VII/49 to the south (KP). Also from Ajar Valley VI/76 (TE), Darunta (E) IV/69 (JV), Kandahar I/1881 (CS) and Shahar Shamba II/1885 (CY).

- 2. <u>Cisticola juncides</u> (Rafinesque) fan-tailed warbler

 C. j. cursitans (Franklin). Recorded breeding in eastern Afghanistan (Vaurie).
- 3. Prinia gracilis (Lichtenstein) graceful prinia

 P. g. lepida Blyth. Resident, breed south of the Hindu Kush. Records from Farah II, IV, V/49 (KP), Jalalabad IX/77 (In), Samuli (SE)

 III/1896 (FM), Kuhak (SW) X/04 (JC), Lashkargah III, V, XI/74, II/75, Kandahar XI/74 (FR).
- 4. Prinia inornata Sykes plain-coloured prinia

 P. i.?terricolor (Hume). A single record from a marsh near Jalalabad

 TX/77 (In). Occurs outside Afghanistan in the plains of Pakistan,

 west to Peshawar.
- 5. Prinia criniger (Hodgson) brown hill warbler
 P. c. striatula (Hume). Recorded from Gusalek (E) III/48 (KP),
 Sarobi II/74, Jalalabad XI/74, I, VI/75, Doab (N) VII/74 (FR).
- 6. Scotocerca inquieta (Cretzschar) scrub warbler

 S. i. platyura Severtzov. Breeds in Paropamisus and Afghan Turkestan, recorded from Kabul IV/37, Doab (N) IV/37, Haibak br V/37 (RM), Obeh (C) VII/49 (KP), Samangan II/74, Mazar-i-Sharif XI/74 (FR).

 S. i. striata (Brooks) breeds in southern Afghanistan (Vaurie).
- 7. Locustella certhiola (Pallas)

 L. c. rubescens Blyth and L. c. centralasiae Sushkin. Occasional visitors (HE). Both subspecies only recorded from Bamian IV/37 (RM).
- 8. Locustella naevia (Boddaert) grasshopper warbler

 L. n. straminea Seebohm. Passage migrant throughout, breeds in the north from Herat to Kunduz, winters in the south (HE). Recorded from Bamian IV/37 (RM), IX/49 (KP), VIII/70 (SM) and only one record of L. n. mongolica Sushkin from Wama, Nuristan V/48 (KP).
- 9. Acrocephalus melanopogon (Temminck) moustached warbler

 A. m. mimica Madarasz. Passage migrant, probably breeding in the north, Danoghori V/37 (RM). Recorded from Jalalabad IX/77 (In), Bamian VIII (SM), Kabul III/80 (JS), Kandahar IV/1879 (St.J), Lashkargah XI/73, II/75, Baghlan II/75 (FR).
- 10. Acrocephalus concinens (Swinhoe) blunt-winged reed warbler

 A. c. harringtoni Witherby. Summer visitor, recorded from the

 Danaghori Plains V/37 (RM).
- 11. Acrocephalus agricola (Jerdon) paddy field warbler

 A. a. brevipennis (Severtzov). Passage migrant, breeds in the NW
 (HE). Recorded from Jalalabad IX/77 (In), Bande Amir VIII, IX/69,
 Kole Hashmat Khan IV, V/64-66 (JN), Kabul VIII-X/79 (JS), Faizabad
 IV/49, Bamian IX, X/49 (KP), Logar VI/34 (Ma), Ghorband (N), Douchy (N),
 Danaghori (N) IV, V/37 (RM).
- 12. Acrocephalus dumetorum Blyth Blyth's reed warbler

 Common passage migrant and summer visitor. Breeds north of Hindu

 Kush and probably also in the south (HE). Recorded from Bande Amir

 VIII, IX/69 (SM), Bamian IX/49 (KP), Ajar Valley V, VII/76-77 (TE),

Mazar-i-Sharif VIII/49 (KP), Kabul VIII-X/79 (JS), Kunduz, Ghorband (N) br V/37 (RM), Kandahar IV/69 (JV), Byan Khel (E) V/1879 (WR), Pashki (E) V/48, Gardez V/49, Obeh VII/49 (KP).

- 13. Acrocephalus scirpaceus (Hermann) reed warbler

 A. s. fuscus (Hemprich and Ehrenberg). Recorded from Herat VII/49

 (KP), Ajar Valley V/77 (TE) and Kole Hashmat Khan br V/67 (AP).
- 14. Acrocephalus stentoreus (Hemprich and clamorous reed warbler
 Ehrenberg)

 A. c. brunnescens (Jerdon). Common passage migrant and summer visitor. Breeds in suitable habitat throughout the country. Project records from Kole Hashmat Khan V-VIII/79 (JS).
- 15. Acrocephalus arundinaceus (Linnaeus) great reed warbler

 A. a. zarudnyi Hartert. A single record from Tirpul (NE) IV/1885
- Hippolais pallida (Hemprich and Ehrenberg) olivaceous warbler

 H. p. elaeica (Lindermeyer). Summer visitor, breeding in the north

 (HE). Recorded from Haibak br V/37 (RM), Maimanah (Jalaiar) VI/1885

 (CY), Shindand, Adreshkan (W) V/69 (JV).
- Hippolais caligata (Lichtenstein)

 H. c. caligata. Rare migrant. H. c. rama (Sykes). Common summer visitor, breeding throughout (HE). Recorded from Bamian IX/49 (KP), Bande Amir VIII, IX/69 (SM), Douchy (N) V/37 (RM), Kila Wali (N), Darband-i-Kil Rekhta (N) IV, V/1885 (CY), Gungazai (SW) V/1881 (CS), Herat, Hasrate Baba (W) V/77 (RP), VII/49 (KP), Kandahar VI/1880 (CS), IV/69 (JV), Farah IV/49, Lashkargah (S), V/49, Ab-i-Estada V/49, Mukur VII/49 (KP), Mazar-i-Sharif, Haibak (N) VII/49 (KP), Kabul IV, V/64-66 (JN), IX/79 (JS), V, VI/75 (FR), VIII/33 (Ma, ssp caligata).
- Hippolais languida (Hemprich and Ehrenberg) Upcher's warbler Passage migrant and summer visitor, probably breeding (KP). Recorded from Paigah Kotal VIII/49 (KP), Ajar Valley VI/76 (TE), Ab-i-Estada, Dashte Nawar V/76 (TE), Kabul IV, V/64-66 (JN), Shar-i-Gholghola (C) VIII/72, Jam VIII/72 (VV), Kambao (NW) V/1885 (JA).
- 19. Sylvia mystacea Menetries Menetries' warbler
 Passage migrant. Recorded from the estuary of the Farah Rud in
 Seistan III/49 (KP), Kuhak (SW) III/05 (JC), Adreshkan (W) V/69 (JV),
 VI/1885 (CY).
- 20. Sylvia nana (Hemprich and Ehrenberg) desert warbler

 S. n. nana. Passage migrant, probably breeding in SW (HE). Recorded from Bagrabad, Faizabad and Farah Rud Estuary III-IV/49, Bamian IX, X/49 (KP), Gulran (NW) III/1885, Tirpul (NW) IV/1885 (JA).
- 21. Sylvia hortensis (Gmelin)

 S. h. jerdoni Blyth. Passage migrant and summer visitor. Probably breeding in NW (KP). Recorded from Obeh (C) VII/49 (KP), Bande Amir V/1885 (JA), Ghorband (C), Bamian IV/37 (RM).

- 22. Sylvia nisoria (Bechstein)

 S. n. merzbacheri Schalow. Passage migrant, breed in Badakhshan (HE).

 Recorded from Pashki (E) V/49 (KP), Bande Amir VIII, IX/69 (SM),

 Wakhan IX/69 (JN), Danaghori (N) V/37 (RM), Kumanibhest (NW) V/1885
- Very common passage migrant. Also summer visitor and breeding in suitable habitat throughout. Three pronounced subspecies recognized of which S. c. minula Hume is the commonest passage migrant in the Kabul area. Vaurie mentions possible breeding in the northwest, Seistan and Turkestan. Project records Kabul VIII-XI/79 IV/80 (JS), Ticehurst and Whistler is commoner in Seistan (IV/49 (KP)).
- 24. Sylvia communis Latham

 S. c. icterops Menetries. Passage migrant, recorded breeding from Amir VIII-IX/70 (SM), Bamian IX/49 (KP), Kabul V/77 (TE), VIII, IX/79 (JS), Adreshkan (W) V/69 (JV).
- 25. Phylloscopus occipitalis (Blyth)

 P. o. occipitalis. Summer visitor and breeding in Nuristan and Safed Koh. Recorded from Wama and Pashki (E) br IV-VII/48 (KP), N Salang VII/70 (SM), Safed Koh br VI/1879 (WR), Chaga Sarai, Kamdesh (E) (NE), Iskarzir (NE) VII/37 (WK).
- 26. Phylloscopus tytleri Brooks

 Breeding in the east. Recorded from Pashki (E) br V, VI/48 (KP),
 N Salang br VII/70 (SM) and Barikot-Kamu (E) br IV/69 (JV).
- Phylloscopus nitidus Blyth

 Passage migrant, breed in the northwest (HE). Recorded from Dashte
 Nawar V/76 (TE), Bande Amir VIII/70, N Salang VII/70 (SM), Akrobat
 (N) IV/37, Douchy-(N) V/37 (RM), Kabul X/79 (JS), Kandahar IV, X/1879
 (St.J).
- 28. Phylloscopus trochiloides Sundevall greenish warbler

 P. t. ludlowi Whistler. Summer visitor in Nuristan and Badakhshan.
 Recorded from Wama and Pashki (E) IV-VI/48 (KP), Small Pamir VIII/72 and Wakhan (JN).

 P. t. viridanus Blyth. Passage migrant, recorded from Pashki (E) VII/48 (KP), Big Pamir VIII/72 (JN), Kabul IX/79 (JS), IV/34 (Ma), Bande Amir VIII, IX/70 (SM). The greenish warbler is also recorded from Wakhan VII/71 (No), Ab-i-Estada V/76 (TE), Chorband (C) IV/34 (Ma).
- 29. Phylloscopus magnirostris Blyth large-billed leaf warbler Passage migrant, breeding in Safed Koh (HE). Recorded from Kabul
- 30. Phylloscopus proregulus (Pallas) yellow-rumped leaf warbler

 P. p. simlaensis Ticehorst. Breeding in Safed Koh (HE).
- 31. Phylloscopus subviridis (Brooke)

 Brooke's willow warbler

 Breed in eastern Afghanistan: Wama and Pashki (E) IV, VI/48 (KP),

 Safed Koh common br IV, V/79 (WR).

- Phylloscopus inornatus Blyth inornate willow warbler

 P. i. humei (Brooks). Passage migrant, breeding in Badakhshan and
 E. Hindu Kush (Charikar), winter visitor in the east and south (HE).

 Recorded from Tilli (NE) VII/48, Bamian IX-X/49, Wama and Pashki (NE)

 IV-V/48 (KP), Bande Amir VIII-IX/70 (SM), Wakhan VII/71 (No), Panjah

 (NE) IV/1874 (Bi), Dara-e-Shaur (Big Pamir) VIII/72 (JN), Jalalabad
- 33. Phylloscopus schwarzi (Radde) Radde's willow warbler The only record is from Danaghori (N) V/37 (RM).
- 34. Phylloscopus griseolus Blyth olivaceous willow warbler Breed in eastern and northeastern Afghanistan, passage migrant (HE). Recorded from the Pech-Parun Valleys (E) br IV-VI/48, Miyan Deh (NE) VII/48 (KP), Byan Khel (E) br VI/1879 (WR), Ghorband br, Bamian IV/37 (RM), Bande Amir VIII/70, N Salang VII/70 (SM), IV-VIII/72, Paghman IV-VIII/72, Darwaz VII/72 (JN).
- 35. Phylloscopus neglectus Hume plain brown willow warbler Breeding in all the mountains, passage migrant (HE). Recorded from Faizabad (SW) IV/49, Obeh (W) VII/49, Sauzak Kotal (W) VII/49 (KP), N Salang VII/70 (SM),?Kunduz IV/78, Dashte Nawar V/77 (TE), Kandahar (St.J), II/69 (JV).
- 36. Phylloscopus sindianus Brooks mountain chiffchaff
 Passage migrant. Recorded from Kabul IV/80 (JS), Ghorband (N) ? br
 IV, V/37 (RM), Helmand (SW) (Ticehurst).
- 37. Phylloscopus collybita (Vieillot) chiffchaff

 P. c. tristis Blyth. Common passage migrant. Recorded from Wakhan Valley (NE) common VII-VIII/69 (No), Sardeh IV/78, Ajar Valley V/77 (TE), Gusalek (E) III/48, Farah Rud (Seistan) III/49, Faizabad (SW) IV/49, Farah IV/49, Bamian IX-X/49 (KP), Kabul III-IV/64-66 common, XI/64-66 less common (JN), common IX-XII/79, III-IV/80 (JS), Kabul Valley, south and west Afghanistan common IV/69 (JV). Whistler, 1944 gives many records throughout the country for April.
- Regulus regulus (Linnaeus)

 R. r. himalayensis Bonaparte. Breeding in the east (Vaurie).

 Recorded from Pashki (E) V, VI/48 (KP). R. r. tristis Pleske. Winter (Ma), Kabul XII/73-III/74, XI/74-III/75, N Salang XII/73, Kunduz XII/73, II/74 (FR).

Family Muscicapidae

- 1. Muscicapa sibirica Gmelin sooty flycatcher

 M. s. gulmergi (Baker). Breeding in the east (HE). Recorded from

 Safed Koh V/1879 (WR) and Pashki (E) V, VI/48 (KP).
- 2. Muscicapa ruficauda Swainson red-tailed flycatcher
 Breeding in the east (HE). Recorded from Safed Koh br V, VI/1879
 (WR), Usman Khel (Gardez) V/49, Pashki (E) V, VI/48 (KP), Pech Valley
 (E) br V, VI/66 (LT), N Salang VII/70 (SM), and ?Kunduz (TE).

3. <u>Muscicapa striata</u> (Pallas)

spotted flycatcher

M. s. neumanni Poche. Summer visitor and passage migrant. Recorded from Pashki (E) V/48 (KP), Pech River (E) V, VI/66 (LT), Sinsoe (E) IX/66 (JN), Kabul IX/79 (JS), Kole Hashmat Khan XI/66 (JN), Jam (C) V/77 (RP), VIII/72, Lashkargah (S) IX/72 (VV), III/74, 75, Pansjir V/75 (FR), Darra-i-Shikari (C) br VI/49, Panjao (C) br VI/49, Obeh (W) VII/49 (KP), Bande Amir (C) VIII/70 (SM), Douchy (N), Danaghori (N) Darre Shikari (N) V/37 (RM), Chil Gaz (NW) V/1885 (JA), Byan Khel (E) br VI/1879 (WR), Darwaz (NE) VII/72 (JN), Kandahar IX/1879 (St.J), IV/1881 (CS).

4. Fidecula superciliaris (Jerdon)

white-browed blue flycatcher

 $\frac{F. \ s.}{River}$ superciliaris. Recorded from Safed Koh VI/1879 (WR), Pech River (E) 2 400 m br V, VI/66 (LT).

5. Fidecula parva (Bechstein)

red-breasted flycatcher

Two subspecies \underline{F} . \underline{p} . \underline{p} and \underline{F} . \underline{p} . $\underline{albicilla}$ (Pallas). Passage migrant. \underline{F} . \underline{p} . $\underline{albicilla}$ is only once recorded from Ghorband (N) IV/37 (RM). \underline{F} . \underline{p} . \underline{p} parva is commonly recorded especially during April throughout Afghanistan up to 2 500 m. Project records from Kabul IX-X/79, IV/80 (JS).

Family Monarchidae

1. Terpsiphone paradisi (Linnaeus)

Asian paradise flycatcher

T. p. leucogaster (Swainson). Summer visitor to central and east Afghanistan. Recorded from Wama (E) IV-V/48, Pech Valley (E) VII/48, Barak (NE) VII/48, Kwaja Chisht (W) VII/49 (KP), N Salang VII/70 (SM), Kabul VI/77 (P&C), VI/79 (JS), Shigul (E) IV/69 (JV), Kandahar IV/1881 (CS), Logar IV/33, Ashraf IX/34 (Ma), Danaghori br, Ghorband br V/37 (RM).

Family Timaliidae

Panurus biarmicus (Linnaeus)

bearded tit

P. b. russicus (Brehm). Probable summer visitor to Seistan and northwestern Afghanistan (HE), reedlands of the Amu Darya (N) (Vaurie).

1. Turdoides caudatus (Dumont)

common babbler

T. c. huttoni Blyth. Resident and breeding in southern Afghanistan (HE). Recorded from Kandahar (Blyth, 1847), XII/1880 (CS), Farah Rud (SW) II-III/49, Farah II-IV, VI/49 (KP), Kuhak common IV/05 (JC), Kandahar, Lashkargah (S) I-XII/73-75 (FR), V/49 (KP).

Garrulax albogularis (Gould)

white-throated laughing thrush

G. a. whisleri Baker. Vaurie gives Afghan border or Mazara (C) eastwards.

2. Garrulax variegatus (Vigors)

variegated laughing thrush

G. v. similis (Hume). Summer visitor to the east (HE). Recorded for Pashki V-VI/48, Stiewe VI/48 (E) (KP), Barikot-Kamu (E) IV/69 (JV), N Salang XII/72, V, XII/73, IV/74, Kabul winter/73, 74, 74/75, Sarobi (E) II/75 (FR).

Garrulax lineatus (Vigors)

streaked laughing thrush

G. 1. bilkevitchi (Zarudnyi). Summer visitor to Badakhshan. Recorded by Vaurie. G. 1. gilgit (Hartert). Summer visitor of Nuristan. Recorded from Gusalek III/48, Wama IV/48 and Pashki V-VII/48 (E) (KP). Also recorded from Bande Amir (C) VIII/72 (VV).

Family Aegithalidae

Aegithalos leucogenys (Horsfield and Moore) white-cheeked tit 1. Resident and breeding in Nuristan. Recorded from Chagha Sarai (Moore, 1855), Pech Valley (E) br VI/66 (LT), Gusalek and Wama (E) br III-V/48 (KP), Barikot (E), Kamu (E), Kamdesh (E) IV/69 (JV), Daulatshah (E) V, VI/37 (WK). Also in Pakistanian Safed Koh br III-V/06 (CW).

Family Paridae

1. Parus rubidiventris Blyth

rufous-bellied tit

- F. r. rufonuchalis Blyth. Resident of northern, central and eastern Afghanistan. Recorded from Gardez V/35 (RM), Pech Valley (E) br VI/66 (LT), Kamdesh (E) IV/69 (JV), Wama, Pashki (E) IV-VII/48, Sauzak Kotal (NW) VII/49 (KP), Khudikhel (SE) V/37, Daulatshah (E), Sirotai (E) V, VI/37, Darami (NE) VIII/37 (WK), N Salang VII/70 (SM), I, III-IV, XI-XII/73-75, Kabul I/74 (FR), Hariab (E) V/1879 (WR), Terak (C) IX/39, Safed Sang (C) IX/39, Burchao (C) IX/39 (WK).
- Parus melanolophus Vigors crested black tit Resident of the east. Recorded from Safed Koh V, VI/1879 (WR), Kamdesh (E) IV/69 (JV), Pech Valley (E) 2 200 m br V, VI/66 (LT), Gusalek (E) II-III/48, Wama, Pashki (E) V/48, Usman Khel (Gardez) br V/49 (KP), V/35 (Blanchard), Khudi Khel V/37, Sirotai (E) VI/37
- 3. Parus cyanus Pallas

azure tit

P. c. flavipectus Severtzov. Resident to the northern Hindu Kush. Recorded from N Salang VII/70 (SM), I/73, III/75, XI/74 (FR), Ajar Valley V/77 (TE), Sufian (NE), Tirgaran (NE) VII/37, Iskan (NE) VIII/37, Gurzan (NE) X/39 (WK).

Parus major Linnaeus

great tit

Four subspecies are recognized: P. m. cashmirensis Hartert (or P. m. decolorans Koelz). Resident of Nuristan and Safed Koh. Recorded from Pech Valley (E) br V, VI/66 (LT), Pech-Parun Valleys (E) II-VI/48 (KP), Chagai Sarai (E) III/1840 (WG), Byan Khel (E) br IV/1879 (WR), Kamdesh, Kamu IV/69 (JV). P. m. decolorans Koelz. Resident of Kabul and eastern Afghanistan (HE). Recorded from Jalalabad, XII/37, Daulatshah, Pactak VI/37 (WK). P. m. ziaratensis Whistler. Resident of the area to the west and southwest of the Hindu Kush. Recorded from Herat VII/49, Khwaja Chisht (W) VII/49 (KP), Kandahar III/1879 (St.J) I/1881 (CS) X-XI/50, Herat X-XI/50 (WK). P. m. ferghanensis is synonym with P. bokharensis (Vaurie). P. major is recorded by FR from Kabul, Sarobi, Jalalabad, Kandahar but only

Parus bokharensis Lichtenstein

P. b. bokharensis. Resident and breeding in Badakhshan, westwards to Afghan Turkestan, the Paropamisus and Iranian border (HE). Recorded from mulberry orchards in N Salang VII/70 (SM), Mazar-i-Sharif II/74 (FR), Balkh IX, XI/37, Taskurghan IX/37, Tukzar IX/39 (WK), Haibak (N) br V/37 (RM), Palunda-Karrez Darra (W) XII/1884 (JA).

Family Sittidae

- 1. Sitta leucopsis Gould white-cheeked nuthatch

 S. 1. leucopsis. Resident of eastern mountainous Afghanistan.

 Recorded from Byan Khel (E), Peiwar (E) br V, VI/1879 (WR), Pech

 Valley (E) br VI/66 (LT), Pashki (E) V-VI/48 (KP), Gardez V-VI/35

 (Blanchard), Sirotai (E) VI/37 (WK).
- 2. Sitta europaea Linnaeus common nuthatch

 S. e. cashmirensis Brooks. Resident of eastern Afghanistan up to

 4 000 m (HE). Recorded from Pech-Parun Valleys (E) III-VII/48 (KP),

 Kamdesh (E) IV/69 (JV), Byan Khel IV/1879, Peiwar Kotal (E) br V/1879

 (WR), Khudi Khel (E) V/37, Daulatshah (E), Kail (E) VI/37 (WK), Pech

 Valley (E) 2 200-2 400 m br V, VI/66 (LT).
- 3. Sitta tephronota Sharpe great nuthatch
 S. t. tephronota. Resident of mountainous Afghanistan. Recorded throughout the Hindu Kush, Badakhshan. Project records from Ajar Valley III/79 (JS).

Family Trichodromadidae

Trichodroma muraria (Linnaeus) wall creeper

T. m. nepalensis Bonaparte. Resident of central, eastern and northeastern Afghanistan (KP). Winter visitor of Jalalabad and Amu Darya
Valley (JN), and of lower levels (Whistler, 1944). Project records
from Qargah XII/79, Ajar Valley III/78 (JS).

Family Certhiidae

Certhia himalayana Vigors

C. h. himalayana. Resident of eastern Afghanistan. Recorded from Pech Valley (E) 2 300 m br V, VI/66 (LT), Pashki (E) V/48 (KP), Kail (E), Sirotai (E), Ferajghan Pass VI/37, Jalalabad XII/37 (WK), Kamdesh (E) IV/69 (JV), Pushut IV/1840 (WG), Peiwar Kotal (E) br VI/1879 (WR), Kail (E), Farajghan Pass (E), Sirotai (E) VI/37, Jalalabad XII/37 (WK), G. h. taeniura Severtzov, resident of Paropamisus and probably Hindu Kush (HE). Recorded from Sauzak Kotal (NW) VII/49 (KP), Terak (C), Safed Sang (W) IX/39, Burchao X/39 (WK). The Himalayan tree creeper is also recorded from N Salang (C) XI/73, Kunduz XII/73 (FR).

Family Remizidae

Remiz pendulinus (Linnaeus) penduline tit

R. p. coronatus (Severtzov). Summer visitor to Badakhshan and winter visitor to southern Afghanistan (HE). Recorded from Tirpul (NW) IV/1885 (JA); Farah Rud Estuary (SW) (as R. p. caspius) III/49 (KP), Sarobi IV/74, Lashkargah II/75 (FR). R. p. nigricans (Zarudnyi) breed in the reedbeds of the Helmand Basin in Iranian Seistan (KP).

Family Nectariniidae

Nectarinia asiatica (Latham)

purple sumbird

N. a. brevirostris Blanford. Summer visitor to the Jalalabad region. Recorded from Kunar Valley VI (HE), Pech Valley (E) VIII/48 Kalaigulaman (Laghman) VI/37 (WK), Khyber Pass, Chagha Sarai (E), Barikot (E), Kamu (E) IV/69 (JV). JV calls it a probable expansion

Family Zosteropidae

Zosterops palpebrosa (Temminck)

oriental white-eye

Z. p. egregia Madarasz. One record from Jalalabad XII/37 (WK).

Family Oriolidae

Oriolus oriolus (Linnaeus)

golden oriole

0. o. oriolus. Passage migrant (HE). 0. o. kundoo Sykes. Common summer visitor and passage migrant from May on. Recorded throughout the country, also in Wakhan.

Family Laniidae

- Lanius isabellinus Hemprich & Ehrenberg isabelline shrike (sometimes treated conspecific with L. collurie). L. i. isabellinus. Winter visitor and common passage migrant throughout the country especially between February and March (FR) but not of the east. Also recorded from Wakhan IX/65 (JN) and Bamian IX-X/49 (KP). L. i. phoenicuroides (Schalow). Summer visitor and passage migrant. Recorded between May and November throughout the country but not from
- Lanius vittatus Valenciennes 2. bay-backed shrike L. v. nargianus Vaurie. Summer visitor to the south and east (HE), recorded from Kabul VI/75 (FR), Chakar Darra (C) VIII/72 (VV), Kole Hashmat Khan V/66 (JN), V/77 (TE), VI/77 (P&C), Khinjan (C) br VII/33 (Ma), Sarobi (E) br VI/72, Arghandab br VI/72 (ES), Lashkargah IV/73, Kandahar V/74 (FR), IV/1881 (CS), Robat (SW) V/1896 (FM), Panjao (C) VIII/72 (VV), Ghorband (C) VI/49 (KP), Ab-i-Estada VI/79 (JS), Karizmir, Baghlan VI/77 (P&C), Tirpul (NW) IV/1885, Kambas (NW)
- 3. Lanius schach Linnaeus

black-headed shrike

L. s. erythronotus (Vigors). Common summer visitor and passage migrant in the whole of Afghanistan. Many records between April and October. Project records from Kabul VII-IX, XI/79 (JS).

Lanius minor Gmelin

lesser grey shrike

L. m. turanicus Fediuschin. Summer visitor to western and northern Afghanistan. Recorded from western Afghanistan, common VIII/72 (VV), IV/69 (JV), V/77 (RP), Obeh (W), Murghab, Pul-i-Khumri (N) VII/VIII/49 (KP), Arghandab (SE), Arghestan (SE) VI/72, Bamian VII/72, Panjao (C), Waras (C) VIII/72 (ES), Danaghori, Tash Kotal br, Kunduz br V/37 (RM), Akinjaram VII/33 (Ma), Kunduz, Pol-i-Chamra (N), Ghazni IV, V/73-75 (FR), Tirpul, Khusan (NW) IV/1885 (JA), Kandahar IV/1881 (CS), ?Ab-i-Estada-Kabul V/76 (TE).

- Lanius excubitor Linnaeus

 L. e. aucheri Bonaparte. Recorded from Shore Kaltegai (NW) V/1885

 (JA), L. e. pallidirostris Cassin. Winter visitor to southern

 Afghanistan (KP), (HE): Kandahar IX/1879 (St.J). Frequently

 reported throughout the country between X-III/73-75 (FR) also from

 Murghab, Guldara (N) VIII/72 (VV), and Pol-i-Chamra (N) VII/74 (FR),

 Kabul Valley IV/77 (TE).
- 6. Lanius senator Linnaeus woodchat shrike
 L. s. niloticus (Bonaparte). Recorded from Farah-Rud Estuary,
 Seistan III/49 (KP), and Girishk II/72 (ES).

Family Dicruridae

- 1. Dicrurus macrocercus (Vieillot)

 B. m. albirictus (Hodgson). Summer visitor of the east (HE). Many records from Nuristan, Sarobi and Jalalabad between May and June. Also recorded from Safed Koh 1879 (WR), Kabul IV, V/74 (FR), Dashte Nawar-Kabul V/76 (TE), Darunta II/79 (JS), Kunar, Pushut (E) III/1840 (WG), Jalalabad VIII/72 (VV), IX/77 (In), br V/37 (RM).
- 2. Dicrurus leucophaeus Vieillot ashy drongo

 D. 1. longicaudatus Jerdon. Summer visitor to eastern Afghanistan, south of the Hindu Kush (Vaurie, HE). Recorded from Khudi Khel (E), Daulatshah (E) IV-VII/37 (WK).

Family Corvidae

- 1. Garrulus lanceolatus Vigors lanceolated jay
 Eastern Afghanistan and Safed Koh. Recorded from Pansjir Valley
 VI (HE), Gusalek, Wama, Pashki (E) II-V/48 (KP), Barikot (E),
 Kamdesh (E) IV/69 (JV).
- Pica pica (Linnaeus)

 P. p. bactriana Bonaparte. Resident and common throughout

 Afghanistan, except the eastern part, where it does not occur and
 the magpie is scarce in the south.
- Nucifraga caryocatactes (Linnaeus) nutcracker

 N. C. multipunctata Gould. Resident in eastern Afghanistan (HE).

 Recorded from Wama, Pashki (E) IV-VI/48 (KP) and Peiwar Kotal (E)

 1879 (WR).
- Pyrrhocorax graculus (Linnaeus) alpine chough

 P. g. digitatus Hemprich & Ehrenberg. In the Hindu Kush from Herat to Pamirs and Nuristan (HE). Recorded from Pech Valley (E)

 IV-VI/48, Weran, Kokcha, Warduj, Sanglich Valleys (NE) VII/48 (KP), Salang VII/70 (SM), Ajar Valley III/79 common (JS), Wakhan, Big and Small Pamir VII-VIII/71 very common (No), Khalachan IX/33 (Ma), Salang II, VI, VII, XI, XII/73-75 (FR).
- Pyrrhocorax pyrrhocorax (Linnaeus) red-billed chough

 P. p. docilis (Gmelin). Resident of the Hindu Kush, Nuristan and Pamirs (HE). Recorded from Pech Valley (E) VI/48, Wera, Kokcha, Sanglich Valleys (NE) VII/48, Maimansh VII/49, Bamian and Darra-i-Shahidan (C) IX-X/49, Kandahar II/49, Farah II/49 (KP), Ajar Valley III/79 (JS), Salang VII/70 (SM), Wakhan, Big and Small Pamir

Salyan - Salian Samangan 36015' 68003' Samuli Afghan-Baluch border Sanglich 36020' 71014' Sardy Lake Kabul Valley Sar ·i-Kol = Zor Kol 37027' 73042' Sarkhad-i-Wakhan 36059' 73026' Sarobi 34040' 69046' Sarghae Wakhan Sar-i-Asia = ?Khush Asia 34059' 62019' Sarai Siah Ab Cave 32012' 62019' Sauzak Pass 62030' 34030' Scham Schirrohr = Shamshir Ghar Seistan SW Afghanistan Senza 35010' 71010' Shaharistan SW Afghanistan Shahar Shamba 35043' 63058' Shahgul = Shigul 34048' 71005' Shar-i-Chasma = Chasma-i-Cher

Shahidan E 34050' 67037' Shahidan S 32031' 64048' Shahrak 34006 64018' Shamshir Ghar 31033' 65028' Shar-i-Gholghala central Afghanistan Shibar Pass 34055' 68018' Shibergan = Shibargan 36040' 65042' Shigul 34048' 71005' Shindand = Sindand 33016' 62005 Shombul 34055' 68018' Shorawak 31048' 64016' Sia Koh Mts SW mountain spur Sim Koh Mts NW mountains Sirotai = Saroti 33028' 69025' Small Pamir 37005' 74020' Spin Daldak 31000' 66023' Stiewe 35030' 70056' Sufak N central Afghanistan

36005' 68035'

Sulaiman Mts Pakistan Supian Badakhshan Surkhab R 34021' 70000' Sust-e-Bala 37000' 72040' Taimani Watt E Afghanistan Tajan R extreme NW Afghanistan Tagau 34050' 69040' Taliqan 36045' 69035' Tarnak Valley north of Kandahar Tashgurghan 36044' 67041' Tash Kotal 36040' 67040' Terghem Qorum 37015' 74020' Tilli 35050' 70038' Tirgaran = Tigaran 36039' 71008' Tirish Mir Pakistan Tirpul 34034' 61020' Tolibai = Tulibai 37010' 72057' Tomam Agha Iran Tsah Gorgan 36045' 67040' Turkestan N plains Unai Pass 34027' 68025' Usman Khel 33052' 69037' Victoria Lake = Sar-i-Kol Waghdjir 37005' 74020' Wakhan E Badakhshan Wama 35007' 70044' Wardah S of Kabul Warduj Badakhshan Wazirabad - Balkh Weran Pass 35033' 70031' Yawkawlang Badakhshan Zabul Iran Zagin - 330 620 Zebak 36032' 71021' Zindijan = Zindajan 34010' 61015' Zor-i-Kol: 37027! 73042!

VII-VIII/71, IX/65 common (No), (JN), Darwaz VII/72 very common (JN), eastern Hindu Kush common br IV/37 (RM), Samangan, Salang, Doab, Kabul, Bamian, Kandahar 73-75 (FR), Kandahar flocks in winter 1879 (St.J), Bande Amir, Bamian VIII/72 (VV).

- 6. Corvus monedula Linnaeus jackdaw
 - C. m. soemmerringii Fischer. Breeding to the north of the Hindu Kush and winter visitor to the south (KP). Recorded from Haibak (N), Maimanah br V/37 (RM), VII, VIII/49 (KP), Faizabad (N), Keshem VI/77 (P&C), VIII/65 (JN), Kabul, Qargah XII-IV/64-66 (JN), II/77 (TE), Kandahar I/1881 (CS), IV/1879 (St.J), Murghab XII/1884 (JA), Maruchat (NW) XII/1884, Karawal (NW) IV/1885 (CY), Logar I/34 (Ma), Kunduz XI/74, Kabul XI/73, I/75, Pol-i-Shamri XII/73, Douchy II/74 (FR).
- 7. Corvus splendens Vieillot housecrow

 C. s. zugmayeri Laubmann. One record only, from Khost (E) VII/64
- 8. Corvus frugilegus Linnaeus rook

 C. f. frugilegus. Common winter visitor, except in the higher ranges between October and April, in the south arriving in January/February, leaving in March.
- 9. Corvus corone Linnaeus crow, hooded crow

 C. c. orientalis Eversmann. Summer visitor and breeding recorded in Qargah 73-75 br (FR), Paghman br 64-66 (JN), Haibak, Kunduz, Upper Kabul Valley, Unai Pass, Bamian, Doab and Danaghori br IV, V/37 (RM), common winter visitor between XI and V, in Jalalabad and Sarobi only in I and II (FR). C. c. sharpii Oates. Hooded crow, breeding in the north: Daulatabad (N) VIII/72 (VV), Mazar-i-Sharif 73-75 (FR), elsewhere between X and III common winter visitor.
- 10. Corvus macrorhynchos Wagler jungle crow

 C. m. intermedius Adams. Resident in the wooded parts of the east

 (KP) and probably southwest (HE). Recorded from Gusalek, Wama,

 Pashki, Stiewe (E) II-VI/48 (KP), Nuristan common IV/69 (JV), Hariab

 Valley (E) br V/1879 (WR).
- 11. Corvus ruficollis Lesson brown-necked raven

 C. r. ruficollis. Resident of arid northern (KP) and southern (HE)

 Afghanistan. Recorded from Andkhui (N) VII/49 (KP), Qargah VIII/74

 (FR), Seistan, Paropamisus (Vaurie).
- 12. Corvus corax Linnaeus raven

 C. c. subcorax Severtzov. Resident and widely distributed, many records throughout the country. In Wakhan probably C. c. tibetanus Hodgson (HE).

Family Sturnidae

Sturnus pagodarum (Gmelin)

Summer visitor and breeding in the east (HE). Recorded from

Jalalabad IK/77 (In), VI/37 (RM), Kabul VI/77 (P&C), /79 (JS), 1840

(WG), br V-VIII/74 (FR), Guldara (C) VIII/72 (VV), V-XI/69 (AP),

Mainakhel, Tagau (E), Chandau (E) VI/37, Marmakhel V/39 (WK).

2. Sturnus vulgaris Linnaeus

starling

- S. v. nobilior Hume. Breeding in the north and migrating to Seistan (HE). Recorded from Arghandab 1840 (WG), Haibak br V/37 (RM), Faizabad (N) VII/48 (KP). S. v. poltaratskyi Finsch. Passage migrant and winter visitor. Recorded from Kandahar 1840 (WG), Vagnami XI/33 (Ma), Gusalek (E) III/48, Bamian X/49 (KP). S. n. porphyronotus Sharpe. Winter visitor (HE). Kabul X/1839, Jalalabad 1840 (WG). The starling is also recorded from Ajar Valley (C) III/79 (JS), Faizabad (N) common VIII/65 (JN), Mukur III/65, Jalalabad II/66, Kandahar XII/64 (JN), Kabul III/78 common, II/77 (TE), X/64, X-XI/65, III/66 (JN), first XII/79 (JS), Kole Hashmat Khan IV/69, Kamdesh (E) ? br IV/69 (JV). FR records the starling as winter visitor between X/73-IV/74 throughout Afghanistan, except in
- 3. Sturnus roseus (Linnaeus) rose-coloured starling Summer visitor and abundant passage migrant, perhaps breeding in central (KP) and north (HE) Afghanistan. Records throughout Afghanistan for IV-V and VII-VIII. Summer records from Pech Valley (E) II-VI/48, Panjao (C) VI/49 (KP). The rose-coloured starling has an east-west migration from India to its breeding area in USSR to
- Acridotheres tristis (Linaneus) 4. common mynah A. t. tristis. Common resident of most of Afghanistan except the central highlands and the steppes in the SW (KP), vigorous expansion into Afghanistan since 1925 (Marien, 1951). Concentrated in urban
- Acridotheres ginginianus (Latham) 5. bank mynah Straggler, recorded from Kandahar, 1840 (WG).

Family Passeridae

Passer ammodendri Gould 1.

saxoul sparrow

- P. a. korejewi Zarudnyi and Härms. Breeding in NW Afghanistan (HE).
- Passer domesticus (Linnaeus) 2. house sparrow P. d. persicus Zarudnyi and Kudashev (intergrading to P. d. indicus Jardine and Selby). Summer visitor to the SW. Records from Faizabad (SW) and Farah (SW) IV/49 (KP). P. d. bactrianus Zarudnyi and Kudashev. Summer visitor and breeding in all provinces except the wooded parts of Nuristan and Seistan. In winter only in lower Kabul and in Kunar Valleys. Many records throughout Afghanistan. Project records Kabul: VII-IX/79, IV/80 (JS). P. d. parkini Whistler. Summer visitor to the Pamirs (HE).
- Passer hispaniolensis (Temminck) Spanish sparrow P. h. transcaspicus Tschusi. Summer visitor and breeding in northern and central Afghanistan. Passage migrant in the south (KP). Often mixed flocks with Passer domesticus, winter visitor to the lower Kabul Valley. A bird ringed in USSR 670 km to the north 8/K/73 was caught in Kunduz 28/X/73. Recorded throughout the country, especially

- 4. Passer rutilans (Temminck) cinnamon sparrow
 P. r. cinnamomeus (Gould). Breeding in Nuristan (HE), recorded from Otipore (E) 1840 (WG) and probably from Kunduz IV/78 (TE).
- Passer moabiticus Tristram

 P. m. yatii Sharpe. Resident in Seistan (KP, HE). Recorded from Lower Farah Rud, Seistan III/49 (KP), De Kamran-De Doda, large flocks X/1884 (JA), Na-i-Ab River (SW) III/03 (JC), Lashkargah ?V/75 (FR).
- 6. Passer montanus (Linnaeus) tree sparrow
 P. m. dilutus Richmond. Resident throughout Afghanistan. Common in villages and nearby fields (KP). Project records from Kabul: III-XII/79 I-IV/80 (JS).
- 7. Petronia brachydactyla Bonaparte pale rock sparrow Recorded from western Afghanistan. JV collected specimens in Shindand and Andreshkan (W) V/69.
- 8. Petronia xanthocollis (Burton) yellow-throated sparrow
 P. x. xanthocollis. Breeding in southern and eastern Afghanistan
 (KP) also passage migrant. Records from Kandahar V/1879 (St.J),
 Laghman V/37 (WK), Farah IV/49 (KP). P. x. transfuga (Hartert)
 summer visitor of southern Afghanistan (Vaurie).
- 9. Petronia petronia (Linnaeus) rock sparrow
 P. p. intermedia Hartert. Common resident in northern and central Afghanistan east to Faizabad and Ishkahim, winter visitor to lower altitudes. Breeding recorded from Salang N IV-VII/73-75 (FR), Dragon (C) VIII/72 (VV), Bamian, Haibak, Kunduz IV/37 (RM).
- Montifringilla theresae Meinertzhagen Theresa's snow finch Endemic for Afghanistan, occurs in the mountains NW of Kabul. Winter migrant to Soviet Union (KP). Records from Shibar Kotal (C), Bamian IV/37 (RM), Sabz Kotal (C) VIII/39 (WK), Shibar Kotal VI/49, Bamian IX-X/49 (KP), VI/73, VII/74 (FR), Unai Pass VI/49 (KP), IV/65 (JN), Bande Amir VI/77 (P&C), VII, VIII/70, Nil Kotal VII, VIII/70 (Wood & al. 78), Dashte Nawar VI/65 (JN), V/77 (TE), N Chakcharan (C) V/77 (RP), Koh-i-Baba (C), Chakcharan (C), Sia Koh (C) VIII/72 (VV).
- 11. Montifringilla nivalis (Linnaeus) snow finch

 M. n. alpicola (Pallas). Resident of the mountains of central, east and northeast Afghanistan uncommon in Nuristan (KP). Many records, breeding recorded from Salang, Bamian 73-75 (FR), Khinjan (C) VII/33 (Ma), Shibar (C), Unai (C) V/37 (RM), Unai-Panjao (C) V-VI/49 (KP). Project record from N Salang VII/79 (JS).

Family Estrildidae

Lonchura punctulata (Linnaeus) spotted mannikin
 L. p. punctulata. Recorded from Jalalabad IX/77 (In).

Family Fringillidae

1. Fringilla coelebs Linnaeus

chaffinch

F. c. coelebs Linnaeus. Winter visitor to northern, eastern and southern Afghanistan. Recorded from Kabul Valley II/77 (TE), Qargah XII/65 (JN), XII/79 (JS), Kabul common winter guest (JN), I-III/80 (JS), Burchao Pass X/39, Maimanah XI/37 (WK). According to FR between X-IV common winter visitor in: Kabul, Kunduz, Salang, Douchy (C), Sarobi, Pol-i-Shamri (C), Samangan, Mazar-i-Sharif, Lashkargah, Baghlan 73-75 (FR).

2. Fringilla montifringilla Linnaeus brambling

Winter visitor. Recorded from Kabul XI, XII/79, I-III/80 (JS), III/78 (TE), IV/37 (RM), Kabul Valley II/77 (TE), Qargah XII/79 (JS), II-III/65 (JN), Burchao Pass (C) X/39. Kang (SW) XI/37, Bala Murghab XI/39, Maimanah XI/39, Mazar-i-Sharif XI/39, Balkh X-XII/39 (WK), Chahiltan (C) XI, XII/33 (Ma), Panjah (NE) IV/1874 (Bi), Bamian IV/37 (RM) Kandahar I/1880 (St.J). Winter visitor to Kabul, Douchy (C), Pol-i-Shamri (C), Lashkargah (S), Salang, Baghlan, often in mixed flocks with chaffinches, buntings, between XI-III/73-75 (FR).

3. Serinus pusillus (Pallas) red-fronted serin

Common resident and breeding bird of mountainous Afghanistan including Nuristan and winter visitor to lower elevations (HE).

Breeding recorded from Doab (C) juv VII/74 (FR), Byan Khel VI/1879 (WR).

4. Carduelis chloris (Linnaeus) greenfinch

C. c. turkestanicus (Zarudnyi). Winter visitor to northern Afghanistan (HE). Recorded from Kabul XII/79 (JS), II, III/74 (FR), Balkh XII/37 (WK).

Carduelis spinoides Vigors

black-headed greenfinch

C. s. spinoides. Stuart Baker (in HE) recorded it from Afghan-Baluchistan border (Iran).

5. <u>Carduelis carduelis</u> (Linnaeus)

goldfinch

- C. c. paropanisi Kollibay. A widely distributed resident and winter visitor throughout the country but rather scarce everywhere (KP). Breeding recorded from Ghorband (C) V/37 (RM), Chahiltan (C) VI/33 (Ma). Project records from Chemptala (Kabul) XI/79 (JS).

 C. c. aubulata (Gloger). Rare winter visitor (HE), only recorded from Bamian IV/37 (RM).
- 6. Carduelis spinus (Linnaeus) siskin
 Recorded from Kabul I, II/75, Salang N XI/74 (FR).
- 7. <u>Carduelis cannabina</u> (Linnaeus) linnet
 - C. c. bella (Brehm). Resident of the high semideserts of the Hindu Kush, in the winter to lower altitudes. Recorded from Sauzak Kotal (W) VII/49 (KP), Ajar Valley III/79 (JS), Bande Amir, Sia Koh (C) VIII/72 (VV), Salang S V/75, Samangan I/75, II/74, Kabul I/75, Mazari-Sharif II/74 (FR), Gandachesma (C), Sufian (C), Tigaran (C) VII/37, Sabz Pass VIII/39, Gharchi (S) Gurzan (S) X/39, Kabul XI/39, Mazar-i-Sharif XII/37 (WK).

- 8. Carduelis flavirostris (Linnaeus) twite

 C. f. korejewi (Zarudnyi and Härms). Widely distributed resident and breeding in the high mountain pastures of the central Hindu Kush and Pamirs. Breeding recorded from Paghman IV/37 (RM), Unai, Shibar (C) VI/49 (KP), summer 64-65 (JN). Easterly records: Dashte Nawar VI/65 (JN), Tolibai (Big Pamir) VIII/71, Kotal-i-Dalez (Wakhan)
- 9. Loxia curvirostra Linnaeus crossbill

 L. c. tianschanica Laubmann. Probably passage migrant (HE), may wander into Afghanistan (Vaurie). Recorded from Kabul I/73 (FR), winter/78 (Haywood). Also Blyth reported them from Afghanistan.
- 10. Leucosticte nemoricola (Hodgson)

 L. n. altaica (Evertsmann). Resident in Badakshan (HE) and Nuristan VII/74 (FR), Wakhan, Small and Big Pamir VII, VIII/71 (No), Qala-i-Weran Valley, Miyan Deh (NE) br VII/48, Pashki, Stiewe E VI/48 (KP), Iskatul (NE), Sanglich (NE) VII/37 (WK).

 L. n. altaica (Evertsmann). Resident in Badakshan (HE) and Nuristan VII/74 (FR), Wakhan, Small and Big Pamir VII, VIII/65 (JN), V/75, Pandja (NE) IV/1874 (Bi), IX/65, Sust-e-Bala (Wakhan) VIII/72 (JN), Kamdesh (E) IV/69 (JV), Sarobi II/72, Yakawlang (C) VIII/72 (ES), Iskatul (NE), Sanglich (NE) VII/37 (WK).
- 11. Leucosticte brandti Bonaparte

 L. b. pamirensis Severtzov. Resident of Badakhshan (HE) and Hindu Kush (JN). Recorded from Salang VII/70 (SM), VI/75, VII/74 (FR), Hindu Kush VI/65, Small Pamir IX/65 (JN), VII/71, Wakhan VII/71 (No), Weran Valley (NE) VII/48 (KP), Shahidan (C) V/72, Hadjijak (C) VIII/72 (ES), VIII/72, Bande Amir VIII/72 (VV), Sanglich (NE) VII/37, Munjan Pass (NE) VII/37 (WK).
- Rhodopechys sanguinea (Gould)

 R. s. sanguinea. Resident of the high mountains of the Hindu Kush and the Pamirs (HE). Recorded from Panjao (C) VI/49 (KP), Hindu Kush VI/65 (JN), Bande Amir VII, VIII/70 (SM), VIII/72 (VV), Bamian (RM), Panjah (NE) IV/1874 (Bi), Wakhan common VII/71 (No), IX/65 (JN), Tolilai (Big Pamir) VIII/71 (No).
- Rhodospiza obsoleta (Lichtenstein)

 (Syn. Rhodopechys obsoleta). Resident of Afghanistan except of the northeast (HE). Recorded from Kandahar IV/65 (JN), XII/1880, III/1881 (CS), IV/69 (JV), Jam V/77 (RP), Adreshkan V/69 (JV), Qala Nau (N) br VII/49 (KP), VII, VIII/70, Bande Amir VII, VIII/70 (SM), Ajar Valley VI/76 (TE), Kabul spring/79 (JS), Lashkargah, Kandahar, Kabul, Kunduz between II-IX/73-75 (FR), Khelat-i-Ghilzai (S) Gogirdak (NW) III/1885 (CY), Khalat-i-Gilzai (S) X/37, Farah, Herat, Balkh XI/37 (WK).
- 14. Bucanetes mongolicus (Swinhoe)

 (Syn. Rhodopechys mongolica). Breeding on rocky slopes at high altitudes of the Hindu Kush (Vaurie). Recorded from Bamian br VI/49, LX-X/49, Shibar (C), Panjao (C) VI/49 (KP), VIII/72 (VV), Ghazni-(Bi), Sanglich (NE), Zebak (NE) VII/37, Sar-i-Asia (NW), Sabz Pass (NW), Balaghli (NW), Andkhui (NW) VIII/37 (WK).

- 15. Bucanetes githagineus (Lichtenstein)

 (Syn. Rhodopechys githaginea). B. g. crassirostris (Blyth). Resident of Afghanistan except of the northeastern part (HE). Recorded from Kabul IV/33, Chazni VI/33, Chakri Minai br VII/33 (Ma), Pansjir V/75 (FR), Kabul Valley II/77 (TE), Haibak (C) br V/37 (RM), V/37 (WK), Chakcharan (C), Farsi (NW) V/75 (RP), VIII/72 (VV), Shin Dand (70 km N of) VII/49 (KP), Adreshkan IV, V/69 (JV), Ab-i-Estada IV/70 (JS), Dashte Nawar V/77 (TE), Chakri Minai (E) VII/37, Ghazni VI/37 (WK).
- 16. Carpodacus erythrinus (Pallas)

 C. e. ferghanensis (Kozlova). Summer visitor and passage migrant.

 Breeding in Badakhshan, Nuristan and eastern Hindu Kush. Many
 IX/79 (JS).

 Common rosefinch

 Roseding in Badakhshan, Nuristan and eastern Hindu Kush. Many
 IX/79 (JS).
- 17. Carpodacus synoicus (Temminck)

 C. s. salimalii (Meinertzhagen). Resident in central Hindu Kush.

 Recorded from Bamian VI, IX-X/49 (KP), VII/74 (FR), IV/37, Shibar (C)

 IV/37, Akrobat IV/37 (RM), Doab VII/74 (FR), Sufak (N), Khami Deh (N)

 IX/39 (WK).
- 18. Carpodacus thura Bonaparte and Schlegel white-browed rosefinch

 C. t. blythi (Biddulph). Breeding in Safed Koh (HE). Not uncommon just over Pakistan border (CW).
- 19. Carpodacus rhodochlamys (Brandt)

 C. r. grandis Blyth. Resident in the mountains of the Hindu Kush and Badakhshan, west till Paropamisus (HE). Recorded from Pashki (E)

 V-VII/49 (KP), Darwaz VII/72 (JN), (but referred to C. r. kotschubeii)

 Salang N V/75 (FR), Pashat (E) 1840 (WG), Peiwar Kotal (E) VI/1879

 (WR), Maimanah XI/37, Safed Sang (N) IX/39, Burchao Pass (N) X/39,

 Gumandra (E) VI/37, Sirotai (E) VI/37, Terak (E) IX/39 (WK).
- 20. Carpodacus rubicilla (Gildenstädt) great rosefinch

 C. r. diabolica (Koelz). Resident in the northeast (HE). Recorded (WK).

 (WK).
- 21. Mycerobas icterioides (Vigors)

 Resident of Safed Koh (HE). Recorded from Peiwar Kotal (E) VI/1879
- 22. Mycerobas carnipes (Hodgson)

 M. c. speculigerus (Brandt). Resident in mountainous Afghanistan from Herat to Badakhshan and in Safed Koh (HE). Recorded from Salang VII/70 (SM), Koh-i-Baba (C) VIII/72, Sia Koh (C) VIII/72 (VV), Byan Khel (E) (WK).
- 23. Coccothraustes coccothraustes (Linnaeus) hawfinch

 C. c. humii Sharpe. Northwest Afghanistan. Recorded from Obeh (W)

 VII749 (KP), Ashroff Pass (W) X/78 (Wassey), Kabul I/74 (FR), Khossik

 Pass (SE) IV/1839 (WG). Also recorded just over the Pakistan border

 of Safed Koh (E) V/06 (CW).

Family Emberizidae

- 1. Emberiza leucocephalos Gmelin pine bunting

 E. 1. leucocephalos. Winter visitor and passage migrant. Recorded from 25 km W of Sarobi (E) I/66 (JN), Pushut 1840 (WG), Gusalek (E) III/48 (KP), Kabul I, II/74 (FR), XI/33, Chahiltan XII/33, Logar (C)
 - III/48 (KP), Kabul I, II/74 (FR), XI/33, Chahiltan XII/33, Logar (CI, XII/34 (Ma), Douchy (C) II/74, Samangan (N) II/74 (FR), Ab-i-Kamarra (W) XII/1884 (JA), Kandahar XII/1879, I/1881 (St.J), I/1882 (CS), Panjah (WE) IV/1884 (Bi).
- Emberiza citrinella Linnaeus yellow hammer
 C. erythrogenys Brehm. Rare winter visitor. Recorded from 25 km of Sarobi I/66 (JN), Kabul II/74, Mazar-i-Sharif II/74 (FR).
- Summer visitor to central and eastern Afghanistan (HE). Recorded from Wama, Pashki (E) IV-VI/48 (KP), Kamdesh (E) IV/69 (JV), Bande Amir VIII/70, Salang VII/70 (SM), br V/73-75, Pansjir (C), Doab (C), Bamian br V/73-75, Sarobi IV/74, Kabul IV/74 (FR), IX, X/79 (JS), Paghman, Darre Shihari (C), Doab (C) IV/37 (RM), Kunduz-Khanabad VIII/72 (VV), Obeh (W) VII/49 (KP), Ajar, Paktia V/77 (TE), Byan Khel (E) br (WR).
- 4. Emberiza cia Linnaeus rock bunting

 E. c. par Hartert. Common summer visitor to Hindu Kush, Badakhshan and Nuristan (HE), (KP). In the winter down to lower altitudes. Project record Kabul, common III/80 (JS).
- 5. Emberiza striolata Lichtenstein house bunting

 E. s. striolata. Rare summer visitor to eastern Afghanistan.

 Recorded from Jalalabad VI/66 (JN), Darunta br VI/66 (JN), IV/69 (JV),
 and probably Herat V/77, Qala Nau br V/77 (RP).
- 6. Emberiza cineracea Brehm ashy-headed bunting
 ?E. c. semenowi Zarudnyi. One record from Sarobi I/72 (ES).
- Passage migrant in northwest Afghanistan (HE). Recorded from Tirpul common IV/1885 (JA).
- 8. Emberiza buchanini Blyth grey-necked bunting

 E. b. buchanini. Summer visitor and breeding in the Hindu Kush and
 Badakhshan. Passage migrant (HE). Recorded from Tera Kotal (Gardez)

 V/49, Maidan (E) VI/49, Panjao (C), Shibar Kotal (C) VI/49, Bamian

 IX/49 (KP), IV/37, Paghman br V/37 Doab (C) IV/37, Ghorband br (C) V/37

 (RM), Pansjir, Salang, Pol-i-Shamri (C), Samangan (N), Doab (C) from

 April 73-75 on (FR), Bande Amir VIII/70 (SM), VIII/72, Jam VIII/72 (VV),

 Kole Hashmat Khan IX/65 (JN), Kabul V/33, Jangal (C) IX/33 (Ma), Obeh

 (W) VII/49 (KP), Kambao (NW) V/1885 (JA), Wakhan IX/65 (JN).
- 9. Emberiza fucata Pallas chestnut-eared bunting

 E. f. arcuata Sharpe. One report from July (Scully), referred in

 Whistler, 1944.
- 10. Emberiza rustica Pallas rustic bunting

 E. r. rustica. Two records only: Kabul III/80 (JS) and Sarobi XII/71

- 11. Emberiza pusilla Pallas little bunting Recorded from Lokhi (SW) XI/71 and Sarobi XII/71 (ES).
- 12. Emberiza schoeniclus (Linnaeus)

 E. s. pallidior Hartert. Winter visitor. Recorded from Farah Rud
 Estuary, Seistan II/49 (KP), Kole Hashmat Khan III/65 (JN), Kabul
 I/74 (FR), Bagrani XI/33 (Ma), Kandahar II/1881 (CS), Chahar Shamba
 (NW), Kara Bel (NW), Gulla Chasma (NW) II, III/1885 (CY).
- 13. Emberiza bruniceps Brandt red-headed bunting
 Summer visitor and passage migrant. Widely distributed as a breeding
 bird around towns and villages between April and August but not in the
 south (KP). Many records throughout Afghanistan. Project records
 from Kabul VIII, IX/79 (JS).
- 14. Emberiza melanocephala Scopolo black-headed bunting

 One record from Melcharez, 60 km S of Kandahar, flocks IV/1881 (CS).
- M. c. buturlini (Johansen). Summer visitor and breeding in northern Afghanistan (HE). Recorded from Danaghori (N) br V/37, Khanabad (N) br V/37 (RM), Kunduz V/75, Samangan (W) II/72, 73 and between I-II/74 from Kunduz, Kabul, Douchy (C). Pol-i-Chara (C) (FR). Unsubstantiated reports from the Ajar Valley X/78 (Wassey) and Chakcharar V/77 (RP).

Appendix 7

REPTILES OF AFGHANISTAN

ORDER CHELONIA

Family Testudinidae

1. Testudo horsfieldii Gray

Afghan turtle

Loamy and sandy steppes throughout Afghanistan up to elevation of 3 000 m. Common at lower elevations.

Family Trionychidae

1. Trionyx gangeticus Cuvier

One recorded, Khost (E) VII, VIII/69 (Schneider).

ORDER SQUAMATA

Family Gekkonidae

Agamura femoralis Smith

Not recorded from Afghanistan but occurs in neighbouring parts of Northwestern Baluchistan (Pakistan).

1. Agamura persica (Duméril)

Afghanistan, south of Hindu Kush and the lowlands in the west along Iranian border; at Paghman up to 2 800 m elevation (LA), Chah-i-Angir (S) 330 m 1950 (LA), Girishk IX/57 (OW).

2. Alsophylax tuberculatus (Blanford)

Seistan, Helmand River basin and higher regions to the east (2 700 m at Ghaom-i-Faringi). Recorded from Kandahar 1 425 m (LA), 30 km S of Mukur 2 300 m VII/64 (CC), Chah-i-Angir (S) 330 m 1950 (LA).

3. Alsophylax pipiens (Pallas)

Vicinity of Kabul. Recorded from Kabul 2 000 m VII, VIII/64 (CC) IV, V/50, Paghman VIII/50 (LA), Oukak (E) 1965 (Mertens).

4. <u>Crossobamon</u> <u>eversmanni</u> (Wiegmann)

Western and northern Afghanistan (LA).

5. Crossobamon lumsdeni (Boulenger)

Only known from Afghan-Baluch border between Nushki and Helmand (Boulenger, 1887).

6. Crossobamon maynardi (Smith)

Helmand River basin (LA), Afghan Baluch Boundary (Maynard and McMahon).

- 7. Cyrtodactylus caspius (Eichwald)
 - In Afghanistan to the north of the Hindu Kush. Recorded from Paghman 2 440 m, Mazar-i-Sharif 457 m IX/65, 100 km E of Faizabad VIII/65 (LA), Bala Murghab 470 m V/64, Mazar-i-Sharif 1964 (BK), Sarkol X/57, Pol-i-Khomri (N) X/57 (OW).
- 8. Cyrtodactylus fedtschenkoi (Strauch)

North of the Hindu Kush and Paghman, west and south to Baluchistan. Recorded from 50 km W of Dilaram (W) 823 m XI/65, Kandahar 1 425 m (LA), 50 km W of Girishk (S) 1 200 m VIII/64 (CC), Kandahar IV/58, Girishk IV/58, Farah (SW) IV/58 (OW), Bala Murghab (NW) (GB).

- Crytodactylus persicus (Nicolsky)
 Recorded from Kabul, 1951 (LA).
- 10. Cyrtodactylus russowii (Strauch)
 Wermuth, 1965, records this species from northern Afghanistan.
- 11. Cyrtodactylus scaber (Heyden)

South and southeast from low elevations up to 1 700 m. Recorded from Kandahar 1 425 m (LA), Herat 1 000 m VIII/64, Khost (E) 1 400 m VII/64 (CC), Jalalabad IV/67 (BK), Sharisafa (E) (LA).

12. Cyrtodactylus watsoni (Murray)

Kabul River valley to the west as far as Jalalabad. Recorded from Jalalabad (LA), also by (GB) as \underline{C} . $\underline{kachhensis}$ (Stoliczka).

- 13. Cyrtodactylus spec. nov.
- Helmand River basin from Farah to Kandahar (LA).
- Eublepharus macularius (Blyth)

 Eastern Afghanistan to the south of the Hindu Kush, even in Kabul townside up to 1 800 m elevation. Recorded from Jalalabad 732 m (LA), Minla (E) 1 200 m, Chirikar 1 800 m VII, VIII/64 (CC), Jalalabad 500 m III/65 (BK), II/57 Gulbahar (Kabul) II/57 (OW).
- 15. Hemidactylus flaviviridis RUppell

Only known from Paghman 2 440 m and Jalalabad 732 m, probably due to human agency (LA). Recorded from Jalalabad II-III/66, IV/67 (BK).

16. Teratoscincus bedriagai Nikolsky

Helmand River basin to at least 1 600 m elevation. Recorded from Qala-i-Kang, 580 m XI/65, Kandahar 1 425 m (LA), Chah-i-Angir (S) 1950 (LA).

- 17. <u>Teratoscincus microlepis</u> Nikolsky Afghan-Baluchistan border (LA).
- 18. Teratoscincus scincus (Schlegel)

Low deserts along the western and southern borders to 1 600 m elevation. Recorded from Qala-i-Kang (SW) 518 m XI/65, 50 m W of Dilaram (SW) (LA), Chah-i-Angir (S), 330 m 1950 (LA); Welmuth gives northern Afghanistan.

Appendix 3

LIST OF OBSERVERS (for literature see Bibliography)

F F R N R	CN C. Naumann JG J. Gaisler JH J. Hassinger JN J. Niethammer KH. Habibi KP K. Paludan KZ K. Zimmermann JO D. Meyer Oehme JO G. Nogge KN. Rodenburg KA V. Aellen	Reptiles & Amphibians	BK CC GB LA OW RP	B. Kral Clark, R.J. et al G. Brück A. Leviton and S. Anderson O. von Wettstein- Westersheim R. Petocz
Birds A B C C C C E F F F H I I J A J C J N A N O P E S W R R R R F S C W W W W W W W W W W W W W W W W W W	J. Biddulph C. Swinhoe C. Whitehead C. Yate E. Smith F. Maywood F. Reeb F. Hue and R.D. T. Inskipp J. Aitchison J. Cumming J. Niethammer et J. Sayer and A. J. Vieillard K. Paludan R. Maconachie N. Annandale G. Nogge Parinder and Croc R. Meinertzhagen R. Petocz	al. van der Zon (partly ok	· '	

Family Scincidae

1. Ablepharus bivittatus (Menetries)

A.b. lundbergi Wettstein. Upland Afghanistan to more than 3 000 m elevation (LA). Recorded from Obeh VII/57 (OW).

2. Ablepharus grayanus (Stoliczka)

In Afghanistan recorded only from the Sabzak Pass 2 190 m VI/59 (OW).

3. Ablepharus pannonicus Lichtenstein

Foothills and mountains in most of Afghanistan to 2 500 m elevation. Recorded from 36 km S of Kabul 2 300 m IV/50 (LA), Qal-i-Chahrak (C) VIII/57, Qal-i-Nau (NW) X/57, Bamian IX/57 (OW).

Eumeces blythianus (Anderson)

Blythbskink

Not yet recorded from Afghanistan but from Khyber Pass Region (Pakistan) (LA).

4. Eumeces schneideri (Daudin)

Schneider's skink

E. s. princeps (Eichwald). Lower steppes north of the Hindu Kush (LA). E. s. zarudnyi. Helmand Basin (LA).

5. Eumeces taeniolatus (Blyth)

Steppes near Pandjvai (Kandahar) and Tajan River on Afghan-Iran-Turkmen border (LA). Recorded from Bala Murghab (NW) 50 m (GB), IV/64, 20 km E of Jalalabad V/67 (BK), Kandahar IV/58 (OW).

6. Mabuya aurata (Linnaeus)

Possibly recorded from Tajan River at the Afghan-Iran-Turkmen borders (LA).

7. Mabuya dissimilis (Hallowell)

Kabul river valley to 1 200 m elevation. Recorded from Jalalabad 732 m (LA), 15 km E of Nimla 1 200 m (CC).

Ophiomurus brevipes (Blanford)

Not yet recorded from Afghanistan but it is found in adjacent areas in Iran and USSR (Terent'ev and Chernov).

8. Ophiomurus tridactylus (Blyth)

snake skink

Sandy areas of Helmand Basin and adjacent areas. Recorded from 30 m E of Girishk 1 100 m VIII/64 (CC), Chah-i-Angir (S), 330 m /50 (LA) Kandahar, Helmand (Boulenger, 1887), Chaman (Alcock & Finn, 1896).

9. Scincella himalayana (Gunther)

Mountainous region of Nuristan (Pashki) 1948 (KP).

Family Lacertidae

1. Acanthodactylus blanfordii Boulenger.

Recorded from Afghanistan (Arnold, pers. inf.).

2. Acanthodactylus cantoris (Glinther) fringe-toed lizard

Kabul River valley and Helmand River basin. Recorded from Jalalabad 732 m (LA), 30 km E of Girishk (S) 1 100 m, 30 km SE Kandahar 1 300 m, Nimla (E) 1 200 m VII-VIII/64 (CC), Chah-i-Angir (S) 330 m 1950 (LA), Jalalabad II/66 (BK).

3. Eremias acutirostris (Boulenger) desert lacertid

Desert basin of southeastern Afghanistan. Recorded from HelmandNushki (Boulenger, 1887).

4. Eremias aporosceles (Alcock and Finn)

Along Afghan-Baluchistan border (LA).

5. Eremias aria Anderson and Leviton

Jalalabad Valley, endemic for Afghanistan. Recorded from Nimla (E) 1 200 m VII/64 (CC).

6. Eremias fasciata Blanford

Southern Afghanistan in the Helmand Basin. Recorded from Chah-i-Angir (S) 330 m 1950 (LA), Helmand Basin 1889 (Boulenger).

7. Eremias grammica (Lichtenstein)

Lowland regions of Afghanistan, north of the Hindu Kush. Recorded from Aq Chah (Mazar-i-Sharif) summer/64 (BK).

8. Eremias guttulata

E. g. watsonana Stoliczka

Common throughout Afghanistan, to 2 600 m elevation (LA).

9. Eremias intermedia (Strauch)

Valley of Tajan River at the Afghan-Iran-Turkmen borders (LA) and recorded from Aq Chah (Mazar-i-Sharif) summer/64 (BK).

10. Eremias lineolata (Nikolsky)

Lowland regions of Afghanistan, north of the Hindu Kush. Record from Bala Murghab (GB).

11. Eremias nigrocellata (Nikolsky)

Lowland regions of northern Afghanistan. Recorded from Faizabad (NE) 2 653 m VIII/65 (LA), Wakhan 3 100 m (RP).

12. Eremias regeli Bedriaga

Kabul River Valley. Recorded from 15 km E of Nimla 1 200 m VII/64 (CC).

13. Eremias scripta (Strauch)

Desert regions in the south (LA) and north. Recorded from Aq Chah (N) summer/64 (BK).

14. Eremias velox (Pallas)

E. v. velox. Northwestern Afghanistan, perhaps south to Seistan Basin. Recorded from Tajan River (NW) (LA) and New Gulran (NW) (LA).

E.v. persica Blanford. Common in southern Afghanistan northeast to Paghman, up to 2 700 m elevation (LA).

15. Ophisops jerdoni Blyth

snake-eyed lacertid

Family Anguidae

1. Ophisaurus apodus (Pallas)

Kabul River valley (LA).

sheltopusik

Northern Afghanistan up to almost 3 000 m elevation. Recorded from Faizabad (NE) 2 653 m VIII/65 (LA).

Family Varanidae

1. Varanus bengalensis (Daudin)

Bengal monitor

V. b. bengalensis. Kabul River valley, up to nearly 3 000 m elevation north of Jalalabad. Recorded from Jalalabad 2 600 m VIII/65 (LA), Kabul-Sarobi 1 200 m, 15 km E of Nimla (E) 1 200 m VII, VIII/64 (CC), Paghman, Jalalabad, Darunta, 80 km ESE of Jalalabad III-IV/66 (BK).

2. Varanus griseus (Daudin)

desert monitor

V. g. caspicus (Eichwald). Semi-desert areas of Murghab river, frequent at 400 m elevation and of Helmand basin and Harirud Valley. Recorded from Darya-i-Murghab River 400 m frequent (BK), Chah-i-Angir (S), 330 m 1950 (LA).

Family Agamidae

1. Agama agilis Olivier

agile agama

A. a. isolepis Boulenger. Semi-arid regions of Afghanistan to about $\frac{1}{2}$ $\frac{1}{500}$ m elevation. Recorded from Paghman 2 440 m VII/65, Maimanah 884 m IX/65, 100 km E of Faizabad (NE) VIII/65, Herat 1 000 m VIII/65 (LA), Herat 1 000 m VIII/64, Shindand (W), 30 km E of Farah (SW), 30 km E of Girishk (S), 30 km SE of Kandahar VII-VIII/64 (CC), 25 km N of Kandahar V/58 (OW), Chah-i-Angir (S) 330 m 1950 (LA), Bala-Murghab (NW) 420 m (GB), VI/64 (BK).

2. Agama agrorensis (Stoliczka)

Kabul River valley to at least 2 000 m elevation. Recorded from Jalalabad 732 m (LA).

3. Agama badakhshana Anderson and Leviton

Endemic for Afghanistan. Mazar-i-Sharif IX/65, 100 km east of Faizabad (NE) VIII/65, Paghman VII/65 (LA), Sanglech River (Pamir) 1978 (RP).

4. Agama caucasica (Eichwald)

Caucasian agama

A. c. caucasica. Afghanistan to 3 000 m elevation. Recorded from Paghman 2 440 m VII/65 (LA), 30 km N of Ghazni 1 400 m, Narai (E) 2 700 m, 45 km NE of Gardez 2 200 m, Paghman 2 500 m VII-VIII/64 (CC), Ghazni IV/50, Paghman VIII/50 (LA).

5. Agama erythrogaster (Nikolsky)

Northern Afghanistan, south through the mountain passes to Paghman, 1 000-2 700 m. Recorded from Maimanah 884 m VII/65, Paghman 2 440 m VII/65 (LA), Kharzar (Mazaradjat, C) VIII/57 (OW).

6. Agama himalayana (Steindachner)

A. h. himalayana. Recorded from Ishkamish (Pamir) (LA), Sanglich River (Pamir) to 3 500 m elevation (RP). Also in the Hindu Kush (Terent'ev and Chernov).

7. Agama lehmanni (Nikolsky)

Only recorded from Mazar-i-Sharif (457 m) IX/65 (LA).

8. Agama nupta De Filippi

A.n. nupta. Afghanistan south of the Hindu Kush, up to 1 800 m elevation. Recorded from 50 km W of Dilaram, 832 m, Kandahar 1 425 m (LA), Nimla (E) 1 100 m E of Kabul VII/64 (CC), Jalalabad III/65, II-III/66 (BK), (GB), Farah (SW) IV/58 (OW).

9. Agama nuristanica Anderson and Leviton

Endemic to Afghanistan. Only known from Kamdesh (Nuristan) at 1 342 m elevation X/65 (LA).

10. Agama ruderata Olivier

A. r. ruderata. Southern Afghanistan to about 2 500 m elevation. Recorded from Chazni 2 500 m, Charikar 1 800 m VIII/64 (CC).

11. Agama sanguinolenta (Pallas)

Northern Afghanistan (Terent'ev and Chernov).

12. Agama tuberculata Hardwicke and Gray

Doubtful record from Kabul (Smith, 1935), but also recorded from Jalalabad (GB).

13. Calotes versicolor (Daudin)

Indian bloodsucker

Only recorded from two localities in Afghanistan, highest at about 2 500 m elevation. Recorded from Kabul-Sarobi, 1 400 m VIII/64 (CC), Jalalabad II/65, 66 (BK).

14. Phrynocephalus clarkorum Anderson and Leviton

Helmand River basin. Recorded from SE of Kandahar, 1 300 m VIII/64 (CC).

15. Phrynocephalus euptilopus Alcock and Finn

Helmand Basin to the southeast of Darweshan (LA).

16. Phrynocephalus interscapularis Lichtenstein

P. i. interscapularis. Lowlands between Andkhui and Mazar-i-Sharif in northern Afghanistan to 700 m elevation. Recorded from Aq Chah (Mazar-i-Sharif) 1964 (BK).

17. Phrynocephalus luteoguttatus Boulenger

Helmand River basin up to about 1 300 m elevation. Recorded from 30 km SE of Kandahar 1 300 m VIII/64 (CC).

18. Phrynocephalus maculatus Anderson

P. m. maculatus. Afghan-Baluch border region and Darweshan in Helmand basin (LA).

19. Phrynocephalus mystaceus (Pallas)

P.m. galli Krassowsky. Sand dunes north of the Hindu Kush. Recorded from Aq Chah (N) summer/64 (BK).

20. Phrynocephalus ornatus Boulenger

Helmand River basin. Recorded from 30 km E of Girishk (S) IV/58 (OW), 1 100 m VIII/64 (CC), Chah-i-Angir (S) 330 m 1950 (LA).

21. Phrynocephalus reticulatus Eichwald

P.r. boettgeri Bedriaga. Northern lowlands of Afghanistan. Recorded from Aq Chah (N) summer/64 (BK).

22. Phrynocephalus scutellatus (Olivier)

Southern Afghanistan to at least 2 400 m elevation. Recorded from SE of Kandahar 1 300 m, Ghazni-Hukur (E) 2 400 m VIII/64 (CC), Chah-i-Angir (S) 330 m 1950, Karabakh (C) (LA).

23. Uromastix asmussi (Strauch)

Single record from Afghan-Baluch border.

24. Uromastix hardwickii Gray

Indian spiny-tailed lizard

Dry clayey soils along Kabul River valley to 700 m elevation. Recorded from 10 km ESE of Jalalabad 650 m II/66, IV/67 (BK).

Family Typhlopidae

1. Typhlops vermicularis Merrem

Vermiform blind snake

One record only, north of Herat (LA).

Family Leptotyphlopidae

1. Leptotyphlops blanfordi (Boulenger)

Kabul River valley (LA) and also recorded from Kuh-i-Malik Siah at the Afghanistan-Iran-Pakistan border (Alcock and Finn, 1896).

Family Boidae

1. Eryx elegans (Gray)

Recorded from Paghman, 2 440 m, VIII/65 (LA).

2. Eryx johnii (Russell)

brown sand snake

Mundi (Hissar) to the south of Kandahar (Murray, 1892).

3. Eryx tataricus (Lichtenstein)

E.t. speciosus Tzarevsky. Northern Afghanistan, Paghman and Southern Afghanistan. Recorded from Paghman 2 440 m VII/65, Maimanah 884 m IX/65 (LA) Bala Murghab 600 m, 40 km SW Jalalabad III/65 (BK), Chah-i-Angir (LA).

Family Colubridae

1. Boiga trigonata Smith

tree snake

B.t. melanocephala Annandale. Southern and western lowlands. Recorded from Chah-i-Angir (S) 330 m 1950 (LA).

2. Coluber harelinii Brandt

racer

Southern lowlands. Recorded from Chah-i-Angir (S) 330 m 1950 (LA), Aq Chah (Mazar-i-Sharif) summer/64 (BK), Bala Murghab 430 m III-IV (GB).

3. Coluber ravergieri Menétries

racer

C.r. plumbens (Cerrov). Northern and eastern part of Afghanistan between 700-2 700 m elevation. Recorded from Paghman 2 440 m VII/65 (LA), Kara Bakh (N of Kabul) (LA), Bamian 2 500 m 1940 (Smith).

4. Coluber rhodorachis (Jan)

race

Lower elevations on both the north and south side of the Hindu Kush. Recorded from Kandahar 1 425 m XI/65, Paghman 2 440 m VII/65 (LA), 15 km SW of Kabul 2 200 m IX/50 (LA), Jalalabad summer/64, Aq Chah (Mazar-i-Sharif) summer/64 (BK).

5. Coluber vetromaculatus Gray and Hardwicke

Boulenger, 1890 includes Afghanistan in the distribution but no documented reports (LA).

Eirenis persica (Anderson)

dwarf snake

Not yet in Afghan records but occurring in neighbouring places of Iran and Pakistan.

6. Elaphe dione (Pallas)

desert rat snake

Bollu-Kuch 10 km SE of Kunduz.

Lycodon striatus

Asian wolf snake

L. s. bicolor (Nikolsky). Not yet recorded from Afghanistan but only Iranian Seistan.

7. Lytorhynchus maynardi Alcock and Finn

Desert basins of southern Afghanistan (LA). Recorded from the Afghan Baluchistan border (Koh- Malik-do-Khand) (Alcock and Finn, 1896).

8. Lytorhynchus ridgewayi Boulenger

Northwestern and southern Afghanistan. Recorded from Chah-i-Angir (S), 330 m 1950 (LA), Chinkilok (Boulenger, 1887).

9. Matrix tessellata (Laurenti)

diced snake

N. t. tessellata. Afghanistan. Recorded from Mazar-i-Sharif 457 m IX/65, Paghman 2 440 m VII/65, Maimanah 884 m IX/65 (LA), Aq Chah (Mazar-i-Sharif) summer/64 (BK).

10. Oligodon arnensis (Shaw)

Semi-deserts, 3 km southeast of Jalalabad 560 m III/65 (BK).

11. Oligodon taeniolatus (Jerdon)

Found in the western Himalayas and recorded from Jalalabad (GB).

12. Psammophis leithii Günther

Semi-desert, 8 km from Jalalabad 650 m III/66 (BK).

13. Psammophis lineolatus (Brandt)

Afghanistan, desert areas. Recorded from Paghman, 2 440 m VII/65 (LA), Bala Murghab (NW) 400 m V/64, Jalalabad II/66, Kunduz V/67, Aq Chah (Mazar-i-Sharif) summer/64 (BK).

14. Psammophis schokari (Forskal)

Southern Afghanistan. Recorded from Chah-i-Angir (S). 330 m 1950 (IA), Helmand and Hamun to Khusan (Boulenger, 1896).

15. Ptyas mucosus (Linnaeus)

Asian rat snake

Afghanistan, including Kabul Valley and between Kabul and Kandahar. Recorded from Herat 1 800 m VITI/65, Kamdesh (E) 1 342 m X/65, Kandahar 1 425 m X/65, Maimanah 884 m IX/65 (LA) 40 km SW of Kabul 2 300 m IX/50 (LA).

16. Spalerosophis diadema (Schlegel)

S. d. schirazana (Jan). Northern lowlands and southern deserts. Recorded from Maimanah 884 m IX/65; Paghman 2 440 m (LA) Jalalabad 560 m summer/65, IV/67 (BK).

17. <u>Xenochrophis piscator</u> (Schneider)

Collected 10 km south of Jalalabad 600 m summer/65, Darunta (E)

II/66 (BK).

Family Elapidae

- 1. Bungarus caeruleus (Schneider) krait
 20 km E of Jalalabad 800 m semi-desert IV/66 (BK).
- 2. Naja oxiana (Eichwald)

 Northwest and south Afghanistan. Recorded from northern slopes of Selsala-Kohi-Beni (Herat Province, 800 m) V/64 (BK), Jalalabad 732 m (LA) (GB).

Family Viperidae

- 1. Agkistrodon halys (Pallas) Halys viper

 A. h. caucasicus Nikolsky. One record from Sauzak Pass near Herat

 1949 (KP).
- 2. Agkistrodon himalayanus (Günther) Himalayan viper
 One recorded from Wama, Nuristan 1948 (KP).
- 3. Echis carinatus (Schneider) saw-scaled viper

 Lower elevations, both north and south of the Hindu Kush. Recorded from Kandahar, 1 425 m, XI/65 Chah-i-Angir (S) 330 m 1950 (LA),

 Jalalabad 620 m III/65 (BK).
- 4. Eristicophis macmahonii Alcock and Finn McMahon's viper

 Southern deserts of Afghanistan in Registan and Dasht-i-Margo, below
 1 300 m (LA).

Pseudocerastes persicus (Duméril, Bibron and Duméril)

- P. p. persicus. Not yet recorded from Afghanistan but recorded from neighbouring desert areas of Iran and Pakistan (LA).
- 5. Vipera lebetina (Linnaeus)

 V. 1. turanica Cernov. Northwestern part of Afghanistan and Kabul River Valley. Recorded from Jalalabad, 732 m (LA).

Appendix 8

AMPHIBIANS OF AFGHANISTAN

ORDER CAUDATA

Family Hynobiidae

1. Batrachuperus mustersi Smith

Endemic, only known from the mountain streams of the Paghman Mountains, 3 000 m (Smith, 1940).

ORDER SALIENTIA

Family Bufonidae

1. Bufo andersonii Boulenger

South of the Hindu Kush in SE Afghanistan and along the Helmand River. To 1 500 m elevation. Recorded from Qala-i-Kang (SW) XI/65, Paghman, Jalalabad 732 m, Kandahar 1 425 m (LA), Khost 1 400 m VII/64 (CC).

2. Bufo viridis Laurenti

Common in suitable places throughout Afghanistan to an elevation of 5 000 m in the Himalayas (LA).

Family Ranidae

1. Rana cyanophlyctis Schneider

South of the Hindu Kush to elevation of 1 500 m. Recorded from Jalalabad, Khost (E), Dilaram (W) 823 m XI/65 (LA).

2. Rana ridibunda Pallas

R. r. ridibunda. North of the Hindu Kush as far south and east as Paghman; also known from Iranian Seistan; up to an elevation of 2 500 m. Recorded from Paghman 2 400 m, Herat 1 800 m VIII/65, Faizabad (NE) VIII/65 (LA), Jalalabad 750 m VII/64, Khost 1 400 m VII/65 (CC).

3. Rana sternosignata Murray

Southeastern Afghanistan, north to Kabul. Elevation between 1 400 and 2 800 m. Recorded from Paghman 2 440 m, Kandahar 1 425 m (LA), 50 km W of Khost 2 700 m (CC), Logar River (E of Kabul) 1 000 m X/50 (LA).

Appendix 9

BIBLIOGRAPHY - ALPHARETICAL

- Abbott, J. Narrative of a Journey from Heraut to Khive. Moscow and St. Petersburgh, 1884 W.H. Allen and Co., London. Vol. I.
- Adams, L. Remarks on the habits and haunts of some of the mammalia found in the various parts of India and the western Himalaya Mountains. Proc. Zool. Soc., London, 26:, 512-531.
- Aellen, V. Contribution à l'étude de la faune d'Afghanistan Chiroptères. Revue Suisse 1959 Zool. 66: 353-386.
- Afghanistan, Cartographic Institute. Afghanistan, physical and political map, 1:1 300 000. 1968 Kabul.
- Afghanistan, Central Statistics Office. Provisional Cazetteer for Afghanistan, Vol. 1, 1975 Afghan Demographic Studies, Prime Ministry, Kabul.
- Afghanistan, Central Statistics Office. Afghan agriculture in figures. Kabul. 168 p. 1978
- Afghanistan, Civil Aviation and Tourism Authority, General Meteorological Directorate.

 Monthly climatological data. Ministry of Transport, General Meteorological

 Directorate, Computer Section. Kabul.
- Afghanistan, Ministry of Planning. Basic directive and indicative figures of the five year 1979 economic and social development plan of the Democratic Republic of Afghanistan during the years 1358-1362 (1979-1984). Kabul. 91 p.
- Afghanistan, topographic maps. 1:100 000 complete; 1:250 000 northern part complete, central, south incomplete; 1:50 000 central and southern part complete.
- Agachanjanc, O.E. Botanical-geographical survey of Afghanistan. Izv. Akad. Nauk. tadzhik. 1958 SSR; Otd. est Nauk. 1: 25 (in Russian).
- Agachanjanc, O.E. Notes on botanical-geographical districts of Tadzhikistan and NE
 1960 Afghanistan. Izv. Akad. Nauk tadzhik. SSR; Otd. s.-sh. bilog. 1,2 (in Russian).
- Agachanjano, O.E. Vegetation of ME Afghanistan. Sb. tr. tadzhik. fil. geogr. obech. SSR. 1961 2 (in Russian).
- Agachanjano, O.E. Botanical-geographical observations in Afghanistan. Bot. Zhurn. 1964 49: 150-155 (in Russian).
- Ahlemann, H. Markhor-Jagd in Muristan. Wild und Hund. 73: 172-177 (Hunting Markhor in 1970 Nuristan).
- Aitchison, C.U. (comp) A collection of treaties, engagements and Sanads relating to India 1933 and neighbouring countries. Government of India, Calcutta, 5th edition. Vol. 13 Persia and Afghanistan. pp. 547.
- Aitchison, J.E.T. The vegetation of Kurram and Hariab valleys. Ind. For. 5: 179-188. 1879

Aitchison, J.E.T. Flora of Afghanistan. J. Linn. Soc.

Part 1, 122 p.

7.

1880 Vo. 16. Part 2, 93 p. Vol. 19.

Aitchison, J.E.T. On the Flora of Kurram Valley. J. Linn. Soc. Bot. 18: 1-122; 19: 139-200. 1881/82 1 map, 3 + 29 plates.

Aitchison, J.E.T. The Botany of the Afghan Delimitation Commission. Trans. Proc. Bot. Soc. 1888 Edinb. 2: Ser 3; 1-139.

Aitchison, J.E.T. The Zoology of the Afghan Delimitation Commission. Trans. Linn. Soc. 1989a Lond. pp. 53 - 142.

Aitchison, J.E.T. A summary of the botanical features of the country traversed by the 1889b Afghan Delimitation Commission during 1884/85. Trans. bot. Soc. Edinb. 17: 421-434.

Aitchison, J.E.T. Notes on the products of W. Afghanistan and NE Persia. Trans. Proc. 1890 bot. Soc. Edinb. 18: 1-228.

Akhtar, S.A. Babar the Great on flamingos. J. Bombay Nat. Hist. Soc. 46: 545-547. 1946

Akhtar, S.A. Ab-i-Estada, a breeding place of the flamingo Phoenicopterus ruber roseus 1947 (Pallas) in Afghanistan. J. Bombay Nat. Hist. Soc. 47: 308-314.

Akhtar, S.A. Bird migration and fowling in Afghanistan. J. Bombay Nat. Hist. Soc. 1955 53: 49-53.

Akram, M. Bibliographie de L'Afghanistan. Centre of Documentation University, Paris. 1947 pp. 504.

Akram, N. Blütenpflanzen im Hindukusch. in Frey, W., Zwischen Munjan und Bashgal, 1967 47-50, OSppingen.

Akram, N. L'Ouest afghan (Stude physique). (The Afghan West (A Physical Study)) Afghan 6, 1951 No. 2, 57-62.

Alcook, A. & Finn, F. An account of the Reptilia collected by Dr. A.H. Maynard, Captain
A.H. KoMahon, C.I.E. and the members of the Afghan Baluch Boundary Commission.
J. Asiat. Soc. Beng. 65: 550-566.

Aleem, A. Markhor in Chitral Gol. Pakist. J. For. (April) pp. 117-128.

Ali, S. The wild ass of Kutch. J. Bombay Nat. Hist. Soc. 46(3): 472-477.
1949

Ali, S. The book of Indian birds. Natural History Society, Bombay.

Ali, S. & Ripley, S. Handbook of the birds of India and Pakistan. Oxford University 1968— Press, Bombay. Vol. 1-10.

Appendix 4

MAMMALS OF AFGHANISTAN

For the taxonomical background G.B. Corbett, 1978, 'The mammals of the Palaearctic Region, a taxonomic review', is followed except for the genus Ovis, where Nadler et al., 1973 are followed, who consider O. orientalis and O. ammon as two species.

Most of the information is from publications on collections made by J. Hassinger, J. Niethammer and J. Gaisler. For bats collections are made by J. Gaisler, D. Meyer Oehme and V. Aellen.

ORDER INSECTIVORA

Family Erinaceidae

- 1. Hemiechinus auritus (Gmelin) long-eared hedgehog
 H. a. auritus Common in the marshes and semi-deserts of lowland
 northern, western and southern Afghanistan.
 - H. a. megalotis (Blyth) Central and eastern mountains up to 3 000 m in altitude, common in Ab-i-Estada (JN).
- 2. Paraechinus hypomelas (Brandt) Brandt's hedgehog
 P. h. hypomelas. Uncommon and only below 1 700 m. Recorded from
 Paktia, Khost, Kandahar, Jalalabad (JN).

Family Soricidae

- 1. Crocidura pergrisea Miller
 - C. p. pergrisea. Recorded from Afghanistan (Spitzenberger, 1971).
- 2. Crocidura russula (Hermann) greater white-toothed shrew

 C. r. gueldenstaedti (Pallas). Widespread in mountainous areas along

 streams, between 600-3 000 m elevation, also in Wakhan (Naumann).
- Crocidura suaveolens (Pallas)

 C. s. suaveolens. By Richter, 1970, considered as subspecies of C. russula. Northern and western steppes of Afghanistan, in humid places between 500-2 500 m elevation. Recorded from Kunduz, Mazar-i-Sharif and Herat (Richter).
- 4. Crocidura zarudnyi Ognev East-Persian white-toothed shrew
 - C. z. zarudnyi. Southern lowlands, Iranian Seistan.
 - C. z. streetorum Hassinger. Mountainous central and northeast Afghanistan up to 2 000 m elevation, widely distributed (JH).

FAO LIBRARY FICHE AN: 8219795

FO:DP/AFG/78/007 Technical Report, Vol. II

NATIONAL PARKS AND WILDLIFE MANAGEMENT

AFGHANISTAN

A CONTRIBUTION TO A CONSERVATION STRATEGY Volume II: APPENDIXES





UNITED NATIONS DEVELOPMENT PROGRAMME

FOOD AND AGRICULTURE ORGANIZATION
OF THE UNITED NATIONS ROME, 1981

NATIONAL PARKS AND WILDLIFE MANAGEMENT

AFGHANISTAN

A CONTRIBUTION TO A CONSERVATION STRATEGY

Volume II: APPENDIXES

Report prepared for
the Government of Afghanistan
by
the Food and Agriculture Organization of the United Nations
acting as executing agency for
the United Nations Development Programme

based on the work of

J.A. Sayer
Team Leader/Ecologist
and
A.P.M. van der Zon
National Park Planner

UNITED NATIONS DEVELOPMENT PROGRAMME
FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS

Rome, 1981

TABLE OF CONTENTS

Volume I: TEXT	Page
PREFACE	
ACTION PLAN - CONCLUCTORS AND	ix
ACTION PLAN - CONCLUSIONS AND RECOMMENDATIONS	1
1. SOCIAL AND ECONOMIC BACKGROUND	9
1.1 The country	•
1.2 The people	9
1.2.1 Origins	_
1.2.2 Demography	9 11
1.3 The economy	:
1.3.1 Introduction	11
1.3.2 Agriculture	11
1.3.3 Livestock	14
1.4 Energy	14
	17
2. GEOGRAPHICAL AND BIOLOGICAL BACKGROUND	20
2.1 Physical environment	20
2.1.1 Climate	- -
2.1.2 Hydrology	20
2.1.3 Geology	22 22
2.1.4 Soils	24
2.2 Biological environment	27
2.2.1 Biogeographical divisions of Afghanistan	
vegetation	27
2.2.3 Fauna	27 32
2.3 Ecological history	32
and activation in the control of the	36
2.3.1 Prehistoric times	
2.3.2 Improvement of irrigation and agricult	36
	36
Tupact of ungilarag	37
2.3.5 Present forest use	37
2.3.6 Original climax vegetation	38 39
3. CONSERVATION NEEDS AND PRIORITIES	
3.1 Farmland	40
2 1 1	40
3.1.1 Traditional farming 3.1.2 Large-scale irrigation	40
3.1.2 Large-scale irrigation projects	40

	Page
3.2 Rangelands	41
3.2.1 Present grazing policy	
3.2.2 Rangeland management	42
3.2.3 Strategy of mortality control	42
3.2.4 Artificial water supplies	43
	44
3.3 Forests	44
3.3.1 Kunar	77
3.3.2 Paktia	45
3.3.3 Nangahar	47
3.3.4 Kabul Greenbelt	47
3.3.5 Pistacchio forests	48
3.3.6 Fuelwood	48
	48
3.4 Wetlands	
	50
3.4.1 River courses	5.0
3.4.2 Lakes and marshes	50
3.4.3 Artificial lakes	52
3.5 Wildlife conservation	54
order conservation	55
3.5.1 Mammals	
3.5.2 Birds	56
3.5.3 Reptiles	63
3.5.4 Hunting	65
-	65
3.6 Conservation of genetic resources	67
4. INSTITUTIONAL AND LEGISLATIVE FRAMEWORK	
	69
4.1 Introduction	60
4.2 Establishment of an environmental unit	69
TO DUVILUMENTAL Project	69
4.4 The Department of Forests and Range 4.5 Legislation	70
4.5 Legislation	70
4.5.1.0	71
4.5.1 Creation of a system of protected areas	71
	71
4.5.3 Other provisions	73
4.6 Training	74
4.7 International problems and many transfer	74
4.8 Education and public awareness	75
	77
4.8.1 Schools	
4.8.2 Educational broaders:	77
· · · · · · · · · · · · · · · · · · ·	77
4.8.4 Conservation education centre	77
·	77
4.9 Tourism	
	78

5. INVENTORY OF SITES, ACTION PLAN	Page
	80
 5.1 Selection of natural sites, criteria 5.2 Natural sites and action plans 	80 80
5.2.1 Big Pamir	00
5.2.2 Ab-i-Estada	80
5.2.3 Ajar Valley	83
5.2.4 Bande Amir	85
5.2.5 Hamun-i-Puzak	89
5.2.6 Nuristan	91
5.2.7 Kole Hashmat Khan	93 94
5.2.8 Dashte Nawar	97
5.2.9 Darqad and Imam Sahib	99
5.2.10 Registan desert	101
5.2.11 Northwest Afghanistan	103
SELECTED REFERENCES	
	105
LIST OF TABLES	
1. Population estimates for Afghanistan, 1975-2025	
2. Distribution of the labour force, 1975-76	13
interchandise exports	13
4. Categories of land	15
5. Average crop area, 1971/72-1976/77	15
o. Estimated livestock populations of Africa.	16
a reduction of electricity, coal natural are a	16 18
	18
9. Forest types and areas in Afghanistan	45
LIST OF MAPS	
1. Topography	0
2. Administrative areas and communications	8 10
3. The distribution of the population	12
	21
5. Rainfall map and diagrams 6. Soil map	23
7. Biogeographical provinces	. 26
8. The natural vegetation	28
9. Forest cover	30
10. Rivers and wetlands	46
ll. Mammal distribution map	51
12. Mammal distribution map	58
13. Mammal distribution map	60
14. Priority conservation areas	62
13. Pamir and Wakhan Corridor and Big Pamir Wildlife Persons	79 81
	84
1. Mai valley Wildlife Reserve	86
IMACIONAL PARK	88
cook give Hamili-1-28DEL) NAG	90
20. Distribution of markhor and Nuristan National Park 21. Kole Hashmat Khan	92
22. Dashte Nawar Flamingo and Waterfowl Sanctuary	95
23. Dailyad and Imam Sahib	98
24. Registan Desert	100
25. Northwest Afghanistan	102
	104

v

		Page
	Volume II: APPENDIXES	
Appendix 1 CONVENTIO SPECIES	N ON INTERNATIONAL TRADE IN ENDANGERED OF WILD FAUNA AND FLORA (CITES)	
	OF LOCATIONS IN AFGHANISTAN	1
		3
Appendix 3 LIST OF OR	BSERVERS	. 8
Appendix 4 MAMMALS OF	FAFGHANISTAN	
Appendix 5 CHECKLIST	OF THE BIRDS OF AFGHANISTAN	9
		23
Appendix 6 BIRDS OF A	FCHANISTAN	33
Appendix 7 REPTILES O	F AFGHANISTAN	
Appendix 8 AMPHIBIANS	OF ADDITION	88
		97
Appendix 9 BIBLIOGRAPH	HY - ALPHABETICAL	20
Appendix 10 BIBLIOGRAPH	Y - SIRIECTS	98
	00000010	141